


Quick Reference Guide

HEDIS® MY 2026



- Meridian Medicaid
- Wellcare Meridian Dual Align (HMO D-SNP)
- YouthCare (Medicaid)
- Ambetter (Marketplace)
- Wellcare (Medicare)
-  Electronic Clinical Data Systems (ECDS)
- HEDIS measure



For more information, visit www.ncqa.org

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA). The HEDIS® measures and specifications were developed by and are owned by NCQA. NCQA holds a copyright in these materials and may rescind or alter these materials at any time. Users of the HEDIS® measures and specifications shall not have the right to alter, enhance, or otherwise modify the HEDIS® measures and specifications, and shall not disassemble, recompile, or reverse engineer the HEDIS® measures and specifications. Anyone desiring to use or reproduce the materials, subject to licensed user restrictions, without modification for an internal non-commercial purpose may do so without obtaining any approval from NCQA. Use of the Rules for Allowable Adjustments of HEDIS® to make permitted adjustments of the materials does not constitute a modification. All other uses, including a commercial use, or any external reproduction, distribution, or publication must be approved by NCQA and are subject to a license at the discretion of NCQA.

HEDIS® MY 2026 Quick Reference Guide

Updated to reflect NCQA HEDIS® Measurement Year (MY) 2026 Technical Specifications

We strive to provide quality healthcare to our membership as measured through HEDIS quality metrics. We created the HEDIS® MY 2026 Quick Reference Guide to help you increase your practice's HEDIS rates and to address care opportunities for your patients. Please always follow state and/or CMS billing guidance and ensure the HEDIS codes are covered prior to submission.

What is HEDIS®?

HEDIS® (Healthcare Effectiveness Data and Information Set) is a set of standardized performance measures developed by the National Committee for Quality Assurance (NCQA) to objectively measure, report, and compare quality across health plans. NCQA develops HEDIS measures through a committee represented by purchasers, consumers, health plans, healthcare providers, and policymakers.

What Are the Scores Used For?

As state and federal governments move toward more quality-driven healthcare, HEDIS rates are becoming more important for both health plans and individual providers. State purchasers of healthcare use aggregated HEDIS rates to evaluate health insurance companies' efforts to improve preventive health outreach for patients.

Physician-specific scores are also used to measure your practice's preventive care efforts. Your practice's HEDIS score determines your rates for physician incentive programs that pay you an increased premium — for example, Pay for Performance or Quality Bonus Funds.

How Are Rates Calculated?

HEDIS rates can be calculated in two ways: administrative data or hybrid data. Administrative data includes claim or encounter data and electronic clinical data systems (ECDS) data. ECDS uses data directly from EHRs, claims, and registries, which reduces manual burden and increases efficiency. Hybrid data consists of both administrative data and a sample of medical record data. Hybrid data requires review of a random sample of member medical records to abstract data for services rendered but that were not reported to the health plan through claims/encounter data. Accurate and timely claim/encounter data can reduce the need for medical record review. If services are not billed, not billed accurately, or not shared through electronic data feeds, they are not included in the calculation.

How Can I Improve My HEDIS® Scores?

- Submit claim/encounter data for each and every service rendered
- Make sure that chart documentation reflects all services billed
- Bill (or report by encounter submission) for all delivered services, regardless of contract status
- Ensure that all claim/encounter data is submitted in an accurate and timely manner
- Include CPT II codes to provide additional details and reduce medical record requests
- Implement a supplemental data feed to capture relevant HEDIS data, such as test results not obtained via claims or services rendered before the member's enrollment, for more accurate reporting

How Does Meridian's Quality Improvement Team Support Providers?

Meridian's Quality Improvement (QI) Program aims to continuously improve the delivery of healthcare services in a low-resource environment to enhance the overall health status of members. Our team partners with providers to fulfill this objective and ensure members receive high-quality, medically appropriate, and cost-effective healthcare. To support providers, we also publish this guide, which contains descriptions of HEDIS measures and coding requirements for capturing performance.

Go to the [Meridian Quality Improvement Resources section](#) of this guide for essential information about additional QI resources and expectations for providers, such as:

- Overview of the Consumer Assessment of Healthcare Providers and Systems (CAHPS®) survey
- CAHPS® Outpatient Mental Health Survey Information
- Health Outcomes Survey (HOS)
- Critical Incidents (CI) reporting
- Cultural Competence training and tips
- Social Determinants of Health (SDOH) screening
- Quality Education Webinars
- Access and Availability Standards
- Caring for Individuals with Intellectual and Developmental Disabilities (IDD)
- Early Periodic Screening, Diagnosis, and Treatment (EPSDT) resources

Contact Information for Provider Services

- ILmeridian.com 866-606-3700
- wellcare.ILmeridian.com (D-SNP) 1-844-536-2175
- ILYouthCare.com 844-289-2264
- ambetterofillinois.com 1-855-745-5507
- wellcare.com/en/Illinois (Medicare Advantage) 1-855-538-0454
(TTY 711)

 For more information, visit www.ncqa.org

Contents

- 2 What is HEDIS®?
- 2 What Are the Scores Used For?
- 3 How Are Rates Calculated?
- 3 How Can I Improve My HEDIS® Scores?
- 3 How Does Meridian's Quality Improvement Team Support Providers?
- 4 Contact Information for Provider Services
- 5 Contents

8 Partnering with the Health Plan on HEDIS

- 8 Introduction
- 9 Provider Incentive Programs ● ● ●
- 11 Medication Safety Measures ● ● ● ● ●
- 11 Electronic Clinical Data Systems (ECDS) ● ● ● ● ● ● ●
- 14 Supplemental Data ● ● ● ●

15 Adult Health

15	Adults' Access to Preventive/Ambulatory Health Services (AAP)	● ● ●
16	Advanced Care Planning (ACP)	● ●
16	Adult Immunization Status (AIS-E)	● ● ● ● ● ●
19	Appropriate Treatment for Upper Respiratory Infection (URI)	● ● ● ●
20	Glycemic Status Assessment for Patients with Diabetes (GSD)	● ● ● ●
20	Blood Pressure Control for Patients with Diabetes (BPD) & Blood Pressure Control for Patients with Diabetes (BPD-E)*	● ● ● ● ● ● ● ●
21	Eye Exam for Patients with Diabetes (EED)	● ● ● ●
22	Kidney Health Evaluation for Patients with Diabetes (KED)	● ● ● ●
22	Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)	●
23	Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)	●
24	Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)	●
24	Colorectal Cancer Screening (COL-E)	● ● ● ● ● ●
25	Care for Older Adults (COA)	●
26	Controlling High Blood Pressure (CBP) & Blood Pressure Control for Patients with Hypertension (BPC-E)*	● ● ● ● ● ● ● ●
27	Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions (FMC)	● ●
28	Persistence of Beta-Blocker Treatment After a Heart Attack (PBH)	● ● ●
29	Pharmacotherapy Management of COPD Exacerbation (PCE)	● ● ●
29	Plan All Cause Readmissions (PCR)	● ● ● ●
30	Transitions of Care (TRC)	● ●
31	Use of Imaging Studies for Low Back Pain (LBP)	● ● ● ●

33 Behavioral Health

33	Adherence to Antipsychotic Medications for People with Schizophrenia (SAA)	● ● ● ●
34	Follow-Up After Emergency Department Visit for Mental Illness (FUM)	● ● ● ●
37	Follow-Up After Emergency Department Visit for Substance Use (FUA)	● ● ● ●
47	Follow-Up After High Intensity Care for Substance Disorder (FUI)	● ● ●
50	Follow-Up After Hospitalization for Mental Illness (FUH)	● ● ● ● ●

54	Initiation and Engagement of Substance Use Disorder (IET)	● ● ● ● ●
57	Pharmacotherapy for Opioid Use Disorder (POD)	● ● ● ●
58	Unhealthy Alcohol Use Screening and Follow-Up (ASF-E)	● ● ● ● ●
59	Screening for Depression and Follow-Up Plan (CDF-AD)	●
59	Screening for Depression and Follow-Up Plan (CDF-CH)	●
60	Depression Screening and Follow-Up for Adolescents and Adults (DSF-E)	● ● ● ●
60	Depression Remission or Response for Adolescents and Adults (DRR-E)	● ● ● ●
61	Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E)	● ● ● ●
62	Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)	● ●
63	General Health	
63	Appropriate Testing for Pharyngitis (CWP)	● ● ● ●
63	Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis (AAB)	● ● ● ● ●
65	Follow-Up After Acute and Urgent Care Visits for Asthma (AAF-E)	● ● ●
72	Tobacco Use Screening and Cessation Intervention (TSC-E)	● ● ● ● ●
73	Pediatric Health	
73	Childhood Immunization Status (CIS-E)	● ● ● ●
78	Follow-Up Care for Children Prescribed ADHD Medication (ADD-E)	● ● ●
79	Immunizations for Adolescents (IMA-E)	● ● ● ●
81	Lead Screening in Children (LSC-E)	● ●
82	Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E)	● ● ●
84	Oral Evaluation, Dental Services (OED)	● ● ●
84	Topical Fluoride for Children (TFC)	● ●
85	Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC)	● ● ●
86	Well-Child and Adolescent Well-Care Visits (W30)	● ● ● /WCV ● ● ●
87	Pharmacy Measures	
87	Use of High-Risk Medications in Older Adults (DAE)	● ●
91	Adherence to Cholesterol Medications (STAT)	● ●
92	Adherence to Diabetes Medications (DIAB)	● ●
93	Adherence to Hypertensive Medications (RASA)	● ●
94	Concurrent Use of Opioids and Benzodiazepines (COB)	● ● ● ●
95	Polypharmacy: Use of Multiple Anticholinergic Medications in Older Adults (POLY-ACH)	● ●
97	Statin Therapy for Patients with Cardiovascular Disease (SPC-E)	● ● ● ●
98	Statin Therapy for Patients with Diabetes (SPD-E)	● ● ● ●
99	Women's Health	
99	Breast Cancer Screening (BCS-E)	● ● ● ●
100	Documented Assessment After Mammogram (DBM-E)	● ● ● ● ●
101	Follow-Up After Abnormal Mammogram Assessment (FMA-E)	● ● ● ● ●
103	Cervical Cancer Screening (CCS-E)	● ● ●
104	Chlamydia Screening (CHL)	● ●
105	Osteoporosis Management in Women Who Had a Fracture (OMW)	● ●

- 106 Osteoporosis Screening in Older Women (OSW) ● ●
- 106 Prenatal and Postpartum Care (PPC) ● ●
- 108 Prenatal Immunization Status (PRS-E) ● ● ●
- 109 Prenatal Depression Screening and Follow-Up (PND-E) ● ● ●
- 109 Postpartum Depression Screening and Follow-Up (PDS-E) ● ● ●
- 110 Consumer Assessment of Healthcare Providers and Systems (CAHPS®) ● ● ● ● ● ●

110 Meridian Quality Improvement Resources

- 110 Consumer Assessment of Healthcare Providers and Systems (CAHPS®) ● ● ● ● ● ●
- 114 CAHPS® Outpatient Mental Health Survey Information ● ● ● ● ● ●
- 115 Health Outcomes Survey (HOS) ● ●
- 119 Critical Incidents (CI) ● ●
- 120 Cultural Competence ● ● ●
- 121 Social Determinants of Health (SDOH) ● ● ● ● ● ●
- 122 Quality Education Webinars (QEW) ● ● ● ● ● ●
- 123 Access & Availability ● ● ● ●
- 132 Caring for and Communicating with Individuals with Intellectual and Developmental Disabilities (IDD) ● ● ● ● ● ●
- 134 Early Periodic Screening, Diagnosis, and Treatment (EPSDT) ● ●

138 Appendix

- 138 Antidepressant Medication List ● ● ● ●
- 139 Approved Depression Screening Instruments, Codes, and Positive Findings ● ● ● ● ●
- 140 Follow-Up Visit ● ● ● ●
- 142 Depression or Other Behavioral Health Condition ● ● ● ●
- 147 Denominator Exclusions ● ● ● ● ●
- 157 Point of Service Code Definitions ● ● ● ● ●

Partnering with the Health Plan on HEDIS



Introduction

This guide contains information about the quality measures for the following Meridian lines of business:

- **Meridian Medicaid Plan (Medicaid)**

The Meridian Medicaid Plan in Illinois provides government-sponsored managed care services to families, children, seniors and individuals with complex medical needs through Medicaid across the state.

- **Wellcare Meridian Dual Align (HMO D-SNP)**

Wellcare By Meridian combines Medicaid and Medicare Advantage benefits into one integrated plan.

- **YouthCare (Medicaid)**

The YouthCare HealthChoice Illinois program is a specialized healthcare program built cooperatively with parents and other stakeholders to improve access, continuity of care, and healthcare outcomes for Department of Children and Family Services (DCFS) youth in care and former youth in care.

- **Ambetter of Illinois (Marketplace)**

Ambetter offers quality and affordable health insurance in Illinois that fits various needs and budgets.

- **Wellcare (Medicare)**

Medicare Advantage (MA) plans that offer Medicare Part A, Part B, and Part D benefits.

Provider Incentive Programs ● ● ●

Partnership for Quality (P4Q) measures are on the provider portal. Visit our [Provider Portal](#) to view the recent P4Q measures available to our Meridian provider partners.

For additional information about provider incentive programs, please contact your organization's assigned Provider Engagement Administrator or representative. If your practice doesn't have an assigned representative, please complete the [Provider Engagement intake form](#) or consult our [Provider Engagement page](#) for additional contact options.

Provider incentive programs apply to non-risk contracted providers only.

HEDIS MY 2026 Summary of Changes ● ● ● ●

The following changes are effective for 2026.

This guide has been updated with information from the release of the HEDIS 2026 Volume 2 Technical Specifications by NCQA and is subject to change.

Retired Measures:

NCQA is retiring the following two measures in MY 2026.

- Asthma Medication Ratio (AMR)
- Medical Assistance with Smoking and Tobacco Use Cessation (MSC)

New Electronic Clinical Data Systems (ECDS) Measures

As NCQA advances its multiyear transition to digital reporting, the following measures can be reported using the ECDS method beginning in MY 2026.

- Blood Pressure Control for Patients with Diabetes (BPD-E)
- Follow-Up After Acute and Urgent Care Visits for Asthma (AAF-E)*
- Tobacco Use Screening and Cessation Intervention (TSC-E)*

** These new measures replace the AMR and MSC measures noted in the "Retired Measures" section above.*

Measures that are now ECDS-only

For three common HEDIS measures below, only the ECDS reporting method can be used. There will no longer be a hybrid sample for the LSC, SPC, and SPD measures beginning in MY 2026.

- Lead Screening in Children (LSC-E)
- Statin Therapy for Patients with Cardiovascular Disease (SPC-E)
- Statin Therapy for Patients with Diabetes (SPD-E)

Revised Measures

Please refer to each measure's listing within the guide for the latest measure description.

- **Adult Immunization Status (AIS-E)** now has a COVID-19 indicator for adults 65 years of age and older.
- **Depression Screening and Follow-Up for Adolescents and Adults (DSF-E); Prenatal Depression Screening and Follow-Up (PND-E); and Postpartum Depression Screening and Follow-Up (PDS-E)** now allow the PROMIS® Emotional Distress instrument as an approved depression screening instrument for adults 18 years of age and older.
- **Follow-Up After High Intensity Care for Substance Use Disorder (FUI)** has been revised in two critical ways:
 - Modified numerators allow substance use disorder diagnoses in any position on the claim.
 - An expanded numerator now includes peer support as an appropriate follow-up visit.
- **Social Needs Screening and Intervention (SNS-E)** added provider assessments to the screening numerators.

Terminology Changes for 2026

NCQA is adopting new terms for consistency with FHIR® data standards and to reflect how measures are used.

Previous Terminology (MY 2025)	New Terminology (MY 2026)
Measurement Year	Measurement Period
Members	Persons
Eligible Population	Initial Population
Required Exclusions	Denominator Exclusions

Medication Safety Measures



In MY 2025, Meridian began emphasizing three additional measures related to medication safety. Find these in the [Pharmacy Measures section](#) of the guide.

- Use of High-Risk Medications in Older Adults (DAE)

Pharmacy Quality Alliance (PQA) Measures

The guide includes two pharmacy-related measures established and updated by the Pharmacy Quality Alliance (PQA), an independent nonprofit national organization dedicated to improving medication safety, adherence, and appropriate use. Visit [PQAalliance.org](#) for more information and resources.

- Concurrent Use of Opioids and Benzodiazepines (COB)
- Polypharmacy: Use of Multiple Anticholinergic Medications in Older Adults (POLY-ACH)

Electronic Clinical Data Systems (ECDS)



Tip: Look for the  icon to quickly identify HEDIS measures that are only reported using the ECDS method.

ECDS (Electronic Clinical Data Systems) are the network of data containing a patient's personal health information and records of their experiences within the healthcare system. They may also support other care-related activities directly or indirectly, including evidence-based decision support, quality management, and outcome reporting. Data in these systems are structured such that automated quality measurement queries can be consistently and reliably executed, providing results quickly and efficiently to the team responsible for patient care.

Establishing an enterprise network of interoperable electronic data systems will foster a patient-centered, team-based approach to improving healthcare quality and communication across healthcare service providers.

ECDS reporting is part of NCQA's larger strategy to enable a Digital Quality System and is aligned with the industry's move towards greater interoperability of health information. Visit [www.ncqa.org/ecds](#) for more information and frequently asked questions about ECDS reporting.

The goal is to promote the integration of clinical information by automatically transferring data for gap closure. ECDS measures allow for plans to view quality care prospectively as opposed to reviewing quality care retrospectively.

ECDS data sources include:

- Enrollment
- Claims
- Encounters
- EHRs
- HIEs
- Registries
- Case Management

Tips for successful gap closure for all measures include:

- Establish electronic data transfers, such as supplemental data files
 - Refer to Supplemental Data source section
- Include CPT II codes on claims
- Ensure CPT II, LOINC, and SNOMED codes are all linked in your provider EMR system
 - LOINC and SNOMED value sets are transferred electronically, not on claims
- Develop and execute a data strategy prior to HEDIS transitioning to ECDS only

ECDS reporting highlights:

The following measures have transitioned to ECDS-only reporting for MY 2026:

- Lead Screening In Children (LSC-E)
- Statin Therapy for Patients with Cardiovascular Disease (SPC-E)
- Statin Therapy for Patients with Diabetes (SPD-E)

Other ECDS measures developed that are currently being measured and reported include:

- Follow-Up After Acute and Urgent Care Visits for Asthma (AAF-E)
- Follow-Up Care for Children Prescribed ADHD Medication (ADD-E)
- Adult Immunization Status (AIS-E)
- Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E)
- Unhealthy Alcohol Use Screening and Follow-Up (ASF-E)
- Breast Cancer Screening (BCS-E)
- Blood Pressure Control for Patients With Hypertension (BPC-E)
- Blood Pressure Control for Patients With Diabetes (BPD-E)
- Cervical Cancer Screening (CCS-E)
- Childhood Immunization Status (CIS-E)
- Colorectal Cancer Screening (COL-E)
- Documented Assessment After Mammogram (DBM-E)
- Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E)

- Depression Remission or Response for Adolescents and Adults (DRR-E)
- Depression Screening and Follow-Up for Adolescents and Adults (DSF-E)
- Follow-Up After Abnormal Mammogram Assessment (FMA-E)
- Immunizations for Adolescents (IMA-E)
- Postpartum Depression Screening and Follow-Up (PDS-E)
- Prenatal Depression Screening and Follow-Up (PND-E)
- Prenatal Immunization Status (PRS-E)
- Social Need Screening and Intervention (SNS-E)
- Tobacco Use Screening and Cessation Intervention (TSC-E)

Visit www.ncqa.org/ecds for more information and FAQs about ECDS reporting.

ECDS Transition Timeline for Hybrid Measures

	MY2025	MY2026	MY2027	MY2028	MY2029
Lead Screening in Children (LSC)		ECDS Only			
Weight Assessment and Counseling for Nutrition and Physical Activity in Children and Adolescents (WCC)			Admin Only		
Prenatal and Postpartum Care (PPC)				Admin Only	
Controlling High Blood Pressure (CBP)	+ECDS				
Blood Pressure Control for Patients with Diabetes (BPD)		+ECDS		ECDS Only	
Glycemic Status Assessment for Patients with Diabetes (GSD)			+ECDS		ECDS Only
Transition of Care (TRC)			+ECDS		ECDS Only
Care for Older Adults (COA)			+ECDS		ECDS Only

Supplemental Data ●●●●●

Supplemental data is data that is captured for HEDIS care gap closure, and used in rate calculation, but is not submitted through claims processing. It allows for services rendered to be captured for gap closure.

Supplemental Data Benefits

- Improve accuracy by minimizing human intervention and error that can occur through chart reviews and data entry
- Increase efficiency of data retrieval to support real-time intervention with patients for care gap closure
- Cost efficient
- Eliminate the burden on provider staff submitting high volumes of medical records
- Increase provider incentive earnings and improve provider performance
- Captures lab results and blood pressure readings, if CPT II codes are not captured on claims

Supplemental Data Implementation

If you are interested in setting up a supplemental data feed, please contact our HEDIS Operations team at ILHEDISOps@mhplan.com

Please include the following information:

- Main point of contact regarding the data feed
- IP Address
- Tax ID Number(s)
- Product lines you plan to submit data for (Ambetter, Meridian, Wellcare, and/or YouthCare)
- List of individuals to be involved in the implementation communication
 - Please include the person responsible for generating the file and submitting the data

Adult Health



Adults' Access to Preventive/Ambulatory Health Services (AAP) ●●●

The AAP measure evaluates the percentage of persons 20 years of age and older who had an ambulatory or preventive care visit. Services that count include outpatient evaluation and management (E&M) visits, consultations, assisted living/home care oversight, preventive medicine, and counseling.

AAP Measure Codes

Description	CPT*	HCPCS*	ICD-10-CM*
Ambulatory Visits	92002, 92004, 92012, 92014, 98000-98016, 98966-98968, 98970-98972, 98980-98981, 99202-99205, 99211-99215, 99242-99245, 99304-99310, 99315-99316, 99341-99345, 99421, 99341, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99429, 99441-99443, 99457, 99458, 99483	G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2250-G2252, S0620-S0621, T1015	

AAP Measure Codes (continued)

Description	CPT*	HCPCS*	ICD-10-CM*
Reason for Ambulatory Visit			Z00.00, Z00.01, Z00.121, Z00.129, Z00.3, Z00.5, Z00.8, Z02.0-Z02.6, Z02.71, Z02.79, Z02.81-Z02.84, Z02.89, Z02.9, Z76.1, Z76.2

*Codes subject to change

Advanced Care Planning (ACP) ●●

The ACP measure evaluates the percentage of adults 66 to 80 years of age with advanced illness, an indication of frailty, or who are receiving palliative care and had advanced care planning, and adults 81 years of age or older who had advanced care planning during the measurement period.

A discussion or documentation about preferences for resuscitation, life-sustaining treatment, and end-of-life care.

ACP Measure Codes

Description	Codes*
Advanced Care Planning	CPT: 99483, 99497 CPT-CAT-II: 1123F, 1124F, 1157F, 1158F HCPCS: S0257 ICD-10: Z66

*Codes subject to change

Adult Immunization Status (AIS-E) ●●●●

AIS-E is an Electronic Clinical Data Systems (ECDS) measure.

AIS-E evaluates the percentage of adults 19 years of age and older who are up to date on recommended routine vaccines for influenza, tetanus and diphtheria (Td) or tetanus, diphtheria and acellular pertussis (Tdap), zoster, pneumococcal, hepatitis B, and coronavirus disease 2019 (COVID-19).

Influenza:

- Persons who received an influenza vaccine on or between July 1 of the year prior to the measurement period and June 30 of the measurement period

Td/Tdap:

- Persons who received at least one Td vaccine or one Tdap vaccine between 9 years prior to the start of the measurement period and the end of the measurement period

Zoster:

- Persons who received two doses of the herpes zoster recombinant vaccine at least 28 days apart, any time on or after the patient's 50th birthday and before or during the measurement period

Pneumococcal:

- Persons who were administered at least one dose of an adult pneumococcal vaccine on or after their 19th birthday and before or during the measurement period

Hepatitis B:

- Persons who received at least two doses of the adult hepatitis B vaccine on or after their 19th birthday, administered at least 28 days apart

COVID-19:

- Persons who were administered at least one dose of a COVID-19 vaccine that occurred **both** on or between July 1 of the year prior to the measurement period through June 30 of the measurement period **and** on or after their 65th birthday

AIS-E Measure Codes

Description	Codes*
Adult Influenza Immunization	CVX: 88, 135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205, 320
Adult Influenza Vaccine Procedure	CPT: 90630, 90653, 90654, 90656, 90658, 90661, 90662, 90673, 90674, 90682, 90686, 90688, 90689, 90694, 90756 SNOMED CT: 86198006
Influenza Virus LAIV Immunization	CVX: 111, 149
Influenza Virus LAIV Vaccine Procedure	CPT: 90660, 90672 SNOMED CT: 787016008
Patients with Anaphylaxis due to Influenza Vaccine	SNOMED CT: 471361000124100
Td Immunization	CVX: 09, 113, 115, 138, 139
Anaphylaxis due to Diphtheria, Tetanus or Pertussis Vaccine	SNOMED CT: 428281000124107, 428291000124105
Encephalitis due to Diphtheria, Tetanus or Pertussis Vaccine	SNOMED CT: 192710009, 192711008, 192712001
Herpes Zoster Recombinant Vaccine Procedure (2 doses)	CVX: 187 CPT: 90750 SNOMED CT: 722215002

AIS-E Measure Codes (continued)

Description	Codes*
Anaphylaxis due to Herpes Zoster Vaccine	SNOMED CT: 471371000124107, 471381000124105
Adult Hepatitis B Vaccine Procedure	CPT: 90739, 90740, 90743, 90744 90746, 90747, 90759
Adult Pneumococcal Immunization	CVX: 33, 109, 133, 152, 215, 216, 327
Adult Pneumococcal Vaccine Procedure	CPT: 90670, 90671, 90677, 90684, 90732 HCPCS: G0009 SNOMED CT: 12866006, 394678003, 871833000, 1119366009, 1119367000, 1119368005, 434751000124102
Patients with Anaphylaxis due to the Pneumococcal Vaccine	SNOMED CT: 471141000124102
Adult COVID-19 Immunization	CVX: 309, 312, 313
Adult COVID-19 Vaccine Procedure	CPT: 91304, 91320, 91322

*Codes subject to change

Appropriate Treatment for Upper Respiratory Infection (URI) ●●●●

The URI measure evaluates the percentage of episodes for persons 3 months of age and older with a diagnosis of upper respiratory infection (URI) that did not result in an antibiotic-dispensing event.

Antibiotic Medications

- Amikacin
- Amoxicillin
- Amoxicillin/Clavulanate
- Ampicillin
- Ampicillin/Sulbactam
- Azithromycin
- Aztreonam
- Cefaclor
- Cefadroxil
- Cefazolin
- Cefdinir
- Cefepime
- Cefixime
- Cefotaxime
- Cefotetan
- Cefoxitin
- Cefpodoxime
- Cefprozil
- Ceftazidime
- Ceftriaxone
- Cefuroxime
- Cephalexin
- Chloramphenicol
- Ciprofloxacin
- Clarithromycin
- Clindamycin
- Dalforpristin/Quinupristin
- Daptomycin
- Dicloxacillin
- Doxycycline
- Erythromycin
- Fosfomycin
- Gemifloxacin
- Gentamicin
- Levofloxacin
- Lincomycin
- Linezolid
- Metronidazole
- Minocycline
- Moxifloxacin
- Nafcillin
- Nitrofurantoin
- Nitrofurantoin, Macrocrystals/
Nitrofurantoin, Monohydrate
- Ofloxacin
- Oxacillin
- Penicillin G Benzathine
- Penicillin G Benzathine/
Penicillin G Procaine
- Penicillin G Potassium
- Penicillin G Procaine
- Penicillin G Sodium
- Penicillin V Potassium
- Piperacillin/Tazobactam
- Rifampin
- Streptomycin
- Sulfadiazine
- Sulfamethoxazole/Trimethoprim
- Tetracycline
- Tobramycin
- Trimethoprim
- Vancomycin

Note: Comprehensive Diabetes Care (CDC) was replaced by the following three measures in MY 2024: Glycemic Status Assessment for Patients with Diabetes (GSD), Blood Pressure Control for Patients with Diabetes (BPD), and Eye Exam for Patients with Diabetes (EED).

Glycemic Status Assessment for Patients with Diabetes (GSD)

The GSD measure evaluates persons 18–75 years of age with diabetes (Type 1 and 2) whose most recent glycemic status (hemoglobin A1C [HbA1c] or glucose management indicator [GMI]) was at the following levels during the measurement period.

Two rates are reported:

- Glycemic Status <8.0%
- Glycemic Status >9.0% (lower rate indicates better performance)

GSD Measure Codes

Description	Codes*
HbA1c Lab Test (used to identify the last assessment of the measurement period, but does not close the care gap without a result)	CPT: 83036, 83037 LOINC: 4548-4, 17855-8, 17856-6, 4549-2, 96595-48
HbA1c Level Greater Than 9.0	CPT-CAT-II: 3046F
HbA1c Level Greater Than or Equal to 7.0 and Less Than 8.0	CPT-CAT-II: 3051F
HbA1c Level Greater Than or Equal to 8.0 and Less Than or Equal to 9.0	CPT-CAT-II: 3052F
HbA1c Level Less Than 7.0	CPT-CAT-II: 3044F
Glucose management indicator (GMI)	LOINC: 97506-0

*Codes subject to change

Blood Pressure Control for Patients with Diabetes (BPD) & Blood Pressure Control for Patients with Diabetes (BPD-E)*

* Note that voluntary ECDS reporting for the BPD-E measure is available beginning in MY 2026.

The BPD measure evaluates the percentage of persons 18 to 75 years of age with diabetes (Types 1 and 2) whose last blood pressure (BP) reading of the measurement period was adequately controlled (<140/90 mm Hg).

- Consider taking two readings at each office visit. Sometimes a second blood pressure reading is lower.
- Do not round blood pressure; always use exact numbers
- Include CPT-CAT-II codes on claims to ensure the blood pressure reading is received by the health plan

BPD Measure Codes

Description	Codes*
Diastolic 80-89	CPT-CAT-II: 3079F
Diastolic Greater Than or Equal to 90	CPT-CAT-II: 3080F
Diastolic Less Than 80	CPT-CAT-II: 3078F
Systolic Greater Than or Equal to 140	CPT-CAT-II: 3077F
Systolic Less Than 130	CPT-CAT-II: 3074F
Systolic 130-139	CPT-CAT-II: 3075F

*Codes subject to change

Eye Exam for Patients with Diabetes (EED) ●●●●

The EED measure evaluates the percentage of persons 18 to 75 years of age with diabetes (Types 1 and 2) who had a retinal eye exam.

- Include CPT-CAT-II codes on claims to ensure the eye exam result is received by the health plan

EED Measure Codes

Description	Codes*
Unilateral Eye Enucleation	CPT: 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114
Automated Eye Exam	CPT: 92229
Eye Exam with Retinopathy	CPT-CAT-II: 2022F, 2024F, 2026F
Eye Exam Without Retinopathy	CPT-CAT-II: 2023F, 2025F, 2033F
Diabetes Mellitus without Complications (Also requires Diabetic Retinal Screening code to close care gap)	ICD-10: E10.9, E11.9, E13.9
Retinal Eye Exams	CPT: 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92137, 92201, 92202, 92230, 92235, 92250, 99203, 99204, 99205, 99213-99215, 99242-99245 HCPCS: S3000, S0621, S0620
Retinal imaging	CPT: 92227-92228
Diabetic Retinal Screening negative in the prior year	CPT-CAT-II: 3072F

*Codes subject to change

Kidney Health Evaluation for Patients with Diabetes (KED) ●●●●

The KED measure evaluates the percentage of persons 18 to 85 years of age with diabetes (Type 1 and Type 2) who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) and a urine albumin-creatinine ratio (uACR), during the measurement period.

KED Measure Codes

Description	Codes*
Estimated Glomerular Filtration Rate (eGFR)	CPT: 80047, 80048, 80050, 80053, 80069, 82565
Urine Albumin-Creatinine Ratio (uACR)	CPT: 82043, 82570

*Codes subject to change

Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD) ●

The SMD measure evaluates the percentage of persons 18 to 64 years of age with schizophrenia or schizoaffective disorder and diabetes who had both an LDL-C test and an HbA1c test.

SMD Measure Codes

Description	Codes*
HbA1C Lab Tests	CPT: 83036, 83037
Most recent hemoglobin A1c (HbA1c) level less than 7.0% (DM)	CPT-CAT-II: 3044F
Most recent hemoglobin A1c level greater than 9.0% (DM)	CPT-CAT-II: 3046F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 7.0% and less than 8.0% (DM)	CPT-CAT-II: 3051F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 8.0% and less than or equal to 9.0% (DM)	CPT-CAT-II: 3052F
LDL-C Lab Tests	CPT: 80061, 83700, 83701, 83704, 83721
Most recent LDL-C less than 100 mg/dL (CAD) (DM)	CPT-CAT-II: 3048F
Most recent LDL-C 100-129 mg/dL (CAD) (DM)	CPT-CAT-II: 3049F
Most recent LDL-C greater than or equal to 130 mg/dL (CAD) (DM)	CPT-CAT-II: 3050F

*Codes subject to change

Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD) ●

The SSD measure evaluates the percentage of persons 18 to 64 years of age with schizophrenia, schizoaffective disorder, or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test.

SSD Measure Codes

Description	Codes*
HbA1C Lab Tests	CPT: 83036, 83037
Most recent hemoglobin A1c (HbA1c) level less than 7.0% (DM)	CPT-CAT-II: 3044F
Most recent hemoglobin A1c level greater than 9.0% (DM)	CPT-CAT-II: 3046F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 7.0% and less than 8.0% (DM)	CPT-CAT-II: 3051F
Most recent hemoglobin A1c (HbA1c) level greater than or equal to 8.0% and less than or equal to 9.0% (DM)	CPT-CAT-II: 3052F
Glucose Lab Tests	CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951

*Codes subject to change

Cardiac Rehabilitation (CRE) ● ● ●

The CRE measure evaluates the percentage of persons 18 years of age and older who attended cardiac rehabilitation following a qualifying cardiac event, such as:

- Coronary artery bypass grafting
- Heart or heart/lung transplantation
- Heart valve repair/replacement
- Myocardial infarction
- Percutaneous coronary intervention

It is important that patients have the following schedule of cardiac rehabilitation:

- **Initiation:** At least 2 sessions within 30 days after the event
- **Engagement:**
 - At least 12 sessions within 90 days after the event
 - At least 24 sessions within 180 days after the event
- **Achievement:**
 - At least 36 sessions within 180 days after the event
 - Encourage patients to have annual testing

CRE Measure Codes

Description	CPT Category*	HCPCS*
Cardiac Rehabilitation	93797, 93798	G0422, G0423, S9472

*Codes subject to change

Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) ●

The SMC measure evaluates the percentage of persons 18-64 years of age with schizophrenia or schizoaffective disorder and cardiovascular disease who had an LDL-C test during the measurement period.

SMC Measure Codes

Description	Codes*
LDL-C Lab Test	CPT: 80061, 83700, 83701, 83704, 83721
LDL-C Test Result or Finding	CPT-CAT-II: 3048F, 3049F, 3050F

*Codes subject to change

Colorectal Cancer Screening (COL-E) ● ● ● ● ●

COL-E evaluates the percentage of persons 45 to 75 years of age who had one or more appropriate screenings for colorectal cancer.

Appropriate screenings are defined by one of the following:

- Colonoscopy during the measurement period or the nine years prior to the measurement period
- CT colonography during the measurement period or the four years prior to the measurement period
- Flexible sigmoidoscopy during the measurement period or the four years prior to the measurement period
- FIT-DNA during the measurement period or the two years prior to the measurement period
- FOBT during the measurement period

COL-E Measure Codes

Description	Codes*
Colonoscopy	CPT: 44388-44392, 44394, 44401-44408, 45378-45382, 45384-45386, 45388-45393, 45398 HCPCS: G0105, G0121 SNOMED CT: 8180007, 12350003, 25732003, 73761001, 174158000, 174171002, 174185007, 235150006, 275251008, 302052009, 311774002, 367535003, 426699005, 443998000, 444783004, 446521004, 446745002, 447021001, 709421007, 710293001, 711307001, 773128008, 7731202006, 789778002, 10371000132109, 1209098000, 851000119109, 48021000087103, 4803100087101
CT Colonography	CPT: 74261-74263 LOINC: 60515-4, 72531-7, 79069-1, 79071-7, 79101-2, 82688-3
sDNA FIT Lab Test	CPT: 81528 LOINC: 77353-1, 77354-9 SNOMED CT: 708699002
Flexible Sigmoidoscopy	CPT: 45330-45335, 45337-45338, 45340-45342, 45346-45347, 45349-45350 HCPCS: G0104 SNOMED CT: 44441009, 396226005, 425634007
FOBT Lab Test	CPT: 82270, 82274 HCPCS: G0328 LOINC: 104738-0, 107189-3, 107190-1, 107191-9, 12503-9, 12504-7, 14563-1, 14564-9, 14565-6, 2335-8, 27396-1, 27401-9, 27925-7, 27926-5, 29771-3, 56490-6, 56491-4, 57905-2, 58453-2, 80372-6
FOBT Test Results or Finding	SNOMED CT: 59614000, 167667006, 389076003, 71711000112103

*Codes subject to change

Care for Older Adults (COA) ●

The COA measure evaluates the percentage of persons 66 years of age and older who had both of the following in the measurement period:

- Medication review
 - Perform an annual medication review of the person's medications, including prescription medications, over-the-counter medications, and herbal or supplemental therapies.
- Functional status assessment

This assessment measures the person's ability to perform daily tasks and helps to identify any functional decline. For Wellcare By Meridian (D-SNP) patients, please indicate in the medical record:

- Activities of Daily Living (ADL) — Note and date when activities like bathing, dressing, eating, transferring, using toilet, and walking were assessed.
- Instrumental Activities of Daily Living (IADL) — Note and date when activities like shopping, driving or using public transportation, meal preparation, housework, taking medications, and using the telephone were assessed.
- Standardized functional status assessments — Note results and dates of assessments like the Assessment of Living Skills and Resources (ALSAR), the Barthel Index and the Physical Self-Maintenance Scale, Bayer Activities of Daily Living (B-ADL) Scale, or Extended Activities of Daily Living (EADL) Scale.

COA Measure Codes

Description	Codes*
Medication Review with Medication List (Requires a code from both value sets)	Medication List: CPT-CAT-II: 1159F HCPCS: G8427 Medication Review: CPT: 90863, 99483, 99605, 99606 CPT-CAT-II: 1160F
Medication Review: Transitional Care Management Services	CPT: 99495, 99496
Functional Status Assessment	CPT: 99483 CPT-CAT-II: 1170F HCPCS: G0438, G0439

*Codes subject to change

Controlling High Blood Pressure (CBP) & Blood Pressure Control for Patients with Hypertension (BPC-E)*



* Note: Introduced in MY 2025, the BPC-E measure is reported using the ECDS method, and the denominator includes a pharmacy data method with a hypertension diagnosis. The CBP measure is expected to be retired by NCQA in MY 2028.

The CBP and BPC-E measures evaluate the percentage of patients 18 to 85 years of age who had a diagnosis of hypertension (HTN) and whose last blood pressure (BP) reading of the measurement year was adequately controlled (<140/90 mm Hg).

- Consider taking two readings at each office visit. Sometimes a second blood pressure reading is lower.
- Do not round blood pressure; always use exact numbers
- Include CPT-CAT-II codes on claims to ensure the blood pressure reading is received by the health plan

CBP Measure Codes

Description	Codes*
Systolic Greater Than/Equal to 140	CPT-CAT-II: 3077F
Systolic 130-139	CPT-CAT-II: 3075F
Systolic Less Than 130	CPT-CAT-II: 3074F
Diastolic Greater Than/Equal to 90	CPT-CAT-II: 3080F
Diastolic 80-89	CPT-CAT-II: 3079F
Diastolic Less Than 80	CPT-CAT-II: 3078F

*Codes subject to change

Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions (FMC) ●●

The FMC measure evaluates Emergency Department (ED) visits for persons 18 years of age and older who have multiple high-risk chronic conditions who had a follow-up service within 7 days of the ED visit on or between January 1 and December 24 of the measurement period where the person was age 18 or older on the date of the visit.

FMC Measure Codes

Description	Codes
BH Outpatient	CPT: 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
Case Management Encounter	CPT: 99366 HCPCS: T1016, T1017, T2022, T2023
Complex Care Management Services	CPT: 99439, 99487, 99489-99491 HCPCS: G0506
Electroconvulsive Therapy (Requires POS code)	CPT: 90870 ICD-10 PCS: GZB0ZZZ, GZB2ZZZ, GZB4ZZZ
Substance Use Disorder Services	CPT: 99408, 99409 HCPCS: G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012
Transitional Care Management Services	CPT: 99495, 99496

FMC Measure Codes (continued)

Description	Codes
Visit Setting Unspecified (Requires POS code)	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255
Outpatient and Telehealth	CPT: 98009-98016, 98966-98968, 98970-98972, 98980-98981, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99421-99423, 99429, 99441-99443, 99455-99458, 99483 HCPCS: G0071, G0402, G0438-G0439, G0463, G2010, G2012, G2250-G2252, T1015
Substance Abuse Counseling and Surveillance	ICD-10-CM: Z71.41, Z71.51
Partial Hospitalization or Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485

*Codes subject to change. Refer to the Appendix for POS codes.

Persistence of Beta-Blocker Treatment After a Heart Attack (PBH)

The PBH measure evaluates the percentage of patients 18 years of age and older during the measurement year who were hospitalized and discharged from July 1 of the year prior to June 30 of the measurement year with a diagnosis of AMI and who received persistent beta-blocker treatment for six months after discharge.

Eligible Medications
<ul style="list-style-type: none"> • Acebutolol • Atenolol • Atenolol-chlorthalidone • Bendroflumethiazide-nadolol • Betaxolol • Bisoprolol • Bisoprolol-hydrochlorothiazide • Carvedilol • Hydrochlorothiazide-metoprolol • Hydrochlorothiazide-propranolol • Labetalol • Metoprolol • Nadolol • Nebivolol • Pindolol • Propranolol • Sotalol • Timolol

Pharmacotherapy Management of COPD Exacerbation (PCE) ●●●

The PCE measure evaluates the percentage of COPD exacerbations for patients 40 years of age and older who had an acute inpatient discharge or ED visit on or between January 1–November 30 and were dispensed appropriate medications.

Two rates are reported:

- Dispensed a systemic **corticosteroid** (or there was evidence of an active prescription) within **14 days of the event**
- Dispensed a **bronchodilator** (or there was evidence of an active prescription) within **30 days of the event**

Eligible Medications	
<ul style="list-style-type: none">• Aclidinium-bromide• Albuterol• Albuterol-ipratropium• Arformoterol• Budesonide-formoterol• Cortisone• Dexamethasone• Formoterol• Formoterol-aclidinium• Formoterol-glycopyrrolate• Formoterol-mometasone• Fluticasone furoate - umeclidinium-vilanterol• Fluticasone-salmeterol	<ul style="list-style-type: none">• Fluticasone-vilanterol• Ipratropium• Levalbuterol• Metaproterenol• Methylprednisolone• Olodaterol• Olodaterol-tiotropium• Prednisolone• Prednisone• Salmeterol• Tiotropium• Umeclidinium• Umeclidinium-Vilanterol

Plan All Cause Readmissions (PCR) ●●●●

The PCR measure evaluates persons 18 years of age and older who had an acute inpatient and observation stay that was followed by an unplanned acute readmission for any diagnosis within 30 days and the predicted probability of an acute readmission.

What Providers Can Do

- See the patient within seven days of discharge
- Educate the patient and family about the diagnosis and care plan
- Review medication list
- Establish care goals with the patient
- Identify barriers to health for the patient that may have contributed to the hospitalization and discuss strategies for addressing them

How Meridian Can Help

- Meridian will notify PCPs during the Transition of Care (TOC) process
- A TOC letter is faxed to the PCP within 24 hours of discharge
- Meridian's Interdisciplinary Care Team (ICT) reviews individual care plans
- Educate patients on appropriate Emergency Department utilization

Transitions of Care (TRC) ● ●

The TRC measure evaluates the percentage of discharges for persons 18 years of age and older who had each of the following:

- Notification of Inpatient Admission (NIA)
- Receipt of Discharge Information (RDI)
- Patient Engagement After Inpatient Discharge (PE)
- Medication Reconciliation Post-Discharge (MRPD)

Remember to:

- Document receipt of notification of inpatient admission on the day of admission through two days after admission (for a total of 3 days)
- Document receipt of notification of discharge on the day of discharge through two days after discharge (for a total of 3 days)
- Document patient engagement within 30 days after discharge (e.g., office visits, visits to the home, telehealth)
- Document medication reconciliation on the date of discharge through 30 days after discharge (for a total of 31 days)

Patient engagement provided within 30 days after discharge. Do not include patient engagement that occurs on the date of discharge. The following meet criteria for patient engagement:

- An outpatient visit
- A telephone visit
- Transitional care management services
- An e-visit or virtual check-in

Medication reconciliation must be conducted by a prescribing practitioner, clinical pharmacist, physician assistant, or registered nurse on the date of discharge through 30 days after discharge (31 total days).

TRC Measure Codes

Description	Codes*
Patient Engagement: Transitional Care Management Services	CPT: 99495-99496
MRPD: Medication Reconciliation Encounter	CPT: 99483, 99495-99496

TRC Measure Codes (continued)

Description	Codes*
MRPD: Medication Reconciliation Intervention	CPT: 99605, 99606 CPT-CAT-II: 1111F SNOMED CT: 430193006, 428701000124107
Outpatient and Telehealth	CPT: 98000-98016, 98966, 98967, 98968, 98970-98972, 98980-98981, 99202-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99429, 99441-99443, 99455-99458, 99483 HCPCS: G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2250, G2251, G2252, T1015

*Codes subject to change

Use of Imaging Studies for Low Back Pain (LBP)

The LBP measure evaluates persons 18 to 75 years of age with a principal diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis. A *lower rate indicates better performance*.

Numerator: An imaging study with a diagnosis of uncomplicated low back pain on the IESD or in the 28 days following the IESD

LBP Measure Codes

Description	Codes*
Imaging Study	CPT: 72020, 72040, 72050, 72052, 72070, 72072, 72074, 72080-72084, 72100, 72110, 72114, 72120, 72125-72133, 72141, 72142, 72146-72149, 72156-72158, 72200, 72202, 72220
Uncomplicated Low Back Pain	ICD-10: M47.26-M47.28, M47.816-M47.818, M47.896-M47.898, M48.061, M48.07, M48.08, M51.16-M51.17, M51.26-M51.27, M51.36-M51.37, M51.86-M51.87, M53.2X6-M53.2X8, M53.3, M53.86-M53.88, M54.16-M54.18, M54.30-M54.32, M54.40-M54.42, M54.5, M54.50, M54.51, M54.59, M54.89, M54.9, M99.03-M99.04, M99.23, M99.33, M99.43, M99.53, M99.63, M99.73, M99.83, M99.84, S33.100A, S33.100D, S33.100S, S33.110A, S33.110D, S33.110S, S33.120A, S33.120D, S33.120S, S33.130A, S33.130D, S33.130S, S33.140A, S33.140D, S33.140S, S33.5XXA, S33.6XXA, S33.8XXA, S33.9XXA, S39.002A, S39.002D, S39.002S, S39.012A, S39.012D, S39.012S, S39.82XA, S39.82XD, S39.82XS, S39.92XA, S39.92XD, S39.92XS

*Codes subject to change

According to the National Committee for Quality Assurance (NCQA), 75 percent of American adults will experience low back pain at some time in their lives, making it one of the most common reasons patients seek healthcare services.

Please consider imaging studies for lower back pain only if red flags are present or if there is no improvement after four weeks.

Alternative Recommendations

Meridian encourages its providers to refer to the Agency for Healthcare Research and Quality website (ahrq.gov) for standards and guidelines in managing your patients' acute lower back pain.

Source: ncqa.org

Behavioral Health



Adherence to Antipsychotic Medications for People with Schizophrenia (SAA) ●●●

The SAA measure evaluates persons 18 years of age and older in the measurement period with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.

Antipsychotic Medications

- Amitriptyline-perphenazine
- Anripiprazole lauroxil
- Albuterol-ipratropium
- Aripiprazole
- Asenapine
- Brexpiprazole
- Cariprazine
- Chlorpromazine
- Clozapine
- Fluphenazine
- Fluphenazine decanoate
- Haloperidol
- Haloperidol decanoate
- Iloperidfone
- Loxapine
- Lumateperone
- Lurasidone
- Molindone
- Olanzapine
- Paliperidone
- Paliperidone palmitate
- Perphenazine
- Prochlorperazine
- Quetiapine
- Risperidone
- Thioridazine
- Thiothixene
- Trifluoperazine
- Ziperidone

Follow-Up After Emergency Department Visit for Mental Illness (FUM) ●●●●

The FUM measure evaluates persons 6 years of age and older with a principal diagnosis of mental illness or self-harm who had a follow-up visit for mental illness after an Emergency Department (ED) visit.

Two rates are reported:

- The percentage of ED visits for which the person received follow-up within 7 days of the visit (8 total days)
- The percentage of ED visits for which the person received follow-up within 30 days of the visit (31 total days)

FUM 7 & 30 Day Follow-up Measure Codes

Description	Additional Criteria (if applicable)	Codes*
BH Outpatient	Requires diagnosis of a mental health disorder	CPT: 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-2020, T1015
Electroconvulsive Therapy	Requires POS code	CPT: 90870 ICD-10: GZB0ZZZ, GZB2ZZZ-GZB4ZZZ
Online Assessment (e-visit or virtual check-in)	Requires diagnosis of a mental health disorder	CPT: 98016, 98970-98972, 98980, 98981, 99421-99423, 99457, 99458 HCPCS: G0071, G2010, G2012, G2250-G2252
Partial Hospitalization or Intensive Outpatient	Requires diagnosis of a mental health disorder	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Telephone Visits	Requires diagnosis of a mental health disorder	CPT: 98008-98015, 98966-98968, 99441-99443

FUM 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Visit Setting Unspecified	Requires POS code; requires diagnosis of a mental health disorder	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255
Peer Support Services	Requires diagnosis of a mental health disorder	HCPCS: G0140, G0177, H0024, H0025, H0038, H0039, H0040, H0046, H2014, H2023, S9445, T1012, T1016, T1017
Psychiatric Collaborative Care Management		CPT: 99492-99494
Residential Behavioral Health Treatment		HCPCS: H0017-H0019, T2048
Mental Health Diagnosis		ICD-10-CM: F03.90, F03.911, F03.918, F03.92, F03.93, F03.94, F03.A0, F03.A11, F03.A18, F03.A2, F03.A3, F03.A4, F03.B0, F03.B11, F03.B18, F03.B2, F03.B3, F03.B4, F03.C0, F03.C11, F03.C18, F03.C2, F03.C3, F03.C4, F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.81, F34.89, F34.9, F39, F40.00,

FUM 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Mental Health Diagnosis (continued)		F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.248, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.81, F43.89, F43.9, F44.0, F44.1, F44.2, F44.4, F44.5, F44.6, F44.7, F44.81, F44.89, F44.9, F45.0, F45.1, F45.20, F45.21, F45.22, F45.29, F45.41, F45.42, F45.8, F45.9, F48.1, F48.2, F48.8, F48.9, F50.00, F50.01, 50.010-F50.014, F50.019-F50.024, F50.02, F50.020-F50.024, F50.029, F50.2, F50.20- F50.25, F50.81, F50.82, F50.89, F50.9, F51.01, F51.02, F51.03, F51.04, F51.05, F51.09, F51.11, F51.12, F51.13, F51.19, F51.3, F51.4, F51.5, F51.8, F51.9, F52.0, F52.1, F52.21, F52.22, F52.31, F52.32, F52.4, F52.5, F52.6, F52.8, F52.9, F53.0, F53.1, F59, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3, F63.81, F63.89, F63.9, F64.0, F64.1, F64.2, F64.8, F64.9, F65.0, F65.1, F65.2, F65.3, F65.4, F65.50, F65.51, F65.52, F65.81, F65.89, F65.9, F66, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F69, F80.0, F80.1, F80.2, F80.4, F80.81, F80.82, F80.89, F80.9, F81.0, F81.2, F81.81, 50.810-F50.814, F50.819, F81.89, F81.9,

FUM 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Mental Health Diagnosis (continued)		F82, F50.83, F50.84, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F88, F89, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9, F95.0, F95.1, F95.2, F95.8, F95.9, F98.0, F98.1, F98.21, F98.29, F98.3, F98.4, F98.5, F98.8, F98.9, F99

*Codes subject to change. Refer to the Appendix for POS codes.

Follow-Up After Emergency Department Visit for Substance Use (FUA) ●●●●

The percentage of Emergency Department (ED) visits among persons 13 years of age and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose, for which there was follow-up.

Two rates are reported:

- The percentage of ED visits for which the person received follow-up within 30 days of the visit (31 total days)
- The percentage of ED visits for which the person received follow-up within 7 days of the visit (8 total days)

FUA 7 & 30 Day Follow-up Measure Codes

Description	Additional Criteria (if applicable)	Codes*
Behavioral Health Assessment	Requires diagnosis of SUD or drug overdose	CPT: 99408, 99409 HCPCS: G0211, G0396, G0397, G0442, G2011, H0001, H0002, H0031, H0049
Online Assessments	Requires provision by a mental health provider	CPT: 98016, 98970-98972, 98980-98981, 99421-99423, 99457, 99458 HCPCS: G0071, G2010, G2012, G2250-G2252
OUD Weekly Drug Treatment Service	Requires diagnosis of SUD or drug overdose	HCPCS: G2067, G2068, G2073, G0533

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
OUD Monthly Office-Based Treatment	Requires diagnosis of SUD or drug overdose	HCPCS: G2069, G2086, G2087
OUD Weekly Non-Drug Service	Requires diagnosis of SUD or drug overdose	HCPCS: G2074-G2077, G2080
Partial Hospitalization or Intensive Outpatient	Requires POS code; requires diagnosis of SUD or drug overdose	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Peer Support Services	Requires diagnosis of SUD or drug overdose	HCPCS: G0177, H0024, H0025, H0038-H0040, H0046, H2014, H2023, S9445, T1012, T1016, T1017
Substance Use Disorder Services		CPT: 99408, 99409 HCPCS: G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012
Substance Use Services		HCPCS: H0006, H0028
Visit Setting Unspecified	Requires POS code; requires diagnosis of SUD or drug overdose	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255 POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 52, 53, 57, 71, 72
Substance Abuse Counseling and Surveillance		ICD-10-CM: Z71.41, Z71.51

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
BH Outpatient	Requires provision by a mental health provider; requires diagnosis of SUD or drug overdose	CPT: 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99483, 99492-99494, 99510 HCPCS: G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0034, H0036, H0037, H0039, H0040, H2000, H2011, H2013-H2020, T1015
Telephone Visits	Requires provision by a mental health provider	CPT: 98008-98015, 98966, 98967, 98968, 99441, 99442, 99443
AOD Medication Treatment		HCPCS: G2069, G2073, G0533, H0020, H0033, J0571, J0572, J0573, J0574, J0575, J2315, Q9991, Q9992, S0109

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
AOD Abuse and Dependence		ICD-10-CM: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132, F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251,

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
AOD Abuse and Dependence (continued)		F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Substance Induced Disorders		ICD-10-CM: F10.90, F10.920, F10.921, F10.929, F10.930, F10.931, F10.932, F10.939, F10.94, F10.950, F10.951, F10.959, F10.96, F10.97, F10.980, F10.981, F10.982, F10.988, F10.99, F11.90, F11.920, F11.921, F11.922, F11.929, F11.93, F11.94, F11.950, F11.951, F11.959, F11.981, F11.982, F11.988, F11.99, F12.90, F12.920, F12.921, F12.922, F12.929, F12.93, F12.950, F12.951, F12.959, F12.980, F12.988, F12.99, F13.90, F13.920, F13.921, F13.929, F13.930, F13.931, F13.932, F13.939, F13.94, F13.950, F13.951, F13.959, F13.96, F13.97, F13.980, F13.981, F13.982, F13.988, F13.99, F14.90, F14.920, F14.921, F14.922, F14.929, F14.93, F14.94, F14.950, F14.951, F14.959, F14.980, F14.981, F14.982, F14.988, F14.99, F15.90, F15.920, F15.921, F15.922, F15.929, F15.93, F15.94, F15.950, F15.951, F15.959, F15.980, F15.981, F15.982, F15.988, F15.99, F16.90, F16.920, F16.921, F16.929, F16.94, F16.950, F16.951, F16.959, F16.980, F16.983, F16.988, F16.99, F18.90, F18.920, F18.921, F18.929, F18.94, F18.950, F18.951, F18.959, F18.97, F18.980, F18.988, F18.99, F19.90, F19.920, F19.921, F19.922, F19.929, F19.930, F19.931, F19.932, F19.939, F19.94, F19.950, F19.951, F19.959, F19.96, F19.97, F19.980, F19.981, F19.982, F19.988, F19.99

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Unintentional Drug Overdose		ICD-10-CM: T40.0X1A, T40.0X1D, T40.0X1S, T40.0X4A, T40.0X4D, T40.0X4S, T40.1X1A, T40.1X1D, T40.1X1S, T40.1X4A, T40.1X4D, T40.1X4S, T40.2X1A, T40.2X1D, T40.2X1S, T40.2X4A, T40.2X4D, T40.2X4S, T40.3X1A, T40.3X1D, T40.3X1S, T40.3X4A, T40.3X4D, T40.3X4S, T40.411A, T40.411D, T40.411S, T40.414A, T40.414D, T40.414S, T40.421A, T40.421D, T40.421S, T40.424A, T40.424D, T40.424S, T40.491A, T40.491D, T40.491S, T40.494A, T40.494D, T40.494S, T40.5X1A, T40.5X1D, T40.5X1S, T40.5X4A, T40.5X4D, T40.5X4S, T40.601A, T40.601D, T40.601S, T40.604A, T40.604D, T40.604S, T40.691A, T40.691D, T40.691S, T40.694A, T40.694D, T40.694S, T40.711A, T40.711D, T40.711S, T40.714A, T40.714D, T40.714S, T40.721A, T40.721D, T40.721S, T40.724A, T40.724D, T40.724S, T40.8X1A, T40.8X1D, T40.8X1S, T40.8X4A, T40.8X4D, T40.8X4S, T40.901A, T40.901D, T40.901S, T40.904A, T40.904D, T40.904S, T40.991A, T40.991D, T40.991S, T40.994A, T40.994D, T40.994S, T41.0X1A, T41.0X1D, T41.0X1S, T41.0X4A, T41.0X4D, T41.0X4S, T41.1X1A, T41.1X1D, T41.1X1S, T41.1X4A, T41.1X4D, T41.1X4S, T41.201A, T41.201D, T41.201S, T41.204A, T41.204D, T41.204S, T41.291A, T41.291D, T41.291S, T41.294A, T41.294D, T41.294S, T41.3X1A, T41.3X1D, T41.3X1S, T41.3X4A, T41.3X4D, T41.3X4S, T41.41XA, T41.41XD, T41.41XS, T41.44XA, T41.44XD, T41.44XS, T41.5X1A,

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Unintentional Drug Overdose (continued)		T41.5X1D, T41.5X1S, T41.5X4A, T41.5X4D, T41.5X4S, T42.3X1A, T42.3X1D, T42.3X1S, T42.3X4A, T42.3X4D, T42.3X4S, T42.4X1A, T42.4X1D, T42.4X1S, T42.4X4A, T42.4X4D, T42.4X4S, T43.601A, T43.601D, T43.601S, T43.604A, T43.604D, T43.604S, T43.621A, T43.621D, T43.621S, T43.624A, T43.624D, T43.624S, T43.631A, T43.631D, T43.631S, T43.634A, T43.634D, T43.634S, T43.641A, T43.641D, T43.641S, T43.644A, T43.644D, T43.644S, T43.651A, T43.651D, T43.651S, T43.654A, T43.654D, T43.654S, T43.691A, T43.691D, T43.691S, T43.694A, T43.694D, T43.694S, T51.0X1A, T51.0X1D, T51.0X1S, T51.0X4A, T51.0X4D, T51.0X4S
AOD Medication Treatment		HCPCS: G0533, G2069, G2073, H0020, H0033, J0571, J0572, J0573, J0574, J0575, J2315, Q9991, Q9992, S0109

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
AOD Abuse and Dependence		ICD-10-CM: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132, F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251,

FUA 7 & 30 Day Follow-up Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
AOD Abuse and Dependence (continued)		F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29

*Codes subject to change. Refer to the Appendix for POS codes.

Follow-Up After High Intensity Care for Substance Disorder (FUI) ●●●

The FUI measure evaluates the percentage of acute inpatient hospitalizations, residential treatment, or detoxification visits for a diagnosis of substance use disorder among persons 13 years of age and older that result in a follow-up visit or service for substance use disorder.

FUI measure reports two rates:

- The percentage of visits or discharges for which the person received follow-up for substance use disorder within the 30 days after the visit or discharge
- The percentage of visits or discharges for which the person received follow-up for substance use disorder within the 7 days after the visit or discharge.

FUI Measure Codes

Description	Additional Criteria (if applicable)	Codes
BH Outpatient	Requires diagnosis of SUD	CPT: 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99510, 99483, 99492-99494, 99510 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
Online Assessment	Requires diagnosis of SUD	CPT: 98016, 98970-98972, 98980-98981, 99421-99423, 99457, 99458 HCPCS: G0071, G2010, G2012, G2250-G2252
Visit Setting Unspecified	Requires POS code; requires diagnosis of SUD	CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255
Partial Hospitalization or Intensive Outpatient	Requires diagnosis of SUD	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485

FUI Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes
Residential Behavioral Health Treatment	Requires diagnosis of SUD	HCPCS: H0017-H0019, T2048
Substance Use Disorder Services	Requires diagnosis of SUD	CPT: 99408, 99409 HCPCS: G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012
Telephone Visits	Requires diagnosis of SUD	CPT: 98008-98015, 98966-98968, 99441-99443
OUD Monthly Office Based Treatment	Requires diagnosis of SUD	HCPCS: G2086-G2087, G2069
OUD Weekly Drug Treatment Service	Requires diagnosis of SUD	HCPCS: G2067, G2068-G2073
OUD Weekly Non-Drug Service	Requires diagnosis of SUD	HCPCS: G2074-G2077, G2080
Substance Use Counseling and Surveillance		ICD-10-CM: Z71.41, Z71.51
AOD Abuse and Dependence		ICD-10: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19,

FUI Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes
AOD Abuse and Dependence (continued)		F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132, F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229,

FUI Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes
AOD Abuse and Dependence (continued)		F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29

*Codes subject to change. Refer to the Appendix for POS codes.

Follow-Up After Hospitalization for Mental Illness (FUH)

The FUH measure evaluates the percentage of discharges for persons 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health service.

Two rates are reported:

- Discharges for which the person received **follow-up within 7 days after discharge**. Do not include visits that occur on the date of discharge.
- Discharges for which the person received **follow-up within 30 days after discharge**. Do not include visits that occur on the date of discharge.

Note: As of MY 2025, follow-up visits can be completed by any qualified provider, including Primary Care Providers (PCPs), if the claim includes a mental health diagnosis and the provider is licensed to bill the appropriate codes.

FUH Measure Codes

Description	Additional Criteria (if applicable)	Codes*
Visit Setting Unspecified	Requires POS code; requires provision by a mental health provider	CPT: 90791, 90792, 90832- 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255
BH Outpatient Visit	Requires provision by a mental health provider	CPT: 98000-98007, 98960- 98962, 99202-99205, 99078, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
Partial Hospitalization/ Intensive Outpatient		HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Residential Behavioral Health Treatment		HCPCS: H0017-H0019, T2048
Electroconvulsive Therapy	Requires POS code	CPT: 90870 ICD-10: GZB0ZZZ, GZB2ZZZ- GZB4ZZZ
Transitional Care	Requires provision by a mental health provider	CPT: 99495, 99496
Telephone Visit with Telehealth	Requires provision by a mental health provider; requires any diagnosis of a mental health disorder	CPT: 98966-98968, 99441- 99443
Psychiatric Collaborative Care Management		CPT: 99492-99494 HCPCS: G0512

FUH Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Peer Support Services	Requires any diagnosis of a mental health disorder	CPT: G0140, G0177, H0024, H0025, H0038, H0039, H0040, H0046, H2014, H2023, S9445, T1012, T1016, T1017
Mental Health Diagnosis		ICD-10-CM: F03.90, F03.911, F03.918, F03.92, F03.93, F03.94, F03.A0, F03.A11, F03.A18, F03.A2, F03.A3, F03.A4, F03.B0, F03.B11, F03.B18, F03.B2, F03.B3, F03.B4, F03.C0, F03.C11, F03.C18, F03.C2, F03.C3, F03.C4, F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.81, F34.89, F34.9, F39, F40.00, F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.248, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12,

FUH Measure Codes (continued)

Description	Additional Criteria (if applicable)	Codes*
Mental Health Diagnosis (continued)		F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.81, F43.89, F43.9, F44.0, F44.1, F44.2, F44.4, F44.5, F44.6, F44.7, F44.81, F44.89, F44.9, F45.0, F45.1, F45.20, F45.21, F45.22, F45.29, F45.41, F45.42, F45.8, F45.9, F48.1, F48.2, F48.8, F48.9, F50.00, F50.01, F50.02, F50.2, F50.81, F50.82, F50.89, F50.9, F51.01, F51.02, F51.03, F51.04, F51.05, F51.09, F51.11, F51.12, F51.13, F51.19, F51.3, F51.4, F51.5, F51.8, F51.9, F52.0, F52.1, F52.21, F52.22, F52.31, F52.32, F52.4, F52.5, F52.6, F52.8, F52.9, F53.0, F53.1, F59, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3, F63.81, F63.89, F63.9, F64.0, F64.1, F64.2, F64.8, F64.9, F65.0, F65.1, F65.2, F65.3, F65.4, F65.50, F65.51, F65.52, F65.81, F65.89, F65.9, F66, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F69, F80.0, F80.1, F80.2, F80.4, F80.81, F80.82, F80.89, F80.9, F81.0, F81.2, F81.81, F81.89, F81.9, F82, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F88, F89, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9, F95.0, F95.1, F95.2, F95.8, F95.9, F98.0, F98.1, F98.21, F98.29, F98.3, F98.4, F98.5, F98.8, F98.9, F99

*Codes subject to change. Refer to the Appendix for POS codes.

Initiation and Engagement of Substance Use Disorder (IET) ●●●●●

The IET measure evaluates the percentage of patients 13 years of age and older with a new episode of substance use disorder that resulted in treatment initiation and engagement who received the following:

- Initiation of substance use disorder treatment: Percentage of patients who initiate treatment through an inpatient new SUD episode that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth, or medication treatment within 14 days
- Engagement of substance use disorder treatment: Percentage of patients who had new SUD episodes that have evidence of treatment engagement within 34 days of the initiation

Alcohol Use Disorder Treatment Medications

- Acamprosate (oral; delayed-release tablet)
- Disulfiram (oral)
- Naltrexone (oral and injectable)

Opioid Use Disorder Treatment Medications

- Buprenorphine (implant)
- Buprenorphine (injection)
- Buprenorphine (sublingual tablet)
- Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)
- Naltrexone (injectable)
- Naltrexone (oral)

IET Medications

Description	Codes*
Buprenorphine Implant	HCPGS: G2070, G2072, J0570
Buprenorphine Injection	HCPGS: G0533, G2069, Q9991, Q9992
Buprenorphine Naloxone	HCPGS: J0572-J0575
Buprenorphine Oral	HCPGS: H0033, J0571
Buprenorphine Oral Weekly	HCPGS: G2068, G2079
Detoxification	HCPGS: H0008, H0009, H0010-H0014
Methadone Oral	HCPGS: H0020, S0109
Methadone Oral Weekly	HCPGS: G2067, G2078
Naltrexone Injection	HCPGS: G2073, J2315

*Codes subject to change

IET OP Measure Codes

Description	Codes*
BH Outpatient	<p>CPT: 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99349, 99350, 99381-99383, 99384-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510</p> <p>HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015</p> <p>Note: Include ICD-10 code for Alcohol abuse and other drug dependence diagnosis</p>
Online Assessments	<p>CPT: 98016, 98970-98972, 98980, 98981, 99421-99523, 99444, 99457, 99458</p> <p>HCPCS: G0071, G2010, G2012, G2250-G2251</p>
Visit Setting Unspecified (Requires POS code)	<p>CPT: 90791, 90792, 90832-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255</p>
Partial Hospitalization or Intensive Outpatient	HCPCS: H0035, H2001, H2012, G0410, G0411
Substance Use Disorder Services	<p>CPT: 99408, 99409</p> <p>HCPCS: G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012</p>
OUD Monthly Office Based Treatment	HCPCS: G2086, G2087, G2069
Telephone Visits	CPT: 98966-98968, 99441-99443
OUD Weekly Drug Treatment Services	HCPCS: G2067, G2073, G0533
OUD Weekly Non-Drug Service	HCPCS: G2074-G2077, G2080
Substance Abuse Counseling & Surveillance	ICD-10-CM: Z71.41, Z71.51
Alcohol Abuse and Dependence	ICD-10-CM: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29
Opioid Abuse and Dependence	ICD-10-CM: F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220

IET OP Measure Codes (continued)

Description	Codes*
Opioid Abuse and Dependence (continued)	F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29
Other Drug Abuse and Dependence	ICD-10-CM: F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132, F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29

For the follow-up treatments, include an ICD-10 diagnosis for Alcohol or Other Drug Dependence from the Mental, Behavioral and Neurodevelopmental Disorder Section of ICD-10 along with a procedure code for the preventive service, evaluation, and management consultation or counseling service.

* Codes listed are subject to change. Meridian recognizes that the circumstances around the services provided may not always directly support/match the codes. It is crucial that the medical record documentation describes the services rendered in order to support the medical necessity and use of these codes. Refer to the Appendix for POS codes.

Pharmacotherapy for Opioid Use Disorder (POD) ●●●●

The POD measure evaluates persons 16 years of age and older with a diagnosis of Opioid Use Disorder (OUD) and a new OUD pharmacotherapy that lasted at least 180 days.

The treatment period of 180+ days begins on the new OUD pharmacotherapy event date through 179 days without a gap in treatment of 8 or more consecutive days.

Opioid Use Disorder Treatment Medications

Description	Prescription	Medication Lists
Antagonist	• Naltrexone (oral)	• Naltrexone Oral Medications List
Antagonist	• Naltrexone (injectable)	• Naltrexone Injection Medications List
Partial agonist	• Buprenorphine (sublingual tablet)	• Buprenorphine Oral Medications List
Partial agonist	• Buprenorphine (injection)	• Buprenorphine Injection Medications List
Partial agonist	• Buprenorphine (implant)	• Buprenorphine Implant Medications List
Partial agonist	• Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)	• Buprenorphine Naloxone Medications List
Agonist	• Methadone (oral)	• NA (refer to Note below)

Methadone is not included on the medication lists for this measure. Methadone for OUD administered or dispensed by federally certified opioid treatment programs (OTP) is billed on a medical claim. A pharmacy claim for methadone would be indicative of treatment for pain rather than OUD.

Unhealthy Alcohol Use Screening and Follow-Up (ASF-E)

ASF-E evaluates the percentage of persons 18 years of age and older who were screened for unhealthy alcohol use using a standardized instrument and, if screened positive, received appropriate follow-up care within 2 months (61 days total).

Eligible standard assessment instruments that have been normalized and validated for the adult patient population with thresholds for positive findings include:

Screening Instrument	Total Score LOINC Codes*	Positive Finding
Alcohol Use Disorders Identification Test (AUDIT) screening instrument	75624-7	Total score ≥ 8
Alcohol Use Disorders Identification Test Consumption (AUDIT-C) screening instrument	75626-2	Total score ≥ 4 for men Total score ≥ 3 for women
Single-question screen (for men): "How many times in the past year have you had 5 or more drinks in a day?" Single-question screen (for women and all adults older than 65 years): "How many times in the past year have you had 4 or more drinks in a day?"	88037-7 75889-6	Response ≥ 1 Response ≥ 1

*Codes subject to change

Description	Codes*
Alcohol Counseling or Other Follow-Up Care	ICD-10-CM: Z71.41 CPT: 99408, 99409 HCPCS: G0396, G0397, G0443, G2011, H0005, H0007, H0015, H0016, H0022, H0050, H2035, H2036, T1006, T1012 SNOMED CT: 20093000, 23915005, 24165007, 64297001, 386449006, 408945004, 408947007, 408948002, 413473000, 707166002, 429291000124102

*Codes subject to change

Screening for Depression and Follow-Up Plan (CDF-AD) ●

This is a CMS core set measure. For more information, visit [cms.gov](https://www.cms.gov).

The CDF-AD measure evaluates persons 18 years of age and older screened for depression who have never had a diagnosis of depression or bipolar disorder on the date of the encounter or 14 days prior to the date of the encounter using an age-appropriate standardized depression screening tool. If positive, a follow-up plan is documented on the date of the eligible encounter.

CDF-B Codes to Document Depression Screen

Code	Description
G8431	Screening for depression is documented as being positive and a follow-up plan is documented
G8510	Screening for depression is documented as negative; a follow-up plan is not required

**Codes subject to change*

Screening for Depression and Follow-Up Plan (CDF-CH) ●

This is a CMS core set measure. For more information, visit [cms.gov](https://www.cms.gov).

The CDF-CH Measure evaluates patients ages 12 to 17 screened for depression who have never had a diagnosis of depression or bipolar disorder on the date of the encounter or 14 days prior to the date of the encounter using an age-appropriate standardized depression screening tool. If positive, a follow-up plan is documented on the date of the eligible encounter.

CDF-B Codes to Document Depression Screen

Code	Description
G8431	Screening for depression is documented as being positive and a follow-up plan is documented
G8510	Screening for depression is documented as negative; a follow-up plan is not required

**Codes subject to change*

Depression Screening and Follow-Up for Adolescents and Adults (DSF-E)

DSF-E evaluates the percentage of persons 12 years of age and older who were screened for clinical depression between January 1 and December 1 of the measurement period, using a standardized instrument and, if screened positive, received follow-up care within 30 days

* Refer to the Appendix for a list of Approved Depression Screening Instruments, Codes, and Positive Findings

Depression Remission or Response for Adolescents and Adults (DRR-E)

DRR-E evaluates the percentage of persons 12 years of age and older with a diagnosis of depression and an elevated PHQ-9 score, who had evidence of response or remission within 120–240 days (4 to 8 months) of the elevated score.

- **Follow-Up PHQ-9:** The percentage of persons who have a follow-up PHQ-9 score documented within 120–240 days (4 to 8 months) after the initial elevated PHQ-9 score.
- **Depression Remission:** The percentage of persons who achieved remission of depression symptoms, as demonstrated by the most recent PHQ-9 total score of <5 during the depression follow-up period (within 120–240 days [4–8 months] after the initial elevated PHQ-9 score).
- **Depression Response:** The percentage of persons who showed response to treatment for depression, as demonstrated by the most recent PHQ-9 total score of at least 50% lower than the PHQ-9 score associated with the diagnosis, documented during the depression follow-up period (within 120–240 days [4–8 months] after the initial elevated PHQ-9 score).

DRR-E Codes

Depression/PHQ-9 Follow-Up, Remission, and Response	LOINC Codes*
Persons 12 years of age and older	44261-6, 89204-2**

*Codes subject to change

**For adolescents ages 12–17 assessed with Patient Health Questionnaire Modified for Teens (PHQ-9M)[®]

Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E)



DMS-E evaluates the percentage of persons 12 years of age and older with a diagnosis of major depression or dysthymia who had an outpatient encounter with a PHQ-9 score present in their record in the same assessment period as the encounter.

The measurement period is divided into three assessment periods with specific dates of service:

- Assessment period 1: January 1–April 30
- Assessment period 2: May 1–August 31
- Assessment period 3: September 1–December 31

DMS-E Codes

Utilization of PHQ-9	LOINC Codes*
Persons 12 years of age and older	44261-6, 89204-2**

*Codes subject to change

**For adolescents ages 12–17 assessed with Patient Health Questionnaire Modified for Teens (PHQ-9M)®

Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) ●●

The APP measure evaluates patients 1 to 17 years of age who had a new prescription for an antipsychotic medication and had documentation of psychosocial care as first-line treatment.

Antipsychotic Medications	
• Aripiprazole	• Molindone
• Aripiprazole lauroxil	• Olanzapine
• Asenapine	• Paliperidone
• Brexpiprazole	• Perphenazine
• Cariprazine	• Pimozide
• Chlorpromazine	• Quetiapine
• Clozapine	• Risperidone
• Fluphenazine	• Thioridazine
• Haloperidol	• Thiothixene
• Iloperidone	• Trifluoperazine
• Loxapine	• Ziprasidone
• Lurasidone	

APP Measure Codes

Description	CPT*	HCPCS*
Psychosocial Care	90832-90834, 90836-90840, 90845-90847, 90849, 90853, 90875, 90876, 90880	G0176, G0177, G0409, G0410, G0411, H0004, H0035-H0040, H2000, H2001, H2011-H2014, H2017-H2020, S0201, S9480, S9484, S9485
Residential Behavioral Health Treatment		H0017, H0018, H0019, T2048

*Codes subject to change

General Health



Appropriate Testing for Pharyngitis (CWP) ●●●●

The CWP measure evaluates the percentage of episodes for persons 3 years of age and older where the person was diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode.

CWP Measure Codes

Description	ICD-10-CM Diagnosis*
Acute pharyngitis	J02.9
Acute tonsillitis	J03.90
Streptococcal sore throat	J02.0

*Codes subject to change

Group A Strep Test Codes

CPT*
87070, 87071, 87081, 87430, 87650-87652, 87880

*Codes subject to change

Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis (AAB) ●●●●

The AAB measure evaluates patients 3 months of age and older who had a diagnosis of acute bronchitis/bronchiolitis that did not result in an antibiotic dispensing event. A *lower rate indicates better performance*.

Treating Uncomplicated Acute Bronchitis

- Avoid prescribing antibiotics
- Treat presented symptoms only
- Prescribe antitussive agents for short-term relief of coughing

AAB Antibiotic Medications

- Amikacin
- Amoxicillin
- Amoxicillin/clavulanate
- Ampicillin
- Ampicillin/sulbactam
- Azithromycin
- Aztreonam
- Cefaclor
- Cefadroxil
- Cefazolin
- Cefdinir
- Cefepime
- Cefixime
- Cefotaxime
- Cefotetan
- Cefoxitin
- Cefpodoxime
- Cefprozil
- Ceftazidime
- Ceftriaxone
- Cefuroxime
- Cephalexin
- Chloramphenicol
- Ciprofloxacin
- Clarithromycin
- Clindamycin
- Dalforpristin/quinupristin
- Daptomycin
- Dicloxacillin
- Doxycycline
- Erythromycin
- Fosfomycin
- Gemifloxacin
- Gentamicin
- Levofloxacin
- Lincomycin
- Linezolid
- Metronidazole
- Minocycline
- Moxifloxacin
- Nafcillin
- Nitrofurantoin
- Nitrofurantoin, macrocrystals/nitrofurantoin, monohydrate
- Ofloxacin
- Oxacillin
- Penicillin G benzathine
- Penicillin G benzathine/penicillin G procaine
- Penicillin G potassium
- Penicillin G procaine
- Penicillin G sodium
- Penicillin V potassium
- Piperacillin/tazobactam
- Rifampin
- Streptomycin
- Sulfadiazine
- Sulfamethoxazole/trimethoprim
- Tetracycline
- Tobramycin
- Trimethoprim
- Vancomycin

Follow-Up After Acute and Urgent Care Visits for Asthma (AAF-E) ●●●

This is a first-year measure in MY 2026, and the Asthma Medication Ratio (AMR) measure was retired.

The AAF-E measure evaluates the percentage of persons 5-64 years of age with an urgent care visit, acute inpatient discharge, observation stay discharge, or ED visit with a diagnosis of asthma that had a corresponding outpatient follow-up visit with a diagnosis of asthma within 30 days.

Description	Codes*
Outpatient and Telehealth	CPT: 98000-98016, 98966-98968, 98970-98972, 98980-98981, 99202-99205, 99211-99215, 99242-99245, 99341-99342, 99344-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99421-99423, 99429, 99441-99443, 99455-99458, 99483 HCPCS: G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2250-G2252, T1015 SNOMED: 50357006, 77406008, 84251009, 86013001, 90526000, 185317003, 185463005, 185464004, 185465003, 209099002, 281036007, 314849005, 386472008, 386473003, 401267002, 439740005, 866149003, 3391000175108, 444971000124105, 456201000124103
Asthma	ICD-10-CM: J45.20, J45.21, J45.22, J45.30, J45.31, J45.32, J45.40, J45.41, J45.42, J45.50, J45.51, J45.52, J45.901, J45.902, J45.909, J45.990, J45.991, J45.998

*Codes subject to change

Social Need Screening and Intervention (SNS-E)

SNS-E evaluates the percentage of persons who were screened, using prespecified instruments, at least once during the measurement period for unmet food, housing, and transportation needs, and who received a corresponding intervention within one month if they screened positive.

Food Screening: Persons with a documented result for food insecurity screening performed between January 1 and December 1 of the measurement period.

Eligible screening instruments with thresholds for positive findings include:

Food Insecurity Instruments	Screening Item LOINC Codes	Positive Finding LOINC Codes
Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	88122-7	LA28397-0 LA6729-3
	88123-5	LA28397-0 LA6729-3
American Academy of Family Physicians (AAFP) Social Needs Screening Tool	88122-7	LA28397-0 LA6729-3
	88123-5	LA28397-0 LA6729-3
American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	88122-7	LA28397-0 LA6729-3
	88123-5	LA28397-0 LA6729-3
Health Leads Screening Panel ^{®1}	95251-5	LA33-6
Hunger Vital Sign ^{TM1} (HVS)	88124-3	LA19952-3
Protocol for Responding to and Assessing Patients' Assets, Risks and Experiences [PRAPARE] ^{®1}	93031-3	LA30125-1
Safe Environment for Every Kid (SEEK) ^{®1}	95400-8	LA33-6
	95399-2	LA33-6
U.S. Household Food Security Survey [U.S. FSS]	95264-8	LA30985-8 LA30986-6
U.S. Adult Food Security Survey [U.S. FSS]	95264-8	LA30985-8 LA30986-6
U.S. Child Food Security Survey [U.S. FSS]	95264-8	LA30985-8 LA30986-6
U.S. Household Food Security Survey—Six-Item Short Form [U.S. FSS]	95264-8	LA30985-8 LA30986-6
We Care Survey	96434-6	LA32-8

Food Insecurity Instruments (continued)	Screening Item LOINC Codes	Positive Finding LOINC Codes
WellRx Questionnaire	93668-2	LA33-6

¹ Proprietary; cost or licensing requirement may be associated with use.

*Codes subject to change

Food Intervention: Persons who received a food insecurity intervention on or up to 30 days after the date of the first positive food insecurity screen (31 days total)

Food Insecurity Procedures Codes*
CPT: 96156, 96160, 96161, 97802, 97803, 97804 HCPCS: S5170, S9470 SNOMED CT: 1759002, 61310001, 103699006, 308440001, 385767005, 710824005, 710925007, 711069006, 713109004, 1002223009, 1002224003, 1002225002, 1004109000, 1004110005, 1148446004, 1162436000, 1230338004, 441041000124100, 441201000124108, 441231000124100, 441241000124105, 441251000124107, 441261000124109, 441271000124102, 441281000124104, 441291000124101, 441301000124100, 441311000124102, 441321000124105, 441331000124108, 441341000124103, 441351000124101, 445291000124103, 445301000124102, 445641000124105, 461481000124109, 462481000124102, 462491000124104, 464001000124109, 464011000124107, 464021000124104, 464031000124101, 464041000124106, 464051000124108, 464061000124105, 464071000124103, 464081000124100, 464091000124102, 464101000124108, 464111000124106, 464121000124103, 464131000124100, 464141000124105, 464151000124107, 464161000124109, 464171000124102, 464181000124104, 464191000124101, 464201000124103, 464211000124100, 464221000124108, 464231000124106, 464241000124101, 464251000124104, 464261000124102, 464271000124109, 464281000124107, 464291000124105, 464301000124106, 464311000124109, 464321000124101, 464331000124103, 464341000124108, 464351000124105, 464361000124107, 464371000124100, 464381000124102, 464401000124102, 464411000124104, 464421000124107, 464431000124105, 464611000124102, 464621000124105, 464631000124108, 464641000124103, 464651000124101, 464661000124104, 464671000124106, 464681000124109, 464691000124107, 464701000124107, 464721000124102, 467591000124102, 467601000124105, 467611000124108, 467621000124100, 467631000124102, 467641000124107, 467651000124109, 467661000124106, 467671000124104, 467681000124101, 467691000124103, 467711000124100, 467721000124108, 467731000124106, 467741000124101, 467751000124104, 467761000124102, 467771000124109, 467781000124107, 467791000124105, 467801000124106, 467811000124109, 467821000124101, 468401000124109, 470231000124107, 470241000124102, 470261000124103, 470281000124108, 470291000124106, 470301000124107, 470311000124105, 470321000124102, 470591000124109, 470601000124101, 470611000124103, 471111000124101, 471121000124109, 471131000124107, 472151000124109, 472331000124100, 551101000124107

*Codes subject to change

Housing Screening: Persons with a documented result for housing instability, homelessness, or housing inadequacy screening performed between January 1 and December 1 of the measurement period

Eligible screening instruments with thresholds for positive findings include:

Housing Instability and Homelessness Instruments	Screening Item LOINC Codes	Positive Finding LOINC Codes
Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	71802-3	LA31994-9 LA31995-6
	99550-6	LA33-6
American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	71802-3	LA31994-9 LA31995-6
	98976-4	LA33-6
Children's Health Watch Housing Stability Vital Signs ^{TM1}	98977-2	≥3
	98978-0	LA33-6
Health Leads Screening Panel ^{®1}	99550-6	LA33-6
Protocol for Responding to and Assessing Patients' Assets, Risks and Experiences [PRA-PARE] ^{®1}	93033-9	LA33-6
	71802-3	LA30190-5
We Care Survey	96441-1	LA33-6
WellRx Questionnaire	93669-0	LA33-6

**Codes subject to change*

Housing Inadequacy Instruments	Screening Item LOINC Codes	Positive Finding LOINC Codes
Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	96778-6	LA28580-1 LA31996-4 LA31997-2 LA31998-0 LA31999-8 LA32000-4 LA32001-2
American Academy of Family Physicians (AAFP) Social Needs Screening Tool	96778-6	LA28580-1 LA32693-6 LA32694-4 LA32695-1 LA32696-9 LA32001-2 LA32691-0
American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	96778-6	LA28580-1 LA31996-4 LA31997-2 LA31998-0 LA31999-8 LA32000-4 LA32001-2 LA33-6
Norwalk Community Health Center Screening Tool [NCHC]	99134-9	LA33-6
	99135-6	LA28580-1 LA31996-4 LA31997-2 LA31998-0 LA31999-8 LA32000-4 LA32001-2

¹Proprietary; cost or licensing requirement may be associated with use.

*Codes subject to change

Housing Intervention: Persons who received an intervention corresponding to the type of housing need identified on or up to 30 days after the date of the first positive housing screen (31 days total)

Inadequate Housing Procedures Codes*

CPT: 96156, 96160, 96161
SNOMED CT: 49919000, 308440001, 710824005, 711069006, 1148446004, 1148813002, 1148815009, 1148823006, 1162436000, 1230338004, 461481000124109, 462481000124102, 462491000124104, 464001000124109, 464011000124107, 464021000124104, 464131000124100, 464161000124109, 464291000124105, 464301000124106, 464311000124109, 464611000124102, 470231000124107, 470431000124106, 470441000124101, 470451000124104, 470461000124102, 470591000124109, 470601000124101, 470611000124103, 471111000124101, 471121000124109, 471131000124107, 472151000124109, 472201000124100, 472211000124102, 472231000124108, 472251000124101, 472331000124100, 472371000124102, 480881000124103, 480891000124100, 480911000124103, 480951000124102, 551041000124105, 551051000124107, 551061000124109, 551071000124102, 551081000124104, 551101000124107

*Codes subject to change

Transportation Screening: Persons with a documented result for transportation insecurity screening performed between January 1 and December 1 of the measurement period

Eligible screening instruments with thresholds for positive findings include:

Transportation Insecurity Instruments	Screening Item LOINC Codes	Positive Finding LOINC Codes
Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	93030-5	LA33-6
	99594-4	LA33-6
American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	99594-4	LA30134-3 LA33093-8
Comprehensive Universal Behavior Screen (CUBS)	89569-8	LA29232-8 LA29233-6 LA29234-4
Health Leads Screening Panel ^{®1}	99553-0	LA33-6
Inpatient Rehabilitation Facility - Patient Assessment Instrument (IRF-PAI)—version 4.0 [CMS Assessment]	93030-5	LA30133-5 LA30134-3
Outcome and assessment information set (OASIS) form—version E—Discharge from Agency [CMS Assessment]	93030-5	LA30133-5 LA30134-3

Transportation Insecurity Instruments (continued)	Screening Item LOINC Codes	Positive Finding LOINC Codes
Outcome and assessment information set (OASIS) form—version E—Resumption of Care [CMS Assessment]	93030-5	LA30133-5 LA30134-3
Outcome and assessment information set (OASIS) form—version E—Start of Care [CMS Assessment]	93030-5	LA30133-5 LA30134-3
Protocol for Responding to and Assessing Patients' Assets, Risks and Experiences [PRAPARE] ^{®1}	93030-5	LA30133-5 LA30134-3
PROMIS ^{®1}	92358-1	LA30024-6 LA30026-1 LA30027-9
WellRx Questionnaire	93671-6	LA33-6

¹ Proprietary; cost or licensing requirement may be associated with use.

*Codes subject to change

Transportation Intervention: Patients who received a transportation insecurity intervention on or up to 30 days after the date of the first positive transportation screen (31 days total)

Transportation Insecurity Procedures Codes*

CPT: 96156, 96160, 96161
 SNOMED CT: 308440001, 710824005, 711069006, 1148446004, 1162436000, 1230338004, 461481000124109, 462481000124102, 462491000124104, 464001000124109, 464011000124107, 464021000124104, 464131000124100, 464161000124109, 464291000124105, 464301000124106, 464311000124109, 464611000124102, 470231000124107, 470591000124109, 470601000124101, 470611000124103, 471111000124101, 471121000124109, 471131000124107, 472151000124109, 472331000124100, 551101000124107, 551111000124105, 551121000124102, 551141000124109, 551161000124108, 551191000124100, 551201000124102, 551211000124104, 551221000124107, 551231000124105, 551241000124100, 551251000124103, 551261000124101, 551271000124108, 551281000124106, 551291000124109, 551301000124105, 551311000124108, 551321000124100, 551331000124102, 551341000124107, 551351000124109, 551361000124106, 551371000124104, 551381000124101, 551401000124101, 551421000124106, 551431000124109, 610961000124100, 610971000124107, 610981000124105, 610991000124108, 611001000124109, 611011000124107, 611021000124104, 611031000124101, 611041000124106, 611051000124108, 611061000124105, 611071000124103, 611081000124100, 611101000124108, 611121000124103, 611281000124107, 611291000124105, 611301000124106, 611311000124109, 611321000124101, 611331000124103, 611341000124108, 611351000124105, 611361000124107, 611371000124100, 611381000124102, 611391000124104, 611401000124102, 611411000124104, 611421000124107, 611431000124105, 611441000124100

*Codes subject to change

Tobacco Use Screening and Cessation Intervention (TSC-E)

This is a first-year measure in MY 2026. Only the Electronic Clinical Data Systems (ECDS) method is used for reporting.

The TSC-E Measure evaluates the percentage of persons 12 years of age and older who were screened for commercial tobacco product use at least once during the measurement period, and who received tobacco cessation intervention if identified as a tobacco user.

Two rates are reported:

- **Tobacco Use Screening:** The percentage of persons who were screened for tobacco use
- **Cessation Intervention:** The percentage of persons who were identified as a tobacco user and who received tobacco cessation intervention

Description	Codes*
Tobacco Use Cessation Counseling	ICD-10-CM: Z71.6

**Codes subject to change*

Pediatric Health



Childhood Immunization Status (CIS-E) ● ● ● ☁

CIS-E evaluates the percentage of children 2 years of age during the measurement year who completed the following immunizations on or before child's second birthday.

Immunization	Required Doses
DTaP (Diphtheria, Tetanus, Acellular Pertussis)	4 doses
PCV (Pneumococcal Conjugate)	4 doses
HiB (Haemophiles Influenza Type B)	3 doses
Hep B (Hepatitis B)	3 doses
IPV (Polio; Inactivated Polio Virus)	3 doses
Flu (Influenza)	2 doses
RV (Rotavirus)	2 or 3 dose series
Hep A (Hepatitis A)	1 dose on or between the 1st and 2nd birthday
MMR (Measles, Mumps, Rubella)	1 dose on or between the 1st and 2nd birthday
VZV (Chicken Pox; Varicella zoster)	1 dose on or between the 1st and 2nd birthday

CIS-E Measure Codes

Description	Codes*
DTaP Immunization	CVX: 20, 50, 106, 107, 110, 120, 146
DTaP Vaccine Procedure	CPT: 90697, 90698, 90700, 90723
Anaphylaxis due to Diphtheria, Tetanus or Pertussis Vaccine	SNOMED CT: 428281000124107, 428291000124105
Encephalitis due to Diphtheria, Tetanus or Pertussis Vaccine	SNOMED CT: 192710009, 192711008, 192712001
HiB Immunization	CVX: 17, 46, 47, 48, 49, 50, 51, 120, 146, 148, 198
HiB Vaccine Procedure	CPT: 90644, 90647, 90648, 90697, 90698, 90748
Anaphylaxis due to the HiB Vaccine	SNOMED CT: 433621000124101
Hepatitis B Immunization	CVX: 08, 44, 45, 51, 110, 146, 198
Hepatitis B Vaccine Procedure	CPT: 90697, 90723, 90740, 90744, 90747, 90748 HCPCS: G0010
History of Hepatitis B Illness	ICD-10-CM: B16.0, B16.1, B16.2, B16.9, B17.0, B18.0, B18.1, B19.10, B19.11 SNOMED CT: 1116000, 13265006, 26206000, 38662009, 50167007, 53425008, 60498001, 61977001, 66071002, 76795007, 111891008, 165806002, 186624004, 186626002, 186639003, 235864009, 235865005, 235869004, 235871004, 271511000, 313234004, 406117000, 424099008, 424340000, 442134007, 442374005, 446698005, 838380002, 1230342001, 153091000119109, 551621000124109
Anaphylaxis due to the Hepatitis B Vaccine	SNOMED CT: 428321000124101
IPV Immunization	CVX: 10, 89, 110, 120, 146, 198
IPV Procedure	CPT: 90697, 90698, 90713, 90723
Anaphylaxis due to IPV Vaccine	SNOMED CT: 471321000124106
MMR Immunization	CVX: 03, 94
MMR Vaccine Procedure	CPT: 90707, 90710

CIS-E Measure Codes (continued)

Description	Codes*
History of Measles Illness	<p>ICD-10-CM: B05.0, B05.1, B05.2, B05.3, B05.4, B05.81, B05.89, B05.9</p> <p>SNOMED CT: 14189004, 28463004, 38921001, 60013002, 74918002, 111873003, 161419000, 186561002, 186562009, 195900001, 240483006, 240484000, 359686005, 371111005, 406592004, 417145006, 424306000, 105841000119101</p>
History of Mumps Illness	<p>ICD-10-CM: B26.0, B26.1, B26.2, B26.3, B26.81, B26.82, B26.83, B26.84, B26.85, B26.89, B26.9</p> <p>SNOMED CT: 10665004, 17121006, 31524007, 31646008, 36989005, 40099009, 44201003, 63462008, 72071001, 74717002, 75548002, 78580004, 89231008, 89764009, 111870000, 161420006, 235123001, 236771002, 237443002, 240526004, 240527008, 240529006, 371112003, 1163539003, 105821000119107</p>
History of Rubella Illness	<p>ICD-10-CM: B06.00, B06.01, B06.02, B06.09, B06.81, B06.82, B06.89, B06.9</p> <p>SNOMED CT: 10082001, 13225007, 19431000, 36653000, 51490003, 64190005, 79303006, 128191000, 161421005, 165792000, 186567003, 186570004, 192689006, 231985001, 232312000, 240485004, 253227001, 406112006, 406113001, 1092361000119109, 10759761000119100</p>
Anaphylaxis due to the MMR Vaccine	SNOMED CT: 471331000124109
Pneumococcal Conjugate Immunization	CVX: 109, 133, 152, 215, 216
Pneumococcal Conjugate Vaccine Procedure	CPT: 90670, 90671, 90677 HCPCS: G0009
Anaphylaxis due to Pneumococcal Vaccine	SNOMED CT: 471141000124102
Varicella Zoster (VZV) Immunization	CVX: 21, 94
Varicella Zoster (VZV) Vaccine Procedure	CPT: 90710, 90716

CIS-E Measure Codes (continued)

Description	Codes*
History of Varicella Zoster (Chicken Pox) Illness	<p>ICD-10-CM: B01.0, B01.11, B01.12, B01.2, B01.81, B01.89, B01.9, B02.0, B02.1, B02.21, B02.22, B02.23, B02.24, B02.29, B02.30, B02.31, B02.32, B02.33, B02.34, B02.39, B02.7, B02.8, B02.9</p> <p>SNOMED CT: 4740000, 10698009, 21954000, 23737006, 24059009, 36292003, 38907003, 42448002, 49183009, 55560002, 87513003, 111859007, 111861003, 161423008, 186524006, 195911009, 230176008, 230198004, 230262004, 230536009, 232400003, 235059009, 240468001, 240470005, 240471009, 240472002, 240473007, 240474001, 309465005, 371113008, 397573005, 400020001, 402897003, 402898008, 402899000, 410500004, 410509003, 421029004, 422127002, 422446008, 422471006, 422666006, 423333008, 423628002, 424353002, 424435009, 424801004, 424941009, 425356002, 426570007, 428633000, 713250002, 713733003, 713964006, 715223009, 723109003, 838357005, 1163465001, 1163483009, 1179456002, 12551000132107, 12561000132105, 12571000132104, 98541000119101, 331071000119101, 681221000119108, 1087131000119102, 15678761000119105, 15678801000119102, 15678841000119100, 15680201000119106, 15680241000119108, 15680281000119103, 15681321000119100, 15681401000119101, 15685081000119102, 15685121000119100, 15685201000119100, 15685281000119108, 15936581000119108, 15936621000119108, 15989271000119107, 15989311000119107, 15989351000119108, 15991711000119108, 15991751000119109, 15991791000119104, 15992351000119104, 16000751000119105, 16000791000119100, 16000831000119106</p>
Anaphylaxis due to the VZV Vaccine	SNOMED CT: 471341000124104
Hep A Immunization	CVX: 31, 83, 85

CIS-E Measure Codes (continued)

Description	Codes*
History of Hepatitis A illness	ICD-10-CM: B15.0, B15.9 SNOMED CT: 16060001, 18917003, 25102003, 40468003, 43634002, 79031007, 111879004, 165997004, 206373002, 278971009, 310875001, 424758008, 428030001, 105801000119103
Anaphylaxis due to hepatitis A vaccine	SNOMED CT: 471311000124103
Influenza Immunization	CVX: 88, 140, 141, 150, 153, 155, 158, 161, 171, 186, 320
Influenza Vaccine Procedure	CPT: 90655, 90657, 90661, 90673, 90674, 90685, 90686, 90687, 90688, 90689, 90756 HCPCS: G0080
Influenza Virus LAIV Immunization (Recommended for children 2 years of age and older. When administered on the child's second birthday, meets the criteria for one of two required doses for Flu immunization.)	CVX: 111, 149
Influenza Virus LAIV Vaccine Procedure	CPT: 90660, 90672
Anaphylaxis due to the Influenza Vaccine	SNOMED CT: 471361000124100
Combination 3	DTaP, IPV, MMR, HiB, hepatitis B, VZV, and pneumococcal indicators
Combination 7	DTaP, IPV, MMR, HiB, hepatitis B, VZV, pneumococcal, hepatitis A, and rotavirus indicators
Combination 10	DTaP, IPV, MMR, HiB, hepatitis B, VZV, pneumococcal, hepatitis A, rotavirus and influenza indicators
Rotavirus Immunization (2 dose schedule)	CPT: 90681 CVX: 119
Rotavirus Immunization (3 dose schedule)	CVX: 116, 122
Anaphylaxis due to the Rotavirus Vaccine	SNOMED CT: 428331000124103

*Codes subject to change

NOTE: Rotavirus is either 2 dose OR 3 dose for compliance

Follow-Up Care for Children Prescribed ADHD Medication (ADD-E)

The ADD-E measure evaluates the percentage of children newly prescribed attention deficit hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed.

Two rates are reported:

- **Initiation Phase:** Percentage of patients 6 to 12 years of age with a prescription dispensed for ADHD medication, who had one follow-up visit with a practitioner with prescribing authority during the 30-day Initiation Phase
- **Continuation and Maintenance (C&M) Phase:** Percentage of patients 6 to 12 years of age with a prescription dispensed for ADHD medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended.

ADD-E Measure Codes

Description	Codes*
Outpatient POS	POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
BH Outpatient Visit	CPT: 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99510, 99483, 99492-99494 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015
Health and Behavior Assessment/Intervention	CPT: 96156, 96158, 96159, 96164, 96165, 96167, 96168, 96170, 96171
Online Assessments	CPT: 98016, 98970-98972, 98980, 98981, 99421-99423, 99457, 99458 HCPC: G0071, G2010, G2012, G2250-G2252

ADD-E Measure Codes (continued)

Description	Codes*
Partial Hospitalization/ Intensive Outpatient	HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 SNOMED CT: 7133001, 305345009, 305346005, 305347001, 391038005, 391042008, 391043003, 391046006, 391047002, 391048007, 391054008, 391055009, 391056005, 391133003, 391150001, 391151002, 391152009, 391153004, 391170007, 391185001, 391186000, 391187009, 391188004, 391191004, 391192006, 391194007, 391195008, 391207001, 391208006, 391209003, 391210008, 391211007, 391228005, 391229002, 391232004, 391252003, 391254002, 391255001, 391256000
Telehealth POS	POS: 02, 10
Telephone Visits	CPT: 98008-98015, 98966-98968, 99441-99443 SNOMED CT: 185317003, 314849005, 386472008, 386473003, 401267002
Visit Setting Unspecified Value Set	CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255 POS: 52, 53

*Codes subject to change

Immunizations for Adolescents (IMA-E)

Beginning in MY 2025, this measure transitioned to the Electronic Clinical Data Systems (ECDS) reporting method only.

IMA-E evaluates the percentage of adolescents 13 years of age in the measurement year who completed the following immunizations on or before the person's 13th birthday.

Immunization	Required Doses
Meningococcal	1 dose between the 11th and 13th birthdays
Tdap (Tetanus, Diphtheria Toxoids and Acellular Pertussis)	1 dose between the 10th and 13th birthdays
HPV (Human Papillomavirus)	2 or 3 doses between the 9th and 13th birthdays

Note: HPV is either two doses with 146 days between the first and second dose **OR** three doses on different dates of service

IMA-E Measure Codes

Description	Codes*
Meningococcal-serogroup A,C,W, and Y Immunization	CVX: 32, 108, 114, 136, 147, 167, 203, 316, 328
Meningococcal Vaccine Procedure	CPT: 90619, 90623, 90624, 90733, 90734 SNOMED CT: 871874000, 428271000124109, 16298691000119102
Anaphylaxis due to the Meningococcal Vaccine	SNOMED CT: 428301000124106
Tdap Immunization	CVX: 115
Anaphylaxis due to the Tetanus, Diphtheria, or Pertussis Vaccine	SNOMED CT: 428281000124107, 428291000124105
Encephalitis due to the Tetanus, Diphtheria, or Pertussis Vaccine	SNOMED CT: 192710009, 192711008, 192712001
HPV Immunization	CVX: 62, 118, 137, 165
HPV Vaccine Procedure	CPT: 90649, 90650, 90651
Anaphylaxis due to the HPV Vaccine	SNOMED CT: 428241000124101

*Codes subject to change

Lead Screening in Children (LSC-E)

Beginning in MY 2026, this measure transitioned to the Electronic Clinical Data Systems (ECDS) reporting method only.

The LSC-E measure evaluates the percentage of children 2 years of age in the measurement year who had one or more capillary or venous lead blood test for lead poisoning on or before patient's second birthday.

Lead Test

CPT: 83655

LOINC: 10368-9, 10912-4, 14807-2, 17052-2, 25459-9, 27129-6, 32325-3, 5674-7, 77307-7

**Codes subject to change*

In order to ensure that children receive appropriate public health follow-up services, physicians and other health providers have an obligation to report blood lead results greater than or equal to 10mg/dL within 48 hours to the Illinois Department of Public Health (IDPH) Childhood Lead Poisoning Reporting System. Providers using the IDPH laboratory are not required to report blood lead results.

Illinois Department of Public Health

Illinois Lead Program

535 W. Jefferson Street

Springfield, IL 62761

P: 217-782-3517

www.idph.state.il.us

The Illinois Department of Healthcare and Family Services (HFS) encourages providers to send all blood lead specimens to the IDPH laboratory for analysis. Providers who utilize the state laboratory for blood lead analysis can order supplies for blood lead specimen collection free of charge by calling the IDPH Laboratory Shipping Section at 217-524-6222, or by downloading the Clinical Supplies Requisition Form from the HFS website: <https://www.dph.illinois.gov/>.

Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E)

The APM-E measure evaluates the percentage of children and adolescents 1 to 17 years of age who had two or more antipsychotic prescriptions and had metabolic testing.

Three rates are reported:

- Percentage of children and adolescents on antipsychotics who received blood glucose testing
- Percentage of children and adolescents on antipsychotics who received cholesterol testing
- Percentage of children and adolescents on antipsychotics who received blood glucose and cholesterol testing

APM-E Measure Codes

Description	Codes*
HbA1c Lab Tests	CPT: 83036, 83037 LOINC: 17855-8, 17856-6, 4548-4, 4549-2, 96595-4 SNOMED CT: 43396009, 313835008
HbA1c Level Greater Than 9.0	CPT-CAT-II: 3046F
HbA1c Level Greater Than or Equal To 7.0 and Less Than 8.0	CPT-CAT-II: 3051F
HbA1c Level Greater Than or Equal To 8.0 and Less Than or Equal To 9.0	CPT-CAT-II: 3052F
HbA1c Level Less Than 7.0	CPT-CAT-II: 3044F
HbA1c Test Result or Finding	CPT-CAT-II: 3044F, 3046F, 3051F, 3052F SNOMED CT: 165679005, 451061000124104
Glucose Lab Tests	CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951 LOINC: 10450-5, 1492-8, 1494-4, 1496-9, 1499-3, 1501-6, 1504-0, 1507-3, 1514-9, 1518-0, 1530-5, 1533-9, 1554-5, 1557-8, 1558-6, 17865-7, 20436-2, 20437-0, 20438-8, 20440-4, 2345-7, 26554-6, 41024-1, 49134-0, 6749-6, 9375-7 SNOMED CT: 22569008, 33747003, 52302001, 72191006, 73128004, 88856000, 104686004, 167086002, 167087006, 167088001, 167095005, 167096006, 167097002, 250417005,

APM-E Measure Codes (continued)

Description	Codes*
Glucose Lab Tests (continued)	271061004, 271062006, 271063001, 271064007, 271065008, 275810004, 302788006, 302789003, 308113006, 313474007, 313545000, 313546004, 313624000, 313626003, 313627007, 313628002, 313630000, 313631001, 313697000, 313698005, 313810002, 412928005, 440576000, 443780009, 444008003, 444127006
Glucose Test Results or Finding	SNOMED CT: 166890005, 166891009, 166892002, 166914001, 166915000, 166916004, 166917008, 166918003, 166919006, 166921001, 166922008, 166923003, 442545002, 444780001, 1179458001
LDL-C Lab Tests	CPT: 80061, 83700, 83701, 83704, 83721 LOINC: 12773-8, 13457-7, 18261-8, 18262-6, 2089-1, 49132-4, 55440-2, 96259-7 SNOMED CT: 113079009, 166833005, 166840006, 166841005, 167074000, 167075004, 314036004
LDL-C Test Result or Finding	CPT-CAT-II: 3048F, 3049F, 3050F
Most recent LDL-C less than 100 mg/dL (CAD) (DM)	CPT-CAT-II: 3048F
Most recent LDL-C 100-129 mg/dL (CAD) (DM)	CPT-CAT-II: 3049F
Most recent LDL-C greater than or equal to 130 mg/dL (CAD) (DM)	CPT-CAT-II: 3050F
Cholesterol Lab Tests	CPT: 82465, 83718, 83722, 84478 LOINC: 2085-9, 2093-3, 2571-8, 3043-7, 9830-1 SNOMED CT: 14740000, 28036006, 77068002, 104583003, 104584009, 104586006, 104784006, 104990004, 104991000, 121868005, 166832000, 166838001, 166839009, 166849007, 166850007, 167072001, 167073006, 167082000, 167083005, 167084004, 271245006, 275972003, 314035000, 315017003, 390956002, 412808005, 412827004, 443915001

APM-E Measure Codes (continued)

Description	Codes*
Cholesterol Test Result or Finding	SNOMED CT: 166830008, 166848004, 259557002, 365793008, 365794002, 365795001, 365796000, 439953004, 707122004, 707123009, 1162800007, 1172655006, 1172656007, 67991000119104

*Codes subject to change

Oral Evaluation, Dental Services (OED) ●●●

The OED measure evaluates the percentage of persons under 21 years of age who received a comprehensive or periodic oral evaluation with a dental provider during the measurement period.

OED Measure Codes

Description	Codes*
Oral Evaluation	CDT: D0120, D0145, D0150

*Codes subject to change

Topical Fluoride for Children (TFC) ●●

The TFC measure evaluates the percentage of children 1 to 4 years of age who received at least two fluoride varnish applications during the measurement period.

TFC Measure Codes

Description	Codes*
Application of Fluoride Varnish	CPT: 99188 CDT: D1206 SNOMED: 313042009

*Codes subject to change

Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) ●●●

The WCC measure evaluates the percentage of children 3 to 17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of each of the following during the measurement period:

- BMI Percentile
- Counseling for Nutrition
- Counseling for Physical Activity

WCC Measure Codes

Description	Codes*
BMI Percentile	ICD-10: Z68.51-Z68.56 LOINC: 59574-4, 59575-1, 59576-9
Nutrition Counseling	CPT: 97802-97804 ICD-10: Z71.3 HCPCS: G0270, G0271, G0447, S9449, S9452, S9470
Physical Activity Counseling	HCPCS: G0447, S9451 ICD-10: Z02.5, Z71.82

*Codes subject to change

Well-Child and Adolescent Well-Care Visits (W30 ●●/WCV ●●●)

The W30/WCV measure evaluates the percentage of persons within designated ages who had comprehensive well-care visit(s) as defined in each measure, with a PCP or an OB/GYN practitioner during the measurement year.

Well-Child Visits in the First 30 Months of Life (W30)

Months of Life: The percentage of children who had the following number of well-child visits with a PCP during the last 15 months

Two rates are reported:

- Well-Child Visits in the First 15 Months
 - Children who turned 15 months old during the measurement period:
Six or more well-child visits
- Well-Child Visits for Age 15 Months–30 Months
 - Children who turned 30 months old during the measurement period:
Two or more well-child visits

W30 Measure Codes

CPT*	HCPCS*	ICD-10*
99381-99385, 99391-99395, 99461	G0438, G0439, S0302, S0610, S0612, S0613	Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z76.1, Z76.2

**Codes subject to change*

(WCV) Child and Adolescent Well-Care Visits: Persons 3 to 21 years of age who had a least one comprehensive well-care visit with a PCP or an OB/GYN

WCV Measure Codes

CPT*	HCPCS*	ICD-10*
99381-99385, 99391-99395, 99461	G0438, G0439, S0302, S0610, S0612, S0613	Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z76.1, Z76.2

**Codes subject to change*

Pharmacy Measures



Use of High-Risk Medications in Older Adults (DAE) ●●

The DAE measure evaluates the percentage of persons 67 years of age and older who had at least two dispensing events for the same high-risk medication class.

Three rates are reported:

1. The percentage of Medicare adults 67 years of age and older who had at least two dispensing events for high-risk medications to avoid from the same drug class
2. The percentage of Medicare adults 67 years of age and older who had at least two dispensing events for high-risk medications to avoid from the same drug class, except for appropriate diagnoses
3. Total rate (the sum of the two numerators divided by the denominator, deduplicating for patients in both numerators)

Note: The measure reflects potentially inappropriate medication use in older adults, both for medications where any use is inappropriate (Rate 1) and for medications where use under all, but specific indications is potentially inappropriate (Rate 2). A lower rate represents better performance.

High-Risk Medications

Drug Class	Prescriptions
Anticholinergics, first-generation antihistamines	<ul style="list-style-type: none">• Brompheniramine• Chlorpheniramine• Cyproheptadine• Dimenhydrinate• Diphenhydramine (oral)• Doxylamine• Hydroxyzine• Meclizine• Promethazine• Triprolidine

High-Risk Medications (continued)

Drug Class	Prescriptions
Anticholinergics, anti-Parkinson agents	<ul style="list-style-type: none"> • Benztropine (oral) • Trihexyphenidyl
Antispasmodics	<ul style="list-style-type: none"> • Atropine • Chlordiazepoxide-clidinium • Dicyclomine • Hyoscyamine • Scopolamine
Antithrombotic	<ul style="list-style-type: none"> • Dipyridamole (oral, excluding extended release)
Cardiovascular, alpha agonists, central	<ul style="list-style-type: none"> • Guanfacine
Cardiovascular, other	<ul style="list-style-type: none"> • Nifedipine (excluding extended release)
Central nervous system, antidepressants	<ul style="list-style-type: none"> • Amitriptyline • Amoxapine • Clomipramine • Desipramine • Imipramine • Nortriptyline • Paroxetine
Central nervous system, barbiturates	<ul style="list-style-type: none"> • Butalbital • Phenobarbital • Primidone
Central nervous system, vasodilators	<ul style="list-style-type: none"> • Ergoloid mesylates (USP)
Central nervous system, other	<ul style="list-style-type: none"> • Meprobamate
Endocrine system, estrogens with or without progestins; include only oral and topical patch products	<ul style="list-style-type: none"> • Conjugated estrogen (USP) • Esterified estrogen (USP) • Estradiol • Estropipate
Endocrine system, sulfonylureas, long-duration	<ul style="list-style-type: none"> • Glimepiride • Glyburide
Endocrine system, desiccated thyroid	<ul style="list-style-type: none"> • Desiccated thyroid
Endocrine system, megestrol	<ul style="list-style-type: none"> • Megestrol
Nonbenzodiazepine hypnotics	<ul style="list-style-type: none"> • Eszopiclone • Zaleplon • Zolpidem

High-Risk Medications (continued)

Drug Class	Prescriptions
Pain medications, skeletal muscle relaxants	<ul style="list-style-type: none"> • Carisoprodol • Chlorzoxazone • Cyclobenzaprine • Metaxalone • Methocarbamol • Orphenadrine
Pain medications, meperidine	<ul style="list-style-type: none"> • Meperidine
Pain medications, other	<ul style="list-style-type: none"> • Indomethacin • Ketorolac, includes parenteral and oral

High-Risk Medications with Days Supply Criteria

Drug Class	Prescriptions	Days Supply Criteria
Anti-Infectives, other	<ul style="list-style-type: none"> • Nitrofurantoin • Nitrofurantoin macrocrystals-monohydrate 	<ul style="list-style-type: none"> • >90 days

High-Risk Medications with Average Daily Dose Criteria

Drug Class	Prescriptions	Days Supply Criteria	Medication Lists
Cardiovascular, other	<ul style="list-style-type: none"> • Digoxin 	<ul style="list-style-type: none"> • >0.125 mg/day 	<ul style="list-style-type: none"> • Digoxin .05 mg per mL • Digoxin .0625 mg • Digoxin .1 mg per mL • Digoxin .125 mg • Digoxin .25 mg • Digoxin .25 mg per mL
Tertiary TCAs (as single agent or as part of combination products)	<ul style="list-style-type: none"> • Doxepin 	<ul style="list-style-type: none"> • >6 mg/day 	<ul style="list-style-type: none"> • Doxepin 3 mg • Doxepin 6 mg • Doxepin 10 mg • Doxepin 10 mg per mL • Doxepin 25 mg • Doxepin 50 mg • Doxepin 75 mg • Doxepin 100 mg • Doxepin 150 mg

High-Risk Medications Based on Prescription and Diagnosis Data

High-risk medications to avoid from the same drug class, except for appropriate diagnoses

Drug Class	Prescriptions
Antipsychotics, first (conventional) and second (atypical) generation	<ul style="list-style-type: none">• Aripiprazole• Aripiprazole lauroxil• Asenapine• Brexpiprazole• Cariprazine• Chlorpromazine• Clozapine• Fluphenazine• Haloperidol• Iloperidone• Loxapine• Lurasidone• Molindone• Olanzapine• Paliperidone• Perphenazine• Pimavanserin• Pimozide• Quetiapine• Risperidone• Thioridazine• Thiothixene• Trifluoperazine• Ziprasidone
Benzodiazepines, long, short and intermediate acting	<ul style="list-style-type: none">• Alprazolam• Chlordiazepoxide• Clobazam• Clonazepam• Clorazepate• Diazepam• Estazolam• Lorazepam• Midazolam• Oxazepam• Temazepam• Triazolam

DAE Measure Codes

Description	Codes*
Schizophrenia	ICD-10-CM: F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9
Bipolar Disorder	ICD-10-CM: F30.10-F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10-F31.13, F31.2, F31.30-F31.32, F31.4, F31.5, F31.60-F31.64, F31.70-F31.78
Other Bipolar Disorder	ICD-10-CM: F31.81, F31.89, F31.9
Alcohol Withdrawal	ICD-10-CM: F10.230-F10.232, F10.239
Benzodiazepine Withdrawal	ICD-10-CM: F13.230-F13.232, F13.239, F13.930-F13.932, F13.939
Generalized Anxiety Disorder	ICD-10-CM: F41.0, F41.1, F41.3, F41.8, F41.9
REM Sleep Behavior Disorder	ICD-10-CM: G47.52
Seizure Disorder	ICD-10-CM: G40.001, G40.009, G40.011, G40.019, G40.101, G40.109, G40.111, G40.119, G40.201, G40.209, G40.211, G40.219, G40.301, G40.309, G40.311, G40.319, G40.401, G40.409, G40.411, G40.42, G40.501, G40.509, G40.801, F40.802, G40.803, G40.804, G40.811- G40.814, G40.821-G40.824, G40.833, G40.834, G40.841-G40.844, G40.89, G40.901, G40.909, G40.911, G40.919, G40.A09, G40.A11, G40.A19, G40.B01, G40.B09, G40.C01, G40.C09, G40.C11, G40.C19

*Codes subject to change

Adherence to Cholesterol Medications (STAT) ● ●

The STAT measure evaluates the percentage of persons 18 years of age and older with the CHOL medication with a Proportion of Days Covered (PDC) \geq 80%.

PDC is calculated utilizing total days supplied of CHOL pharmacy claims/date of first RASA fill to the end of the reporting period.

Each medication claim must be submitted to the health plan (cash payment/samples/filled at out-of-network pharmacy do not count).

CHOL Medications

Prescriptions
<ul style="list-style-type: none"> • Atorvastatin (Lipitor®) • Fluvastatin (Lescol®) • Lovastatin (Mevacor®) • Pravastatin (Pravachol®) • Rosuvastatin (Crestor®) • Simvastatin (Zocor®)

Adherence to Diabetes Medications (DIAB) ●●

The DIAB measure evaluates the percentage of patients 18 years of age and older with a diabetes medication with a Proportion of Days Covered (PDC) $\geq 80\%$.

PDC is calculated utilizing total days supplied of diabetes pharmacy claims/date of first diabetes fill to the end of the reporting period.

Each medication claim must be submitted to the health plan (cash payment/samples/filled at out-of-network pharmacy do not count).

Diabetes Medications:

Category	Medication
Sulfonylureas	Glipizide and glyburide
Biguanides	Metaform
Thiazolidinediones	Actos (pioglitazone)
Alpha-glucosidase inhibitors	Precose (acarbose)
Glucagon-like peptide 1 (GLP-1) agonists	Adlyxin (lixisenatide), Byetta, Bydureon (exenatide), Ozempic (semaglutide), Tanzeum (albiglutide), Trulicity (dulaglutide) and Victoza (liraglutide)
Sodium-glucose cotransporter 2 (SGLT2) inhibitors	Farxiga (dapagliflozin), Invokana (canagliflozin) and Jardiance (empagliflozin)

Adherence to Hypertensive Medications (RASA) ●●

The RASA measure evaluates the percentage of persons 18 years of age and older with a RASA medication with a Proportion of Days Covered (PDC) $\geq 80\%$.

PDC calculated utilizing total days supplied of hypertensive pharmacy claims/date of first RASA fill to the end of the reporting period.

Each medication claim must be submitted to the health plan (cash payment/samples/filled at out-of-network pharmacy do not count).

RASA Medications

Description	Prescription
Direct Renin Inhibitor	<ul style="list-style-type: none">aliskiren (+/- hydrochlorothiazide)
ARB Medications and Combinations	<ul style="list-style-type: none">Azilsartan (+/- chlorthalidone)candesartan (+/- hydrochlorothiazide)erosartan (+/- hydrochlorothiazide)Irbesartan (+/- hydrochlorothiazide)losartan (+/- hydrochlorothiazide)olmesartan (+/- amlodipine, hydrochlorothiazide)Telmisartan (+/- amiodipine hydrochlorothiazide)valsartan (+/- amlodipine, hydrochlorothiazide nebivolol)
ACE Inhibitor Medications	<ul style="list-style-type: none">benazepril (+/- amlodipine, hydrochlorothiazide)captopril (+/- hydrochlorothiazide)enalapril (+/- hydrochlorothiazide)fosinopril (+/- hydrochlorothiazide)lisinopril (+/- hydrochlorothiazide)moexipril (+/- hydrochlorothiazide)perindopril (+/- amlodipine)quinapril (+/- hydrochlorothiazide)ramipriltrandolapril (+/- verapamil)

Concurrent Use of Opioids and Benzodiazepines (COB) ●●●●

Tip: COB is a Pharmacy Quality Alliance (PQA) quality measure. Learn more and obtain additional resources at PQAalliance.org.

The COB measure evaluates the percentage of persons 18 years of age and older with concurrent use of prescription opioids and benzodiazepines during the measurement period.

Benzodiazepine medications^{1,2}

- Alprazolam
- Chlordiazepoxide
- Clobazam
- Clonazepam
- Clorazepate
- Diazepam
- Estazolam
- Flurazepam
- Lorazepam
- Midzolam
- Oxazepam
- Quazepam
- Temazepam
- Triazolam

¹ Includes combination products

² Excludes injectable formulations

Opioid medications^{1,2}

- Benzhydrocodone
- Buprenorphine
- Butorphanol
- Codeine
- Dihydrocodeine
- Fentanyl
- Hydrocodone
- Hydromorphone
- Levorphanol
- Meperidine
- Methadone
- Morphine
- Opium
- Oxycodone
- Oxymorphone
- Pentazocine
- Tapentadol
- Tramadol

¹ Includes combination products and prescription opioid cough medications

² Excludes the following: injectable formulations; sublingual sufentanil (used in a supervised setting); and single-agent and combination buprenorphine products used to treat opioid use disorder (i.e., buprenorphine sublingual tablets, Probuphine® Implant kit subcutaneous implant, and all buprenorphine/naloxone combination products).

Polypharmacy: Use of Multiple Anticholinergic Medications in Older Adults (POLY-ACH) ●●

Tip: POLY-ACH is a Pharmacy Quality Alliance (PQA) quality measure. Learn more and obtain additional resources at PQAalliance.org.

The POLY-ACH measure evaluates the percentage of persons 65 years of age and older with concurrent use of 2 or more anticholinergic medications.

POLY-ACH anticholinergic medications^{1,2}

Description	Prescriptions
Antihistamine medications	<ul style="list-style-type: none">BrompheniramineChlorpheniramineCyproheptadineDimenhydrinate³Diphenhydramine (oral)DoxylamineHydroxyzineMeclizineTriprolidine
Antiparkinsonian agent medications	<ul style="list-style-type: none">BenztropineTrihexyphenidyl
Skeletal muscle relaxant medications	<ul style="list-style-type: none">CyclobenzaprineOrphenadrine
Antidepressant medications	<ul style="list-style-type: none">AmitriptylineAmoxapineClomipramineDesipramineDoxepin (>6 mg/day)⁴ImipramineNortriptylineParoxetine
Antipsychotic medications	<ul style="list-style-type: none">ChlorpromazineClozapineOlanzapinePerphenazineDarifenacinFesoterodineFlavoxateOxybutyninSolifenacinTolterodineTrospium

POLY-ACH anticholinergic medications (continued)

Description	Prescriptions
Antimuscarinic (urinary incontinence) medications	<ul style="list-style-type: none"> • Darifenacin • Fesoterodine • Flavoxate • Oxybutynin • Solifenacin • Tolterodine • Trospium
Antispasmodic medications	<ul style="list-style-type: none"> • Atropine² • Clidinium-chlordiazepoxide⁵ • Dicyclomine • Homatropine² • Hyoscyamine • Scopolamine²
Antiemetic medications	<ul style="list-style-type: none"> • Prochlorperazine • Promethazine

¹ Includes combination products and prescription opioid cough medications

² Excludes ophthalmics, injectable formulations, sublingual sufentanil, and buprenorphine products used to treat opioid use disorder

³ There are no active NDCs for dimenhydrinate

⁴ Includes doxepin when the daily dose is >6 mg/day

⁵ Chlordiazepoxide is not a target medication as a single drug

Statin Therapy for Patients with Cardiovascular Disease (SPC-E)

Beginning in MY 2026, this measure transitioned to the Electronic Clinical Data Systems (ECDS) reporting method only.

The SPC-E measure evaluates males 21 to 75 years of age and females 40 to 75 years of age who were identified as having Clinical Atherosclerotic Cardiovascular Disease (ASCVD).

Two rates are reported:

- **Received Statin Therapy:** Persons who were dispensed at least one high-intensity or moderate-intensity statin medication during the measurement period
- **Statin Adherence:** Persons who remained on a high-intensity or moderate-intensity statin medication for at least 80% of the treatment period

SPC-E Medications

Description	Prescription
High-intensity statin therapy	<ul style="list-style-type: none">• Atorvastatin 40-80 mg
High-intensity statin therapy	<ul style="list-style-type: none">• Amlodipine-atorvastatin 40-80 mg
High-intensity statin therapy	<ul style="list-style-type: none">• Rosuvastatin 20-40 mg
High-intensity statin therapy	<ul style="list-style-type: none">• Simvastatin 80 mg
High-intensity statin therapy	<ul style="list-style-type: none">• Ezetimibe-simvastatin 80 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Atorvastatin 10-20 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Amlodipine-atorvastatin 10-20 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Rosuvastatin 5-10 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Simvastatin 20-40 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Ezetimibe-simvastatin 20-40 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Pravastatin 40-80 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Lovastatin 40 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Fluvastatin 40-80 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Pitavastatin 1-4 mg

Statin Therapy for Patients with Diabetes (SPD-E) ●●●●●

Beginning in MY 2026, this measure transitioned to the Electronic Clinical Data Systems (ECDS) reporting method only.

The SPD-E measure evaluates persons 40 to 75 years of age with diabetes (Types 1 and 2) who do not have clinical atherosclerotic cardiovascular disease (ASCVD).

Two rates are reported:

- **Received Statin Therapy:** Persons who were dispensed at least one statin medication of any intensity during the measurement period
- **Statin Adherence 80%:** Persons who remained on a statin medication of any intensity for at least 80% of the treatment period

SPD Medications: High, Moderate, and Low Intensity Statin Medications

Description	Prescription
Amlodipine Atorvastatin High Intensity Medications	amlodipine/atorvastatin
Amlodipine Atorvastatin Moderate Intensity Medications	amlodipine/atorvastatin
Atorvastatin High Intensity Medications	atorvastatin
Ezetimibe Simvastatin High Intensity Medications	ezetimibe/simvastatin
Fluvastatin Moderate & Low Intensity Medications	fluvastatin
Lovastatin Moderate & Low Intensity Medications	lovastatin
Pitavastatin Moderate Intensity Medications	pitavastatin
Pravastatin Moderate & Low Intensity Medications	pravastatin
Rosuvastatin High & Moderate Intensity Medications	rosuvastatin
Simvastatin High, Moderate, & Low Intensity Medications	simvastatin

Women's Health



Breast Cancer Screening (BCS-E)

The BCS-E measure evaluates the percentage of persons 40 to 74 years of age who were recommended for routine breast cancer screening and had a mammogram any time on or between October 1 two years prior to the measurement period and the end of the measurement period.

Persons recommended for routine breast cancer screening include:

- Administrative Gender of Female at any time in the patient's history
- Sex Assigned at Birth of Female at any time in the patient's history
- Sex Parameter for Clinical Use of Female during the measurement period

BCS-E Measure Codes

Description	Codes*
Mammography	CPT: 77061-77063, 77065-77067 LOINC: 24604-1, 24605-8, 24606-6, 24610-8, 26175-0, 26176-8, 26177-6, 26287-3, 26289-9, 26291-5, 26346-7, 26347-5, 26348-3, 26349-1, 26350-9, 26351-7, 36319-2, 36625-2, 36626-0, 36627-8, 36642-7, 36962-9, 37005-6, 37006-4, 37016-3, 37017-1, 37028-8, 37029-6, 37030-4, 37037-9, 37038-7, 37052-8, 37053-6, 37539-4, 37542-8, 37543-6, 37551-9, 37552-7, 37553-5, 37554-3, 37768-9, 37769-7, 37770-5, 37771-3, 37772-1, 37773-9, 37774-7, 37775-4, 38070-9, 38071-7, 38072-5, 38090-7, 38091-5, 38807-4, 38820-7, 38854-6, 38855-3, 39150-8, 39152-4, 39153-2, 39154-0, 42168-5, 42169-3, 42174-3, 42415-0, 42416-8, 46335-6, 46336-4, 46337-2, 46338-0, 46339-8, 46342-2, 46350-5, 46351-3, 46354-7, 46355-4, 46356-2, 46380-2, 48475-8, 48492-3, 69150-1, 69251-7, 69259-0, 72137-3, 72138-1, 72139-9, 72140-7, 72141-5, 72142-3, 86462-9, 86463-7, 91517-3, 91518-1, 91519-9, 91520-7, 91521-5, 91522-3

*Codes subject to change

Documented Assessment After Mammogram

(DBM-E) ●●●●●

DBM-E evaluates the percentage of episodes of mammograms in the form of a BI-RADS assessment within 14 days of the mammogram for persons 40-74 years of age.

DBM-E Measure Codes

Description	Codes*
BI-RADS Assessment	SNOMED CT: 397138000, 397140005, 397141009, 397143007, 397144001, 397145000, 6111000179101, 6121000179106, 6131000179108, 6141000179100
Mammography	CPT: 77061-77063, 77065-77067 LOINC: 24604-1, 24605-8, 24606-6, 24610-8, 26175-0, 26176-8, 26177-6, 26287-3, 26289-9, 26291-5, 26346-7, 26347-5, 26348-3, 26349-1, 26350-9, 26351-7, 36319-2, 36625-2, 36626-0, 36627-8, 36642-7, 36962-9, 37005-6, 37006-4, 37016-3, 37017-1, 37028-8, 37029-6, 37030-4, 37037-9, 37038-7, 37052-8, 37053-6, 37539-4, 37542-8, 37543-6, 37551-9, 37552-7, 37553-5, 37554-3, 37768-9, 37769-7, 37770-5, 37771-3, 37772-1, 37773-9, 37774-7, 37775-4, 38070-9, 38071-7, 38072-5, 38090-7, 38091-5, 38807-4, 38820-7, 38854-6, 38855-3, 39150-8, 39152-4, 39153-2, 39154-0, 42168-5, 42169-3, 42174-3, 42415-0, 42416-8, 46335-6, 46336-4, 46337-2, 46338-0, 46339-8, 46342-2, 46350-5, 46351-3, 46354-7, 46355-4, 46356-2, 46380-2, 48475-8, 48492-3, 69150-1, 69251-7, 69259-0, 72137-3, 72138-1, 72139-9, 72140-7, 72141-5, 72142-3, 86462-9, 86463-7, 91517-3, 91518-1, 91519-9, 91520-7, 91521-5, 91522-3

*Codes subject to change

Follow-Up After Abnormal Mammogram Assessment (FMA-E)

FMA-E evaluates the percentage of episodes for persons 40–74 years of age with inconclusive or high-risk BI-RADS assessments that received appropriate follow-up within 90 days of the assessment.

FMA-E Measure Codes

Description	Codes*
Breast Biopsy	CPT: 19081, 19083, 19085, 19100, 19101 SNOMED CT: 10940003, 116219004, 116220005, 116334007, 1179705005, 1179707002, 1179705005, 117970700, 1179708007, 12131000087109, 1220570007, 1220571006, 1220572004, 1264555004, 1264556003, 1264555004, 1264556003, 1268323005, 1268996004, 1332066007, 1332067003, 1333891002, 1333892009, 1333893004, 1333894005, 1333895006, 1333896007, 1333897003, 1333898008, 1333899000, 1333900005, 1333901009, 1333902002, 1333903007, 1333904001, 1333905000, 1333906004, 1334078005, 1334079002, 1334080004, 1356791009, 172086006, 2131000087106, 2141000087100, 237372000, 237375003, 237376002, 237377006, 237378001, 237379009, 265253005, 274331003, 2841000087108, 287553003, 28768007, 303689004, 305051000000107, 305071000000103, 306371000000109, 306651000000105, 306671000000101, 308041000000102, 305011000000108, 306381000000106, 306641000000107, 307298009, 307971000000105, 307981000000107, 387736007, 42125001, 432109009, 432157003, 432337008, 432550005, 433008009, 433685008, 433805008, 442963006, 445171002, 445437001, 44578009, 448336005, 448689003, 4541000087104, 4551000087101, 5181000087103, 709628007, 711508007, 723990008, 725936002, 736615002, 770568001, 770569009, 770570005, 771086002, 771625002, 785800009, 786883001, 866232001, 872731000000104
Breast Ultrasound	CPT: 76641, 76642 LOINC: 24599-3, 24601-7, 26215-4, 26216-2, 26288-1, 26290-7, 42132-1, 105420-4, 105421-2
High Risk BI-RADS	SNOMED CT: 397144001, 397145000, 6121000179106, 6131000179108, 6141000179100
Inconclusive BI-RADS	SNOMED CT: 397138000

FMA-E Measure Codes (continued)

Description	Codes*
Mammography	CPT: 77061-77063, 77065-77067 LOINC: 24604-1, 24605-8, 24606-6, 24610-8, 26175-0, 26176-8, 26177-6, 26287-3, 26289-9, 26291-5, 26346-7, 26347-5, 26348-3, 26349-1, 26350-9, 26351-7, 36319-2, 36625-2, 36626-0, 36627-8, 36642-7, 36962-9, 37005-6, 37006-4, 37016-3, 37017-1, 37028-8, 37029-6, 37030-4, 37037-9, 37038-7, 37052-8, 37053-6, 37539-4, 37542-8, 37543-6, 37551-9, 37552-7, 37553-5, 37554-3, 37768-9, 37769-7, 37770-5, 37771-3, 37772-1, 37773-9, 37774-7, 37775-4, 38070-9, 38071-7, 38072-5, 38090-7, 38091-5, 38807-4, 38820-7, 38854-6, 38855-3, 39150-8, 39152-4, 39153-2, 39154-0, 42168-5, 42169-3, 42174-3, 42415-0, 42416-8, 46335-6, 46336-4, 46337-2, 46338-0, 46339-8, 46342-2, 46350-5, 46351-3, 46354-7, 46355-4, 46356-2, 46380-2, 48475-8, 48492-3, 69150-1, 69251-7, 69259-0, 72137-3, 72138-1, 72139-9, 72140-7, 72141-5, 72142-3, 86462-9, 86463-7, 91517-3, 91518-1, 91519-9, 91520-7, 91521-5, 91522-3

*Codes subject to change

Cervical Cancer Screening (CCS-E) ●●●

The CCS-E measure evaluates the percentage of persons 21 to 64 years of age who were recommended for routine cervical cancer screening and screened for cervical cancer using either of the following criteria:

- Persons 21 to 64 years of age who had cervical cytology performed within the last 3 years
- Persons 30 to 64 years of age who had cervical high-risk human papillomavirus (hrHPV) testing performed within the last 5 years
- Persons 30 to 64 years of age who had cervical cytology/high risk human papillomavirus (hrHPV) co-testing within the last 5 years

CCS-E Measure Codes

Description	Codes*
Cervical Cytology Lab Test (20-64)	CPT: 88141-88143, 88147, 88148, 88150, 88152-88153, 88164-88167, 88174, 88175 HCPCS: G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091 LOINC: 10524-7, 18500-9, 19762-4, 19764-0, 19765-7, 19766-5, 19774-9, 33717-0, 47527-7, 47528-5
Cervical Cytology Result or Finding	SNOMED CT: 168406009, 168407000, 168408005, 168410007, 168414003, 168415002, 168416001, 168424006, 250538001, 269957009, 269958004, 269959007, 269960002, 269961003, 269963000, 275805003, 281101005, 309081009, 310841002, 310842009, 416030007, 416032004, 416033009, 439074000, 439776006, 439888000, 441087007, 441088002, 441094005, 441219009, 441667007, 700399008, 700400001, 1155766001, 62051000119105, 62061000119107, 98791000119102
High-Risk HPV Lab Tests (30-64)	CPT: 87624, 87625, 87626, 0502U HCPCS: G0476 LOINC: 104132-6, 104170-6, 104752-1, 104766-1, 104783-6, 21440-3, 30167-1, 38372-9, 59263-4, 59264-2, 59420-0, 69002-4, 71431-1, 75694-0, 77379-6, 77399-4, 77400-0, 82354-2, 82456-5, 82675-0, 95539-3

*Codes subject to change

Chlamydia Screening (CHL) ●●

The CHL measure evaluates the percentage of persons 16 to 24 years of age who were recommended for chlamydia screening, identified as sexually active, and who had at least one test for chlamydia.

Meridian covers all types of chlamydia screenings. This includes traditional methods, as well as urine screening (bill with CPT code 87110) for men and women. The advantage to urine screening is that it is simple, quick to administer, and has a higher accuracy rate than other methods.

CHL Measure Codes

CPT*

87110, 87270, 87320, 87490-87492, 87810

*Codes subject to change

Osteoporosis Management in Women Who Had a Fracture (OMW) ●●

The OMW measure evaluates the percentage of women 67 to 85 years of age who suffered a fracture and had either a bone mineral density (BMD) test or a prescription for a drug to treat osteoporosis in the six months after the fracture.

OMW Measure Codes

Description	Codes*
Bone Mineral Density Tests	CPT: 76977, 77080, 77081, 77085, 77086 ICD10PSC: BP48ZZ1, BP49ZZ1, BP4GZZ1, BP4HZZ1, BP4LZZ1, BP4MZZ1, BP4NZZ1, BP4PZZ1, BQ00ZZ1, BQ01ZZ1, BQ03ZZ1, BQ04ZZ1, BR00ZZ1, BR07ZZ1, BR09ZZ1, BR0GZZ1 LOINC: 100225-2, 101804-3, 101805-0, 104938-6, 24701-5, 24890-6, 24966-4, 38261-4, 38262-2, 38623-0, 38264-8, 38265-5, 38266-3, 38267-1, 46278-8, 46279-6, 46383-6, 80932-7, 80933-5, 80934-3, 80935-0, 80936-8, 80937-6, 80938-4, 80939-2, 80940-0, 80941-8, 80942-6, 80943-4, 80944-2, 80945-9, 80946-7, 80947-5, 80948-3, 80949-1, 80950-9, 80951-7, 80952-5, 80953-3, 80954-1, 80955-8, 80956-6, 83311-1, 85385-3, 85386-1, 85387-9, 85388-7, 85389-5, 85390-3, 85391-1, 85392-9, 85393-7, 85394-5
Osteoporosis Medications Therapy	HCPCS: J0897, J1740, J3110, J3111, J3489
Long-Acting Osteoporosis Medications during Inpatient Stay	HCPCS: J0897, J1740, J3489

*Codes subject to change

Osteoporosis Medications

Prescriptions
<ul style="list-style-type: none">• Abaloparatide• Alendronate• Alendronate-cholecalciferol• Denosumab• Ibandronate• Raloxifene• Risedronate• Romosozumab• Teriparatide• Zoledronic acid

Osteoporosis Screening in Older Women (OSW) ●●

The OSW measure evaluates the percentage of women 65 to 75 years of age who received osteoporosis screening.

OSW Measure Codes

Description	Codes*
Osteoporosis Screening Tests	CPT: 76977, 77078, 77080, 77081, 77085

*Codes subject to change

Prenatal and Postpartum Care (PPC) ●●

The PPC measure evaluates the percentage of deliveries of live births on or between October 8 of the year prior to the measurement period and October 7 of the measurement period. For these persons, the measure assesses the following facets of prenatal and postpartum care.

Timeliness of Prenatal Care: Percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date, or within 42 days of enrollment in the organization

Postpartum Care: Percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery

PPC Measure Codes

Prenatal:

Description	Codes*
Prenatal Visits with <i>Pregnancy Diagnosis</i> (Visit must be performed in the first trimester [13 weeks], on or before the enrollment start date, or within 42 days of enrollment if already pregnant at the time of enrollment with Meridian)	CPT: 98966-98968, 98970-98972, 98980-98981, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99421-99423, 99441-99443, 99457, 99458, 99483 HCPCS: G0463, T1015, G0071, G2010, G2012, G2250, G2251, G2252
Stand-Alone Prenatal Visits	CPT: 99500 CPT-CAT-II: 0500F, 0501F, 0502F HCPCS: H1000, H1001, H1002, H1003, H1004
Prenatal Bundled Services	CPT: 59400, 59425, 59426, 59510, 59610, 59618 HCPCS: H1005

PPC Measure Codes (continued)

Postpartum:

Description	Codes*
Cervical Cytology Lab Test	CPT: 88141-88143, 88147, 88148, 88150, 88152-88153, 88164-88167, 88174, 88175 HCPCS: G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091
Encounter for Postpartum Care (must be on or between 7 days and 84 days after delivery)	ICD-10: Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2
Postpartum Bundled Services	CPT: 59400, 59410, 59510, 59515, 59610, 59614, 59618, 59622
Postpartum Care	CPT: 57170, 58300, 59430, 99501 CPT-CAT II: 0503F HCPCS: G0101

*Codes subject to change

NOTE: When using the Prenatal Visit codes, remember to also include a Pregnancy Diagnosis code.

For information about Global Maternity Billing, please refer to the HFS website (hfs.illinois.gov) and the Illinois Association of Medicaid Health Plans (IAMHP) Comprehensive Billing Manual at iamhp.org/providers.

Prenatal Immunization Status (PRS-E) ● ☁

PRS-E evaluates the percentage of deliveries in the measurement period in which persons had received influenza and tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccinations.

Influenza:

- Persons who received an adult influenza vaccine on or between July 1 of the year prior to the measurement period and the delivery date

Tdap:

- Persons who received at least one Tdap vaccine during the pregnancy (including on the delivery date)

PRS-E Codes

Description	Codes*
Adult Influenza Immunization	CVX: 88, 135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205
Adult Influenza Vaccine Procedure	CPT: 90653, 90656, 90658, 90661, 90662, 90673, 90674, 90682, 90686, 90688, 90689, 90694, 90756
Deliveries where persons had anaphylaxis due to the influenza vaccine on or before the delivery date	SNOMED CT: 471361000124100
Tdap Vaccine Procedure	CVX: 115 CPT: 90715 SNOMED CT: 390846000, 412755006, 412756007, 412757003, 428251000124104, 571571000119105
Anaphylaxis due to Diphtheria, Tetanus, or Pertussis Vaccine	SNOMED CT: 428281000124107, 428291000124105
Encephalitis due to Diphtheria, Tetanus or Pertussis Vaccine	SNOMED CT: 192710009, 192711008, 192712001

*Codes subject to change

The Advisory Committee on Immunization Practices (ACIP) clinical guidelines recommend that all women who are pregnant or who might be pregnant in the upcoming influenza season receive inactivated influenza vaccines. ACIP also recommends that pregnant women receive one dose of Tdap during each pregnancy, preferably during the early part of gestational weeks 27–36, regardless of their prior history of receiving Tdap.

Prenatal Depression Screening and Follow-Up (PND-E)

PND-E evaluates the percentage of deliveries in which persons were screened for clinical depression while pregnant using a standardized instrument and, if screened positive, received follow-up care within 30 days.

- **Deliveries between January 1 and December 1 of the measurement period:** Screening should be performed between the pregnancy start date and the delivery date (including on the delivery date)
- **Deliveries between December 2 and December 31 of the measurement period:** Screening should be performed between the pregnancy start date and December 1 of the measurement period

* Refer to the Appendix for a list of Approved Depression Screening Instruments, Codes, and Positive Findings

**Refer to the Appendix for a list of follow-up visit codes

Postpartum Depression Screening and Follow-Up (PDS-E)

PDS-E evaluates the percentage of deliveries in which persons were screened for clinical depression during the postpartum period (7 to 84 days following delivery), using a standardized instrument and, if screened positive, received follow-up care within 30 days.

* Refer to the Appendix for a list of Approved Depression Screening Instruments, Codes, and Positive Findings

**Refer to the Appendix for a list of follow-up visit codes

Meridian Quality Improvement Resources



Consumer Assessment of Healthcare Providers and Systems (CAHPS®) ● ● ● ●

Every year, a random sample of patients are surveyed about their experience with their providers, services, and health plan. It is an important component of ensuring that patients are satisfied, not only with their health outcomes but also with their healthcare experience.

The CAHPS survey is applicable to **Meridian**, **Wellcare**, and **YouthCare**. For **Ambetter**, the CAHPS survey is referred to as the Qualified Health Plan (QHP) Enrollee Survey. The CAHPS and the QHP surveys allow patients to evaluate the aspects of care delivery that matter the most to them. At Meridian, we are committed to partnering with our providers to deliver outstanding patient experience.

As a provider, you are the most critical component of that experience. We want to ensure that you know exactly how your patients are evaluating your care. Please take a moment to review and to familiarize yourself with some of the key topics included in the survey.

Survey Measure & Description	Daily Practice Tips
<p>Getting Needed Care: This measure assesses the ease with which patients received the care, tests, or treatment they needed. It also assesses how often they were able to get a specialist appointment scheduled when needed.</p>	<ul style="list-style-type: none"> • Office staff should help coordinate specialty appointments for urgent cases • Encourage patients and caregivers to view results on the patient portal when available • Inform patients of what to do if care is needed after hours • Offer appointments or refills via text and/or email • Offer alternative appointment types to expand access to care (e.g., telephone, telehealth, telemedicine, and patient portals)
<p>Getting Care Quickly: This measure assesses how often patients got the care they needed as soon as they needed it and how often appointment wait times exceeded 15 minutes.</p>	<ul style="list-style-type: none"> • Ensure a few appointments each day are available to accommodate urgent visits • Offer appointments with a nurse practitioner or physician assistant for short notice appointments • Maintain an effective triage system to ensure that frail and/or very sick patients are seen right away or provided alternate care via phone and urgent care • Keep patients informed if there is a longer wait time than expected, and give them an option to reschedule
<p>Care Coordination: This measure assesses providers' assistance with managing the disparate and confusing health care system, including access to medical records, timely follow-up on test results, and education on prescription medications.</p>	<ul style="list-style-type: none"> • Ensure there are open appointments for patients recently discharged from a facility • Integrate PCP and specialty practices through EMR or fax to get reports promptly • Ask patients if they have seen any other providers; discuss visits to specialty care as needed • Encourage patients to bring in their medications to each visit • Share test results and explain what they mean • Provide instructions for follow-up questions or concerns

Survey Measure & Description	Daily Practice Tips
<p>How Well Doctors Communicate: This measure assesses patients' perception of the quality of communication with their doctor. Consider using the Teach-Back Method to ensure patients understand their health information.</p>	<ul style="list-style-type: none"> • Use body language to show you are engaged and listening to the patient (sit down and face the patient; maintain eye contact) • Don't rush; explain things in a way the patient can understand • Be empathetic with bad news. Be sensitive and acknowledge family members or caregivers in the room with the patient • Summarize the visit and next steps in a way the patient can understand • Let the patient know when you will see them next <p>What is Teach-back?</p> <ul style="list-style-type: none"> • Asking a patient (or family member) to explain in their own words what they need to know or do, in a caring way • A way to check for understanding and, if needed, re-explain and check again
<p>Rating of Health Care Quality: The CAHPS® survey asks patients to rate the overall quality of their health care on a 0-10 scale.</p>	<ul style="list-style-type: none"> • Encourage patients to make their routine appointments for checkups or follow up visits as soon as they can – weeks or even months in advance • Ensure that open care gaps are addressed during each patient visit • Make use of the provider portal when requesting prior authorizations

On the following page are examples of **satisfaction categories and survey questions** for which your patients are asked to respond; provider discussion questions; and Provider Tips. We hope this tool will provide reinforcement opportunities for your relationship with the patients you serve.

Sample Questions	Provider Tips
<p>Health Promotion Discussion Questions</p> <p>Any problems with your work or daily activities due to physical problems?</p> <p>Any problems with your work or daily activities due to stress?</p> <p>Anything bothering you or stressful?</p> <p>Are you sad or depressed?</p> <p>Do you use tobacco? (Always/Sometimes/Never)</p> <p>Do you drink alcohol? (Always/Sometimes/Never)</p> <p>Do you exercise? (Always/Sometimes/Never)</p> <p>Do you take aspirin? (Always/Sometimes/Never)</p> <p>Do you or anyone in your family have high blood pressure, high cholesterol or had a heart attack?</p> <p>Have you had a flu shot in the past calendar year? If not, Why?</p>	<ul style="list-style-type: none"> • Complete and document any health assessment on patient • Discuss with patient the benefits of exercise and encourage them to start, increase or maintain physical activity and document discussion • Discuss the risks of tobacco use and recommend medication to assist in stopping • Discuss issues associated with drinking too much alcohol, if necessary • Discuss the risks and benefits of aspirin to prevent heart attack or stroke • Screen patient for high blood pressure and cholesterol • Recommend and/or administer the flu shot during flu season
<p>Medication Discussion Questions</p> <p>Are you currently on any prescription medications from another doctor? If so, what?</p> <p>How long have you been on the medication?</p>	<ul style="list-style-type: none"> • Document all prescription medication patient is taking • Discuss options and reasons to take alternate medications if patient is not getting positive results for symptoms • Discuss reasons with patient why they may need to stop taking a particular medication • Discuss the benefits and risks of taking a medicine • Discuss patient's preference on what medication they feel would be best for them • Review medications prescribed by PCP and specialists and verify results

Sample Questions	Provider Tips
<p>Access to Care Discussion Questions</p> <p>Are you satisfied with the timeframe it took to schedule your appointment?</p> <p>Were you able to get your appointment as soon as you needed?</p> <p>Are you satisfied with the coordination of care you receive, coordinating visits with specialists, non-emergency transportation (if needed) and providing lab or test results?</p>	<ul style="list-style-type: none"> Evaluate office procedures to improve getting patients scheduled as quickly as possible for their symptoms Determine why patient perceives difficulty in getting timely care, if necessary Educate patient on timeframes for getting appointments according to their symptoms Assist in coordination of non-emergency transportation, if necessary Use patient experience surveys or post-visit survey results to identify opportunities for improvement

CAHPS® Outpatient Mental Health Survey Information



Annually, NCQA directs health plans to conduct a survey about their members' experience with behavioral health services. The Outpatient Mental Health Survey measures members' experiences and identifies opportunities for health plans and providers to improve quality of care and access to mental health and substance use services.

Composite Measure	Sample Question
Getting Appointments for Prescription Medications	<p>How difficult was it for you to make an appointment with the person who prescribes your mental health medicine?</p> <p><i>(Very difficult, Somewhat difficult, Not very difficult, Not difficult at all)</i></p>
Getting Mental Health Counseling	<p>How difficult was it for you to make an appointment with your mental health counselor?</p> <p><i>(Very difficult, Somewhat difficult, Not very difficult, Not difficult at all)</i></p>
Communication with Mental Health Counselor	<p>How often did your main mental health counselor listen carefully to you?</p> <p>How often did your main mental health counselor show respect for what you had to say?</p> <p><i>(Never, Sometimes, Usually, Always)</i></p>

Composite Measure	Sample Question
Goal Setting	How much did your main mental health counselor consider what is important to you when setting the goals for treatment? <i>(Not at all, A little, Some, A lot)</i>
Getting Help Between Appointments	When you contacted your main mental health counselor between your appointments, how often did you get the help or support you needed? <i>(Never, Sometimes, Usually, Always)</i>
Rating of Mental Health Counselor	What number would you use to rate your main mental health counselor? <i>(0-10)</i>
Unmet Need for Mental Health Services	Did you get all the mental health services you needed? <i>(Yes, No)</i>
Financial Barriers to Mental Health Services	How difficult was it for you to pay for the mental health services you received? <i>(Very difficult, Somewhat difficult, Not very difficult, Not difficult at all)</i>

Provider Tips

- Offer extended hours, telehealth, and various treatment options when possible.
- Let patients know your office hours and how to get after-hour care.
- Partner with the health plan and coordinate care with other specialists and primary care physicians to address whole-person health and access to timely care.
- Assess cultural and linguistic needs, and ask your patients what is important to them.
- Obtain release of information forms and explain the purpose of releasing information to other providers.
- Include family/caregivers/identified support in the treatment plan.
- Invite questions and encourage your patient to take notes.
- Use the “teach-back” method.

Health Outcomes Survey (HOS)

The Medicare Health Outcomes Survey (HOS) measures a health plan's success in improving and maintaining the functional status of members ages 65 and older. HOS is an annual survey that is conducted from July through November with a random sample of Medicare patients. The same patients are surveyed again two years later to assess changes in health status. HOS measures patients' perception of their physical & mental health and overall quality of life. **HOS results impact Centers for Medicare & Medicaid Services (CMS) Star Ratings.**

HOS is one of two annual member surveys that monitor patient/member satisfaction that can affect CMS Star Ratings. There are five HOS measures – two functional health measures and three HEDIS® Effectiveness of Care measures in the annual Medicare Part C Star Ratings. For more information, visit the Medicare HOS website at hosonline.org/en/.

The Five HOS Measures

HOS Measure	Survey Question
Improving or Maintaining Physical Health (functional health measure)	"During the past four weeks, have you accomplished less than you would like or been limited in your work or other regular daily activities as a result of your physical health?"
Improving or Maintaining Mental Health (functional health measure)	"During the past four weeks, how much of the time has your physical health or emotional problems interfered with your social activities?"
Monitoring Physical Activity (HEDIS Effectiveness of Care measure)	"In the past 12 months, did you talk with a doctor or other health provider about your level of exercise or physical activity?"
Improving Bladder Control (HEDIS Effectiveness of Care measure)	"Many people experience leakage of urine, also called urinary incontinence. In the past six months have you experienced leaking of urine?"
Reducing the Risk of Falling (HEDIS Effectiveness of Care measure)	"Has your doctor or other health provider done anything to help prevent falls or treat problems with balance or walking?"

Ways Providers Can Impact HOS Outcomes

- Ask questions
- Remember each HOS measure addresses a different aspect of patient care and patient-provider interaction
- Encourage patients to take actions aligning with the HOS measures
- Encourage your office staff to help patients fill out the HOS (HOS is administered by phone and mail)

HOS Measure 1: Improving or maintaining physical health

This measure assesses the percentage of persons whose physical health was the same or better after two years.

Recommendations	Discussion questions
<ul style="list-style-type: none">Assess your patients' pain and functional status using standardized toolsProvide interventions to improve physical health, such as disease management, pain management, physical therapy, or care managementPromote self-management support strategies, such as goal-setting, action planning, problem solving, and follow-up to help patients take an active role in improving their health	<ul style="list-style-type: none">How far can you walk?Are you able to shop and cook your own food?Do you have trouble with stairs?Does pain limit your activities?

HOS Measure 2: Improving or maintaining mental health

This measure assesses the percentage of patients whose behavioral health was the same or better after two years.

Recommendations	Discussion questions
<ul style="list-style-type: none">Assess your patients' symptoms of depression with the PHQ-2 and, when appropriate, PHQ-9Refer patients to behavioral health services or manage depression and anxiety treatment as indicatedPromote web-based programs, such as mystrength.com, that provide a range of evidence-based behavioral health self-care resourcesUse motivational interviewing to improve treatment engagement and behavioral and physical health outcomes	<ul style="list-style-type: none">How is your energy throughout the day?What do you like to do to socialize?Does alcohol ever get in the way of other important things in life?

HOS Measure 3: Monitoring physical activity

This measure assesses the percentage of patients who discussed exercise with their doctor or other health care provider and were advised to start, increase, or maintain their physical activity within the year.

Recommendations	Discussion questions
<ul style="list-style-type: none">• Assess your patients' current physical activity level• Discuss health benefits and advise patients to start, maintain, or increase physical activity as appropriate for their individual health status• Develop physical activity plans with patients that match their abilities.• Refer patients with limited mobility to physical therapy to learn safe and effective exercises• Encourage participation in a gym, fitness and exercise programs, and local community resources	<ul style="list-style-type: none">• What does your exercise routine look like?• What activities do you enjoy?• Do you feel better when you are more active?

HOS Measure 4: Improving bladder control

This measure assesses the percentage of patients with urinary incontinence (UI) who discussed the problem and treatment options with their health care providers.

Recommendations	Discussion questions
<ul style="list-style-type: none">• Assess concerns with UI in the last six months and document discussion on the My Wellness and Prevention Checklist• Assess the severity of the condition and the impact of UI on patients' quality of life. Involve patients in decisions about treatment options that work best for them. These options include behavioral (such as bladder training and pelvic muscle rehabilitation), pharmacological, and surgical therapies• Have informative brochures and materials visible and available as discussion starters	<ul style="list-style-type: none">• Have you had leakage in the past six months?• How often have you had leakage and when does it occur?• Has leakage affected your daily life (such as social withdrawals, depression, or sleep deprivation)?

HOS Measure 5: Reducing risk of falling

This measure assesses the percentage of patients with falling, walking, or balance problems who discussed these topics with their providers and received treatment within the year.

Recommendations	Discussion questions
<ul style="list-style-type: none">• Assess fall risk by asking patients about falling, gait, and balance problems• Provide fall prevention interventions, such as promoting regular exercise, strengthening and balance activities (tai chi, yoga), promoting regular eye exams, and providing appropriate educational materials• Promote home safety, such as:<ul style="list-style-type: none">– Removal of throw rugs and clutter to reduce tripping– Installing handrails on stairs and grab bars in the bathroom• Use of non-slip mats in the tub or shower• Use of night lights to keep halls well lit	<ul style="list-style-type: none">• Have you had a fall in the past year?• Can you tell me about the circumstances of the fall?• Do you think the fall could have been prevented?• Have you felt dizzy, or had problems with balance or walking in the past year?• Have you had any vision problems?• Have you had a recent eye exam?

Critical Incidents (CI) ● ●

A critical incident (CI) is any alleged or actual event that poses a risk of serious harm, injury, or death of the member.

Critical incidents include, but are not limited to:

• Abuse	• Medication Management
• Fraud	• Restraint/Seclusion/or Other
• Neglect	Restrictive Intervention
• Exploitation	• Medical Emergency/Injury/Illness
• Behavioral Health Issues	• Missing Person/Elopement
• Unanticipated Death	• Environmental/Unsafe
• Legal/Criminal Activity	Housing/Displacement

Types of abuse can include physical abuse, verbal abuse, sexual abuse, or harassment, and mental or emotional abuse.

Reporting Critical Incidents

1. Identify the appropriate CI type
2. Complete a Critical Incident Reporting Form if incident occurred within the past year
 - Submit within 24 hours of discovering the incident
3. Email Critical Incident Reporting Form to criticalincidents@mhplan.com

The Critical Incident Reporting Form can be found here:

ILmeridian.com/providers/resources/forms-resources.html.

Cultural Competence ●●●

Cultural Competence is a set of attitudes, behaviors, and policies that enable people to work effectively in cross-cultural situations. We serve a diverse patient population. The ability to understand and relate to different cultures can help you communicate effectively with your patients. All Meridian network providers are contractually required to complete the online Cultural Competency training annually.

Access the training material on our website at ILmeridian.com/providers/resources/provider-training/annual-training.html

Tips for Providing Culturally Competent Care

Consider population-specific conditions: Low-income/low-literacy, education, disability, race, ethnicity, religion, spirituality, age, gender identity, and urbanicity.

Ask about cultural practices: Spiritual traditions, dietary restrictions, and more may impact a patient's clinical experience.

Practice transcultural techniques: Approach a new patient slowly, be respectful, sit in a quiet setting, and sit a comfortable distance away.

Ensure patient's understanding of care: Lack of accessible medication instructions in a patient's language can impact quality of care. Ensure patient comprehension by providing translated educational materials and utilizing qualified medical interpreters when necessary.

Things to Remember

- 1 in 4 Americans live with a disability and are twice as likely to find his or her provider's skills or facilities inadequate.
- 1 in 5 Americans speak a language other than English at home. Language barriers can prevent patients from effectively conveying their ailments and understanding their care plans.
- 3.6 million Americans miss or delay medical care because they lack reliable transportation.
- Invest in Americans with Disabilities Act (ADA)-approved renovations and train staff on disabilities, challenges, and rights.
- Speak slowly, summarize, demonstrate, and use appropriate terminology when providing instructions. Ensure that patients understand the instructions at the end of the visit.
- Call Meridian's transportation service at **866-796-1165** at least three business days prior to a patient's appointment.

Sources: CDC.gov, census.gov, ncbi.nlm.nih.gov

Social Determinants of Health (SDOH) ● ● ● ●

Social determinants of health are the everyday factors that can impact how healthy a person is regardless of their clinical care. These everyday factors include where a person lives, grows, plays, works, and worships which directly impact financial stability, access to adequate education, access to healthcare, community infrastructure, and social and community context. Addressing SDOH improves medication adherence, supports adherence to treatment plans, and reduces avoidable healthcare utilization.

What can you do?

- Screen for SDOH during all patient visits. Consider using a standard SDOH screener such as PRAPARE (find eligible screening instruments under the [Social Needs Screening and Intervention \(SNS-E\) measure](#))
- Document findings in the patients' medical record using Z-Codes
- Refer patients to community resources and/or care coordination

Reference the [Social Needs Screening and Intervention \(SNS-E\) measure](#) to ensure your practice is effectively capturing and sharing this information with the health plan.

If you are interested in learning more about SDOH and what you can do to help improve patient outcomes, please refer to the following additional resources:

- [World Health Organization: SDOH overview](#)
- [CDC: Social Determinants of Health](#)
- [CMS: Social Drivers of Health and Health-Related Social Needs](#)
- [Office of Disease Prevention and Health Promotion: SDOH Literature Summaries](#)
- [Illinois Department of Public Health: Understanding SDOH](#)
- [Agency for Healthcare Research and Quality: SDOH & Practice Improvement](#)
- [American Heart Association: Social Determinants of Health in Cardiology](#)

Quality Education Webinars (QEWebinars)



To support our providers in their quality improvement efforts, Meridian's Quality Improvement team hosts a series of webinars on topics related to improving patients' quality of care.

We hope these sessions will assist provider teams to improve HEDIS® scores and drive better incentive payments to your practice.

All office staff can attend, including providers, administrative staff, and quality teams. Participants can watch the webinars remotely using the Microsoft Teams or Zoom meeting invitation link provided. If you are interested in being added to the invite distribution list, please email ILHEDISOps@mhplan.com. [Sign up for health plan news](#) to find out about upcoming webinars and receive HEDIS tips.

Sample webinar topics include:

- Behavioral Healthcare
- HEDIS® Pay for Quality (P4Q) Program
- HEDIS® Measure & Exclusion Criteria
- Best Practices for Closing Care Gaps
- CAHPS® Survey and Patient Satisfaction
- Z-codes and Social Determinants of Health
- Health Outcomes Survey (HOS)
- Start Smart for Your Baby® program
- Overview of Illinois family of plans: Meridian Medicaid Plan, Wellcare, YouthCare, Wellcare Meridian Dual Align (HMO D-SNP), and Ambetter
- Risk Adjustment
- Member Outreach with Care Coordination
- Vendor Updates and Outreach

Access & Availability ● ● ●

To ensure medical services are available to our members on a timely basis, Primary Care Provider (PCP), behavioral health, and specialist appointment availability standards for Meridian and its family of plans are established and noted in the following section. Information is included for all plans, including the Meridian Medicaid Plan, YouthCare, Wellcare By Meridian (D-SNP), Ambetter, and Wellcare.

Each year, our Quality Improvement team conducts a telephone audit using the standards below set forth by NCQA, CMS, and/or state regulations. These audits are conducted to monitor provider compliance with Illinois Medicaid contract requirements. The process of conducting the annual audits is outlined below.

IDENTIFY: Meridian audits a sample of its contracted PCPs, behavioral health practitioners, and specialists.

OUTREACH: Conducted by a third-party vendor via phone, up to three attempts are made to reach a live person.

ANALYZE: Analysis is performed based on all data collected.

REPORT: Letters are mailed to offices indicating the results of the audit. Any offices that did not meet the standards are placed on a corrective action plan and are asked to identify ways to improve their appointment availability.

1. **Medicaid** Appointment and Timely Access to Care Standards

Primary Care Appointments	
Type of Care/Appointment	Length of Wait Time
Preventative/Routine Care (Child < 6 months)	Within two (2) weeks of request
Preventative/Routine Care (Child \geq 6 months)	Within five (5) weeks of request
Preventative/Routine Care (Adult)	Within five (5) weeks of request
Urgent/Non-Emergent (Medically Necessary) Care	Within twenty-four (24) hours of request
Non-Urgent/Non-Emergent Conditions	Within three (3) weeks of request
Initial Prenatal w/o Problems (First Trimester)	Within two (2) weeks of request
Prenatal (Second Trimester)	Within one (1) week of request
Prenatal (Third Trimester)	Within three (3) calendar days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Medicaid Appointment and Timely Access to Care Standards (continued)

Behavioral Health Appointments	
Type of Care/Appointment	Length of Wait Time
Life-Threatening Emergency	Immediate admittance or referred to the Emergency Room
Non-Life-Threatening Emergency	Within six (6) hours of request
Urgent Care Visit	Within forty-eight (48) hours of request
Initial Visit for Routine Care	Within ten (10) business days of request
Follow-Up Visit for Routine Care	Within twenty (20) business days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Specialty Care Appointments	
Type of Care/Appointment	Length of Wait Time
Routine Care (Child < 6 months)	Within two (2) weeks of request
Routine Care (Child \geq 6 months)	Within five (5) weeks of request
Routine Care (Adult)	Within five (5) weeks of request
Urgent/Non-Emergent (Medically Necessary) Care	Within one (1) business day of request
Non-Urgent/Non-Emergent Conditions	Within three (3) weeks of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Medical coverage 24 hours a day, 7 days a week

Medicaid Primary Care After-Hours Requirements

Acceptable after-hours access mechanisms include:

- Answering service
- On-call pager/cellular connection
- Call forwarding to practitioner's home/other location
- Published after-hours telephone number and recorded voice message directing patients to a practitioner for urgent and non-life-threatening conditions.

The message should not instruct patients to obtain treatment at the Emergency Room for non-life-threatening emergencies but MUST direct patients in a medical emergency to call 911 or go to the nearest Emergency Room/Urgent Care.

Voice Message MUST contain ONE of the following:

- Message forwards to on-call practitioner
- Message forwards to an answering service
- Message gives the on-call practitioner's number
- Message gives the on-call practitioner's pager number
- Message refers patient to another office, practitioner, or on-call service

Message may not only direct patients to the Emergency Room. The patient must be able to leave a message for an on-call doctor, speak with an on-call doctor, or be forwarded to an on-call doctor.

2. **YouthCare** Appointment and Timely Access to Care Standards

Primary Care Appointments	
Type of Care/Appointment	Length of Wait Time
Preventative/Routine Care (Child < 6 months)	Within two (2) weeks of request
Preventative/Routine Care (Child \geq 6 months)	Within five (5) weeks of request
Preventative/Routine Care (Adult)	Within five (5) weeks of request
Urgent/Non-Emergent (Medically Necessary) Care	Within twenty-four (24) hours of request
Non-Urgent/Non-Emergent Conditions	Within three (3) weeks of request
Initial Prenatal w/o Problems (First Trimester)	Within two (2) weeks of request
Prenatal (Second Trimester)	Within one (1) week of request
Prenatal (Third Trimester)	Within three (3) calendar days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Behavioral Health Appointments	
Type of Care/Appointment	Length of Wait Time
Life-Threatening Emergency	Immediate admittance or referred to the Emergency Room
Non-Life-Threatening Emergency	Within six (6) hours of request
Urgent Care Visit	Within forty-eight (48) hours of request
Initial Visit for Routine Care	Within ten (10) business days of request
Follow-Up Visit for Routine Care	Within twenty (20) business days of request

YouthCare Appointment and Timely Access to Care Standards (continued)

Behavioral Health Appointments	
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients
Specialty Care Appointments	
Type of Care/Appointment	Length of Wait Time
Preventative/Routine Care (Child < 6 months)	Within two (2) weeks of request
Preventative/Routine Care (Child ≥ 6 months)	Within five (5) weeks of request
Urgent/Non-Emergent (Medically Necessary) Care	Within one (1) business day of request
Non-Urgent/Non-Emergent Conditions	Within three (3) weeks of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Medical coverage 24 hours a day, 7 days a week

YouthCare Primary Care After-Hours Requirements

Acceptable after-hours access mechanisms include:

- Answering service
- On-call pager/cellular connection
- Call forwarding to practitioner's home/other location
- Published after-hours telephone number and recorded voice message directing patients to a practitioner for urgent and non-life-threatening conditions.

The message should not instruct patients to obtain treatment at the Emergency Room for non-life-threatening emergencies but MUST direct patients in a medical emergency to call 911 or go to the nearest Emergency Room/Urgent Care.

Voice Message MUST contain ONE of the following:

- Message forwards to on-call practitioner
- Message forwards to an answering service
- Message gives the on-call practitioner's number
- Message gives the on-call practitioner's pager number
- Message refers patient to another office, practitioner, or on-call service

Message may not only direct patients to the Emergency Room. The patient must be able to leave a message for an on-call doctor, speak with an on-call doctor, or be forwarded to an on-call doctor.

3. Wellcare By Meridian D-SNP Appointment and Timely Access to Care Standards

Primary Care Appointments	
Type of Care/Appointment	Length of Wait Time
Preventative/Routine Care	Within twenty-five (25) business days of request
Urgent/Non-Emergent (Medically Necessary) Care	Within one (1) business days of request
Non-Urgent/Non-Emergent Conditions (Medically Necessary)	Within seven (7) business days of request
Non-Urgent/Non-Emergent Conditions	Within three (3) weeks of request
Initial Prenatal w/o Problems (First Trimester)	Within two (2) weeks of request
Prenatal (Second Trimester)	Within one (1) week of request
Prenatal (Third Trimester)	Within three (3) days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Behavioral Health Appointments	
Type of Care/Appointment	Length of Wait Time
Life-Threatening Emergency	Immediate admittance or referred to the Emergency Room
Non-Life-Threatening Emergency	Within six (6) hours of request
Urgent Care Visit	Within forty-eight (48) hours of request
Initial Visit for Routine Care	Within ten (10) business days of request
Follow-Up Visit for Routine Care	Within twenty (20) business days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Specialty Care Appointments	
Type of Care/Appointment	Length of Wait Time
Routine Care (Adult)	Within twenty-five (25) business days of request
Urgent/Non-Emergent (Medically Necessary) Care	Within one (1) business days of request

Wellcare By Meridian D-SNP Appointment and Timely Access to Care Standards (continued)

Specialty Care Appointments	
Non-Urgent/Non-Emergent Conditions (Medically Necessary)	Within seven (7) business days of request
Non-Urgent/Non-Emergent Conditions	Within three (3) weeks of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Wellcare By Meridian D-SNP Primary Care After-Hours Requirements

Acceptable After-Hours access mechanisms include:

- Answering service
- On-call pager/cellular connection
- Call forwarding to practitioner's home/other location
- Published after-hours telephone number and recorded voice message directing patients to a practitioner for urgent and non-life-threatening conditions.

The message should not instruct patients to obtain treatment at the Emergency Room for non-life-threatening emergencies but MUST direct patients in a medical emergency to call 911 or go to the nearest Emergency Room/Urgent Care.

Voice Message MUST contain ONE of the following:

- Message forwards to on-call practitioner
- Message forwards to an answering service
- Message gives the on-call practitioner's number
- Message gives the on-call practitioner's pager number
- Message refers patient to another office, practitioner, or on-call service

Message may not only direct patients to the Emergency Room. The patient must be able to leave a message for an on-call doctor, speak with an on-call doctor, or be forwarded to an on-call doctor.

4. **Marketplace (Ambetter) Appointment and Timely Access to Care Standards**

Primary Care Appointments	
Type of Care/Appointment	Length of Wait Time
Preventative/Routine Care	Within fifteen (15) calendar days of request
Urgent/Non-Emergent (Medically Necessary) Care	Within twenty-four (24) hours of request
Sick Care	Within twenty-four (24) hours of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Behavioral Health Appointments	
Type of Care/Appointment	Length of Wait Time
Life-Threatening Emergency	Immediate admittance or referred to the Emergency Room
Non-Life-Threatening Emergency	Within six (6) hours of request
Urgent Care Visit	Within forty-eight (48) hours of request
Initial Visit for Routine Care	Within ten (10) business days of request
Follow-Up Visit for Routine Care	Within ten (10) business days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Specialty Care Appointments	
Type of Care/Appointment	Length of Wait Time
Routine Care	Within thirty (30) calendar days of request
Urgent Care Visit	Within forty-eight (48) hours of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Medical coverage 24 hours a day, 7 days a week

Marketplace (Ambetter) Primary Care After-Hours Requirements

Acceptable After-Hours access mechanisms include:

- Answering service
- On-call pager/cellular connection
- Call forwarding to practitioner's home/other location
- Published after-hours telephone number and recorded voice message directing patients to a practitioner for urgent and non-life-threatening conditions.

The message **should not instruct patients to obtain treatment at the Emergency Room for non-life-threatening emergencies** but MUST direct patients in a medical emergency to call 911 or go to the nearest Emergency Room/Urgent Care.

Voice Message MUST contain ONE of the following:

- Message forwards to on-call practitioner
- Message forwards to an answering service
- Message gives the on-call practitioner's number
- Message gives the on-call practitioner's pager number
- Message refers patient to another office, practitioner, or on-call service

Message may not only direct patients to the Emergency Room. The patient must be able to leave a message for an on-call doctor, speak with an on-call doctor, or be forwarded to an on-call doctor.

5. **Wellcare (Medicare)** Appointment and Timely Access to Care Standards

Primary Care Appointments	
Type of Care/Appointment	Length of Wait Time
Preventative/Routine Care	Within thirty (30) business days of request
Urgent/Non-Emergent (Medically Necessary) Care	Within twenty-four (24) hours of request
Sick Care	Within seven (7) calendar days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Behavioral Health Appointments	
Type of Care/Appointment	Length of Wait Time
Life-Threatening Emergency	Immediate admittance or referred to the Emergency Room
Non-Life-Threatening Emergency	Within six (6) hours of request
Urgent Care Visit	Within forty-eight (48) hours of request

Wellcare (Medicare) Appointment and Timely Access to Care Standards (continued)

Behavioral Health Appointments	
Initial Visit for Routine Care	Within ten (10) business days of request
Follow-Up Visit for Routine Care	Within thirty (30) business days of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Specialty Care Appointments	
Type of Care/Appointment	Length of Wait Time
Routine Care	Within thirty (30) calendar days of request
Urgent Care Visit	Within twenty-four (24) hours of request
Office Wait Time	Within thirty (30) minutes
Different Hours for Member Plans?	No, hours must be the same for all members and patients

Medical coverage 24 hours a day, 7 days a week

Wellcare (Medicare) Primary Care After-Hours Requirements

Acceptable After-Hours access mechanisms include:

- Answering service
- On-call pager/cellular connection
- Call forwarding to practitioner's home/other location
- Published after-hours telephone number and recorded voice message directing patients to a practitioner for urgent and non-life-threatening conditions.

The message should not instruct patients to obtain treatment at the Emergency Room for non-life-threatening emergencies but MUST direct patients in a medical emergency to call 911 or go to the nearest Emergency Room/Urgent Care.

Voice Message MUST contain ONE of the following:

- Message forwards to on-call practitioner
- Message forwards to an answering service
- Message gives the on-call practitioner's number
- Message gives the on-call practitioner's pager number
- Message refers patient to another office, practitioner, or on-call service

Message may not only direct patients to the Emergency Room. The patient must be able to leave a message for an on-call doctor, speak with an on-call doctor, or be forwarded to an on-call doctor.

Caring for and Communicating with Individuals with Intellectual and Developmental Disabilities (IDD)

As a provider, it is important to be aware of the following health disparities individuals with intellectual and developmental disabilities may experience:

- Fewer preventive screenings than the general population
- Financial and transportation limitations
- Lack of access to providers with specialized training or experience caring for individuals in these populations

Complete an Annual Functional Status Assessment

This assessment measures the patient's ability to perform daily tasks and helps to identify any functional decline. For Meridian patients, please indicate one of the following in the medical record:

- Notation and date that Activities of Daily Living (ADL) were assessed: Bathing, dressing, eating, transferring, using toilet, walking
- Notation and date that Instrumental Activities of Daily Living (IADL) were assessed: Shopping, driving or using public transportation, meal preparation, housework, taking medications, using the telephone
- Result of a standardized functional status assessment and the date it was performed: Assessment of Living Skills and Resources (ALSAR), Barthel ADL Index Physical Self-Maintenance (ADLS) Scale, Bayer Activities of Daily Living (B-ADL) Scale, Extended Activities of Daily Living (EADL) Scale
- Chronic conditions at a younger age
- A higher risk of obesity
- A greater incidence of mental illness
- An accelerated aging process

It is important that providers and health plans adapt and coordinate care for this population by communicating appropriately and respectfully about individuals with disabilities.

Individuals with intellectual and developmental disabilities often go through cognitive and behavioral changes. These include anxiety, depression, dementia, self-injurious behavior, and other factors. These changes may result from pain and discomfort related to other medical issues.

As a provider, it is important to be aware of the following ways to communicate and to emphasize the person first, not the disability:

- Emphasize abilities, not limitations
- Do not use language that suggests the lack of something
- Emphasize the need for accessibility, not the disability
- Do not use offensive language
- Avoid language that implies negative stereotypes
- Do not portray people with disabilities as inspirational only because of their disability

Tips on Using People-First Language

This chart provides tips on what language to use when communicating with an individual with a disability.

Tips	Use	Do Not Use
Emphasize abilities, not limitations	Person who uses a wheelchair	Confined or restricted to a wheelchair, wheelchair bound
	Person who uses a device to speak	Can't talk, mute
Do not use language that suggests the lack of something	Person with a disability	Disabled, handicapped
	Person of short stature	Midget
	Person with cerebral palsy	Cerebral palsy victim
	Person with epilepsy or seizure disorder	Epileptic
	Person with multiple sclerosis	Afflicted by multiple sclerosis
Emphasize the need for accessibility, not the disability	Accessible parking or bathroom	Handicapped parking or bathroom
Do not use offensive language	Person with a physical disability	Crippled, lame, deformed, invalid, spastic
	Person with an intellectual, cognitive, developmental disability	Slow, simple, moronic, defective, afflicted, special person
	Person with an emotional or behavioral disability, mental health impairment, or a psychiatric disability	Insane, crazy, psycho, maniac, nuts
Avoid language that implies negative stereotypes	Person without a disability	Normal person, healthy person
Do not portray people with disabilities as inspirational only because of their disability	Person who is successful, productive	Has overcome his/her disability, is courageous

Source: [cdc.gov/disability-and-health/articles-documents/communicating-with-and-about-people-with-disabilities.html](https://www.cdc.gov/disability-and-health/articles-documents/communicating-with-and-about-people-with-disabilities.html)

Medicare-Medicaid Plan Codes

Description	CPT Category*	CPT II Category*	HCPCS*
Functional Status Assessment	99483	1170F	G0438, G0439

*Codes subject to change

Resources for Patients and Providers

- Illinois Department of Healthcare and Family Services (HFS): hfs.illinois.gov/medicalclients/disabled.html
- Illinois Department of Human Services (IDHS): **217-782-3075**
- For a copy of current clinical practice guidelines, visit our website at ILmeridian.com.
- Meridian Provider Resources: ilmeridian.com/providers.html

Early Periodic Screening, Diagnosis, and Treatment (EPSDT) ●●

The Early Periodic Screening, Diagnosis, and Treatment (EPSDT) program promotes the physical, mental, social, emotional, and behavioral health of children under the age of 21 through recommended well-child screenings.

These services give children early access to preventative and comprehensive healthcare to help prevent disease and identify medical, developmental, and social-emotional concerns in their early stages – when they are more effectively treated.

Mandatory Screening Components

An EPSDT screening for Medicaid members must include:

- Initial and interval history**, including a comprehensive health and developmental history of physical and mental development
- An unclothed physical exam**, documented in the member's medical chart
- Measurements**: Weight, length, head circumference, body mass index (BMI) percentile*, and blood pressure
- Nutrition/obesity prevention**: Assess and educate, with specific documentation on nutrition and physical activity*
- Oral health assessment*** and age-appropriate fluoride varnish application

Specific Screenings

- Caregiver and/or adolescent depression screening
- Developmental surveillance/screening
- Autism screening
- Psychosocial/behavioral assessment
- Alcohol and drug use assessment
- Sensory screenings for vision and hearing*

Immunizations and Age-Appropriate Screenings

- Review immunization history and administer immunizations at recommended ages or as needed
- Offer anticipatory guidance and health education at every visit
- Conduct age-appropriate laboratory tests, including:
 - Lipid screening
 - Hemoglobin/hematocrit
 - Lead blood testing (2)/risk screening
 - Newborn screening: blood, hearing, critical congenital heart disease
 - Sexually transmitted infections and HIV screening
 - Cervical dysplasia screening
 - TB testing
- Provide other medically necessary health care, diagnostic services, and treatment measures

* *These areas are frequently non-compliant or not properly documented to show compliance during medical record reviews.*

Billing Codes for New or Established Patients

Ages	New	Established
<1	99381	99391
1–4	99382	99392
5–11	99383	99393
12–17	99384	99394
18–21	99385	99395
Fluoride Varnish: 99188		

EPSTD/Bright Futures Referral Codes

When making an EPSDT referral, add the appropriate Y modifier to line 10d of the CMS 1500.

Referral Code	Referral Type
YD	Dental referral
YM	Medical referral
YB	Behavioral health referral
YV	Vision referral
YH	Hearing referral
YO	Other referral

Online Resources

- Meridian Early Periodic Screening, Diagnosis, and Treatment (EPSDT) toolkit: ILmeridian.com/content/dam/centene/meridian/il/pdf/Meridian_EPSTD_Provider%20Toolkit_0811.pdf
- American Academy of Pediatrics Recommendations for Preventive Pediatric Care (Periodicity Schedule): aap.org/en/practice-management/care-delivery-approaches/periodicity-schedule/
- Bright Futures, a national health promotion and preventive initiative led by the American Academy of Pediatrics: aap.org/en/practice-management/bright-futures
- Centers for Disease Control and Prevention (CDC) vaccination schedules: cdc.gov/vaccines/index.html

Care for Kids Periodicity Schedule

KEY: ● To be performed ○ Assess risk ↔ Screen at least once during time period indicated

Appendix

Antidepressant Medication List

Description	Prescription
Miscellaneous antidepressants	<ul style="list-style-type: none">BupropionVortioxetine
Monoamine oxidase inhibitors	<ul style="list-style-type: none">IsocarboxazidPhenelzine
Phenylpiperazine antidepressants	<ul style="list-style-type: none">Nefazodone
Psychotherapeutic combinations	<ul style="list-style-type: none">Amitriptyline-chlordiazepoxideAmitriptyline-perphenazineFluoxetine-olanzapine
SNRI antidepressants	<ul style="list-style-type: none">DesvenlafaxineDuloxetine
SSRI antidepressants	<ul style="list-style-type: none">CitalopramEscitalopramFluoxetine
Tetracyclic antidepressants	<ul style="list-style-type: none">Maprotiline
Tricyclic antidepressants	<ul style="list-style-type: none">AmitriptylineAmoxapineClomipramineDesipramineDoxepin (>6 mg)
	<ul style="list-style-type: none">VilazodoneSelegilineTranylcypromineTrazodoneLevomilnacipranVenlafaxineFluvoxamineParoxetineSertralineMirtazapineImipramineNortriptylineProtriptylineTrimipramine

Approved Depression Screening Instruments, Codes, and Positive Findings ●●●

A standard assessment instrument that has been normalized and validated for the appropriate clinical population. Eligible screening instruments with thresholds for positive findings include:

Instruments for Adolescents (≤17 years)	Total Score LOINC Codes*	Positive Finding
Patient Health Questionnaire (PHQ-9) [®]	44261-6	Total score ≥ 10
Patient Health Questionnaire Modified for Teens (PHQ- 9M) [®]	89204-2	Total score ≥ 10
Patient Health Questionnaire-2 (PHQ-2) ^{®1}	55758-7	Total score ≥ 3
Beck Depression Inventory— Fast Screen (BDI-FS) ^{®1,2}	89208-3	Total score ≥ 8
Center for Epidemiologic Studies Depression Scale—Revised (CESD-R)	89205-9	Total score ≥ 17
Edinburgh Postnatal Depression Scale (EPDS)	71354-5	Total score ≥ 10
PROMIS Depression	71965-8	Total score (T Score) ≥ 60

¹ Brief screening instrument; all other instruments are full-length

² Proprietary; cost or licensing requirement may be associated with use

*Codes subject to change

Instruments for Adults (18+ years)	Total Score LOINC Codes*	Positive Finding
Patient Health Questionnaire (PHQ-9) [®]	44261-6	Total score ≥ 10
Patient Health Questionnaire-2 (PHQ-2) ^{®1}	55758-7	Total score ≥ 3
Beck Depression Inventory—Fast Screen (BDI-FS) ^{®1,2}	89208-3	Total score ≥ 8
Beck Depression Inventory (BDI-II)	89209-1	Total score ≥ 20
Center for Epidemiologic Studies Depression Scale—Revised (CESD-R)	89205-9	Total score ≥ 17
Duke Anxiety-Depression Scale (DUKE-AD) ^{®2}	90853-3	Total score ≥ 30
Edinburgh Postnatal Depression Scale (EPDS)	71354-5	Total score ≥ 10
My Mood Monitor (M-3) [®]	71777-7	Total score ≥ 5
PROMIS Depression	71965-8	Total score (T Score) ≥ 60

Instruments for Adults (18+ years)	Total Score LOINC Codes*	Positive Finding
Clinically Useful Depression Outcome Scale (CUDOS)	90221-3	Total score ≥ 31
Geriatric Depression Scale Short Form (GSD)	48545-8	Total score >5
Geriatric Depression Scale Long Form (GSD)	48544-1	Total score >10

¹ Brief screening instrument. All other instruments are full-length.

² Proprietary; cost or licensing requirement may be associated with use.

*Codes subject to change

Follow-Up Visit ●●●●

Description	Codes*
An outpatient, telephone, e-visit, or virtual check-in follow-up visit with a diagnosis of depression or other behavioral health condition	CPT: 98960-98962, 98966-98968, 98970-98972, 98980, 98981, 99078, 99202-99205, 99211-99215, 99242-99245, 99341, 99342, 99344, 99345 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99441-99443, 99457, 99458, 99483 HCPCS: G0071, G0463, G2010, G2012, G2250-G2252 T1015 SNOMED CT: 42137004, 50357006, 86013001, 90526000, 108220007, 108221006, 185317003, 185389009, 281036007, 314849005, 386472008, 386473003, 390906007, 401267002, 406547006, 870191006
Depression Case Management Encounter	CPT: 99366, 99492-99494 HCPCS: T1016, T1017, T2022, T2023 SNOMED CT: 1344983001, 1344984007, 1344994002, 1345003009, 1345013001, 1345014007, 182832007, 225333008, 385828006, 386230005, 409022004, 410216003, 410219005, 410328009, 410335001, 410346003, 410347007, 410351009, 410352002, 410353007, 410354001, 410356004, 410360001, 410363004, 410364005, 410366007, 416341003, 416584001, 424490002, 425604002, 737850002, 621561000124106, 661051000124109, 662081000124106, 662541000124107

Description	Codes*
Behavioral Health Encounter	<p>CPT: 90791, 90792, 90832-90834, 90836-90839, 90845-90847, 90849, 90853, 90865, 90867-90870, 90875, 90876, 90880, 90887, 99484, 99492, 99493</p> <p>HCPGS: G0155, G0176, G0177, G0409-G0411, G0511, G0512, H0002, H0004, H0031, H0034-H0037, H0039, H0040, H2000, H2001, H2010-H2020, S0201, S9480, S9484, S9485</p> <p>SNOMED CT: 5694008, 10197000, 10997001, 38756009, 45392008, 79094001, 88848003, 90407005, 91310009, 165171009, 165190001, 225337009, 370803007, 372067001, 385721005, 385724002, 385725001, 385726000, 385727009, 385887004, 385889001, 385890005, 401277000, 410223002, 410224008, 410225009, 410226005, 410227001, 410228006, 410229003, 410230008, 410231007, 410232000, 410233005, 410234004, 439141002</p> <p>ICD-10-CM: Z71.82</p>
A dispensed antidepressant medication Antidepressant Medications List	See Antidepressant Medications List .
Documentation of additional depression screening on a full-length instrument indicating either no depression or no symptoms that require follow-up (i.e., a negative screen) on the same day as a positive screen on a brief screening instrument. ***	

*Codes subject to change

Depression or Other Behavioral Health Condition

Depression or Other Behavioral Health Condition Codes*

ICD-10-CM: F01.51, F01.511, F01.518, F06.4, F10.180, F10.280, F10.980, F11.188, F11.288, F11.988, F12.180, F12.280, F12.980, F13.180, F13.280, F13.980, F14.180, F14.280, F14.980, F15.180, F15.280, F15.980, F16.180, F16.280, F16.980, F18.180, F18.280, F18.980, F19.180, F19.280, F19.980, F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.8, F32.81, F32.89, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.8, F34.81, F34.89, F34.9, F39, F40.00, F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.248, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.8, F43.81, F43.89, F43.9, F44.89, F45.21, F51.5, F53, F53.0, F53.1, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3, F63.81, F63.89, F63.9, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9, O90.6, O99.340, O99.341, O99.342, O99.343, O99.344, O99.345

Depression or Other Behavioral Health Condition Codes*

SNOMED CT: 109006, 162004, 281004, 600009, 832007, 899001, 1145003, 1196001, 1376001, 1380006, 1383008, 1499003, 1686006, 1816003, 1855002, 1973000, 2312009, 2403008, 2506003, 2618002, 2815001, 3109008, 3158007, 3530005, 3914008, 4306003, 4441000, 4926007, 4932002, 4997005, 5095008, 5158005, 5444000, 5464005, 5507002, 5509004, 5510009, 5703000, 6348008, 7025000, 7052005, 7200002, 7291006, 7397008, 7461003, 7794004, 8185002, 8635005, 8837000, 9167000, 9340000, 9674006, 9760005, 10278007, 10327003, 10586006, 10875004, 10981006, 11806006, 11941006, 12939007, 12969000, 13127006, 13313007, 13438001, 13581000, 13601005, 13670005, 13746004, 14070001, 14077003, 14144000, 14183003, 14291003, 14495005, 14784000, 15193003, 15277004, 15639000, 15945005, 15977008, 16295005, 16506000, 16805009, 16966009, 16990005, 17155009, 17226007, 17262008, 17496003, 17782008, 17961008, 18003009, 18085000, 18260003, 18478005, 18573003, 18653004, 18689007, 18818009, 18941000, 19300006, 19445006, 19527009, 19694002, 19766004, 20010003, 20250007, 20385005, 20876004, 20960007, 21586000, 21634003, 21897009, 21900002, 22121000, 22230001, 22407005, 22419002, 23148009, 23560001, 23645006, 24121004, 24125008, 24315006, 24781009, 25501002, 25766007, 25922000, 26025008, 26203008, 26453000, 26472000, 26516009, 26530004, 26665006, 26714005, 27387000, 27544004, 27956007, 28357009, 28368009, 28475009, 28663008, 28676002, 28864000, 28884001, 29212009, 29599000, 29733004, 29929003, 30059008, 30310000, 30336007, 30491001, 30509009, 30520009, 30605009, 30687003, 30935000, 31027006, 31177006, 31358003, 31373002, 31446002, 31611000, 31648009, 31658008, 31715000, 31781004, 32009006, 32174002, 32358001, 32388005, 32552001, 32721004, 32875003, 32880007, 33078009, 33135002, 33323008, 33380008, 33449004, 33736005, 33871004, 34116005, 34315001, 34938008, 35218008, 35252006, 35253001, 35481005, 35489007, 35607004, 35722002, 35827000, 35846004, 35919005, 36170009, 36217008, 36474008, 36583000, 36622002, 36923009, 37331004, 37739004, 37746008, 37868008, 37872007, 37941009, 38295006, 38328002, 38368003, 38451003, 38547003, 38694004, 39003006, 39465007, 39610001, 39807006, 39809009, 39951001, 40379007, 40568001, 40571009, 40673001, 40926005, 40987004, 41021005, 41083005, 41526007, 41552001, 41832009, 41836007, 42344001, 42594001, 42810003, 42868002, 42925002, 43150009, 43497001, 43568002, 43614003, 43769008, 44031002, 44124003, 44376007, 44966003, 45479006, 45677003, 45912004, 46206005, 46229002, 46244001, 46721000, 46975003, 47372000, 47447001, 47505003, 47664006, 47916000, 48500005, 48589009, 48826008, 48937005, 49271002, 49468007, 49481000, 49512000, 49564006, 50026000, 50320000, 50705009, 50722006, 50933003, 50983008, 51133006, 51443000, 51493001, 51637008, 51771007, 52702003, 52824009,

Depression or Other Behavioral Health Condition Codes*

52866005, 52910006, 52954000, 53049002, 53050002, 53467004, 53607008, 53936005, 53956006, 54319003, 54417002, 54502004, 54587008, 54761006, 55009008, 55341008, 55516002, 55668003, 55967005, 56576003, 56627002, 56641006, 56882008, 57194009, 57588009, 57715001, 58214004, 58329000, 58535001, 58647003, 58703003, 59216005, 59617007, 59645001, 59651006, 59923000, 60099002, 60123008, 61104008, 61144001, 61157009, 61180001, 61212007, 61403008, 61569007, 61831009, 61901004, 62351001, 63181006, 63204009, 63249007, 63393005, 63395003, 63412003, 63649001, 63701002, 63778009, 63909006, 63983005, 64060000, 64165008, 64731001, 64905009, 65042007, 65064003, 65179007, 66307007, 66344007, 66347000, 66381006, 66590003, 66631006, 66936004, 67002003, 67123006, 67195008, 67711008, 68019004, 68569003, 68618008, 68890003, 68963006, 68995007, 69322001, 69361009, 69392006, 69479009, 69482004, 70328006, 70546001, 70691001, 70747007, 70814008, 70932007, 71103003, 71294008, 71328000, 71336009, 71787009, 71959007, 71961003, 71984005, 72366004, 72861004, 73097000, 73462009, 73471000, 73867007, 74010007, 74266001, 74686005, 74732009, 74850006, 74934004, 75084000, 75122001, 75360000, 75752004, 75837004, 76105009, 76129002, 76236006, 76441001, 76566000, 76812003, 76868007, 77355000, 77475008, 77486005, 77675002, 77815007, 77911002, 78004001, 78269000, 78358001, 78640000, 78667006, 79204003, 79298009, 79524000, 79584002, 79842004, 79866005, 80711002, 80868005, 81319007, 81463002, 82218004, 82339009, 82415003, 82494000, 82738004, 82998009, 83176005, 83225003, 83253003, 83367009, 83458005, 83501007, 83631006, 83746006, 83890006, 84466009, 84760002, 84788008, 84984002, 85061001, 85080004, 85248005, 85821003, 85861002, 86058007, 87132004, 87203005, 87414006, 87512008, 87798009, 87842000, 87950005, 87991007, 88740003, 88845000, 88939009, 88975006, 89415002, 89451009, 89618007, 89948007, 90755006, 90790003, 93461009, 95637005, 106015009, 109805003, 111475002, 111477005, 111479008, 111480006, 111482003, 111483008, 111484002, 111485001, 111487009, 111490003, 111491004, 126943008, 128293007, 129104009, 129604005, 129606007, 162218007, 162313000, 187921002, 191447007, 191455000, 191457008, 191458003, 191459006, 191466007, 191471000, 191475009, 191476005, 191478006, 191480000, 191483003, 191485005, 191486006, 191492000, 191493005, 191494004, 191495003, 191496002, 191499009, 191501001, 191508007, 191526005, 191527001, 191531007, 191542003, 191547009, 191548004, 191554003, 191555002, 191559008, 191561004, 191562006, 191563001, 191564007, 191565008, 191567000, 191569002, 191570001, 191572009, 191574005, 191577003, 191583000, 191584006, 191586008, 191588009, 191590005, 191592002, 191593007, 191595000, 191597008, 191604000, 191610000, 191611001, 191613003, 191616006, 191618007, 191620005, 191621009, 191623007,

Depression or Other Behavioral Health Condition Codes*

191621009, 191623007, 191625000, 191627008, 191629006, 191630001, 191634005, 191636007, 191638008, 191639000, 191641004, 191643001, 191659001, 191667009, 191668004, 191670008, 191672000, 191676002, 191678001, 191680007, 191687005, 191689008, 191690004, 191692007, 191693002, 191696005, 191697001, 191714002, 191722009, 191724005, 191725006, 191726007, 191736004, 191737008, 191738003, 191753006, 191765005, 191766006, 191772006, 191773001, 191774007, 191787001, 191811004, 191812006, 191813001, 191966002, 192014006, 192037000, 192038005, 192039002, 192041001, 192042008, 192044009, 192046006, 192049004, 192051000, 192052007, 192054008, 192056005, 192057001, 192058006, 192061007, 192063005, 192064004, 192065003, 192069009, 192072002, 192073007, 192079006, 192080009, 192082001, 192092009, 192097003, 192099000, 192100008, 192108001, 192110004, 192119003, 192127007, 192131001, 192132008, 192147004, 192362008, 192527004, 192562009, 192611004, 192616009, 192631000, 197480006, 199257008, 199259006, 199260001, 199261002, 199262009, 229623002, 229699006, 229700007, 229701006, 229715008, 229740001, 229744005, 229745006, 230334008, 230455006, 231437006, 231438001, 231442003, 231443008, 231444002, 231445001, 231446000, 231449007, 231452004, 231453009, 231454003, 231455002, 231456001, 231467000, 231485007, 231487004, 231489001, 231494001, 231496004, 231500002, 231504006, 231516000, 231520001, 231521002, 231522009, 231523004, 231524005, 231525006, 231526007, 231527003, 231528008, 231530005, 231536004, 231537008, 231538003, 231539006, 231540008, 231541007, 231542000, 231864002, 233690008, 235004005, 237351003, 237352005, 238961003, 238966008, 238967004, 238972008, 238973003, 238974009, 238976006, 238977002, 238978007, 238979004, 247803002, 247804008, 267320004, 268612007, 268617001, 268619003, 268621008, 268622001, 268624000, 268633003, 268634009, 268658008, 268661009, 268662002, 268664001, 268666004, 268667008, 268668003, 268669006, 268673009, 268734000, 270901009, 270902002, 270905000, 271428004, 271952001, 274948002, 274952002, 274953007, 275446004, 275473003, 275474009, 276296007, 276297003, 278506006, 278508007, 278852008, 278853003, 279611005, 280032002, 280943007, 280949006, 280994000, 286933003, 300706003, 300979000, 307417003, 310495003, 310496002, 310497006, 31173003, 312098001, 312936002, 313182004, 313915006, 317816007, 318784009, 319768000, 320751009, 357705009, 370143000, 371024007, 371026009, 371596008, 371599001, 371600003, 371604007, 371631005, 373618009, 386701004, 386810004, 386820009, 386821008, 386822001, 386823006, 386824000, 386825004, 402732001, 402733006, 402735004, 402736003, 403578003, 403579006, 403590001, 403593004, 403594005, 403595006, 404906000, 406506008, 408856003, 408857007, 416340002, , 417294004, 420774007, 425914008, 425919003, 426174008, 426881004, 427469003, 427975003, 428687006, 428703001, 429672007, 430744005, 430751001, 430852001, 430909002, 432091002, 439960005, 441704009, 441711008, 441716003,

Depression or Other Behavioral Health Condition Codes*

441833000, 442057004, 442245003, 442314000, 442351006, 443919007, 446175003, 446180007, 450714000, 473452003, 473456000, 473457009, 609225004, 698689005, 698690001, 698692009, 698693004, 698696007, 698697003, 698699000, 698700004, 698701000, 698946008, 698947004, 698951002, 699241002, 702356009, 702450004, 702732007, 702815001, 708037001, 712823008, 712824002, 712850003, 712884004, 713491003, 713583005, 713862009, 714829008, 715737004, 715924009, 716096005, 716199000, 716578009, 718393002, 719592004, 719600006, 720451004, 720452006, 720453001, 720454007, 720455008, 722287002, 723332005, 723899008, 723900003, 723901004, 723903001, 723905008, 723913009, 724673008, 724674002, 724675001, 724693000, 724706006, 724718002, 724719005, 724730008, 724733005, 724734004, 724735003, 724736002, 724737006, 724738001, 724739009, 724740006, 724741005, 724742003, 726772006, 733623005, 735549008, 735550008, 737225007, 762327001, 762332000, 762347009, 762509000, 762516004, 765176007, 766824003, 767631007, 767632000, 767633005, 767635003, 767636002, 770790004, 771448004, 771512003, 782501005, 783089006, 789061003, 789399002, 870260008, 870261007, 870262000, 870263005, 870264004, 870265003, 870266002, 870267006, 870268001, 870269009, 870270005, 870280009, 870282001, 870303005, 870304004, 870305003, 870306002, 870307006, 870308001, 1153575004, 1204417003, 1220650006, 1254652005, 251000119105, 261000119107, 271000119101, 281000119103, 7031000119100, 18491000119109, 21071000119101, 21831000119109, 22641000119109, 22981000119106, 23601000119102, 23741000119105, 31081000119101, 39951000119105, 60401000119104, 62211000119103, 62231000119108, 89381000119107, 89501000119108, 92501000119101, 94631000119100, 97571000119109, 104851000119103, 130121000119104, 133091000119105, 133121000119109, 153071000119108, 270291000119109, 288271000119103, 288421000119100, 288511000119108, 288751000119101, 288851000119106, 288861000119108, 368361000119108, 174000100004102, 10811121000119102, 10811161000119107, 16238741000119105, 16264621000119109, 16264821000119108, 16264901000119109, 16265061000119105, 16265301000119106, 16265951000119109, 16266831000119100, 16266991000119108, 19887002, 24109003, 54307006, 58963008, 102916005, 280947008, 403592009, 444613000, 1343347009, 1343597009, 1343598004, 1343608003, 1343614005, 1343615006, 1344950008

*Codes subject to change

Denominator Exclusions



Exclusions	Applicable Measures	Codes*
Hospice	All Measures	CPT: 99377, 99378 HCPCS: G9473, G9474, G9475, G9476, G9477, G9478, G9479, Q5003, Q5004, Q5005, Q5006, Q5007, Q5008, Q5010, S9126, T2042, T2043, T2044, T2045, T2046, G0182
Palliative Care	AAF-E, BCS-E, BPC-E, BPD-E, CBP, CCS-E, , COL-E, CRE, EED, GSD, KED, LBP, OMW, OSW, SPC-E, SPD-E, TSC-E	HCPCS: G9054, M1017 ICD-10: Z51.5
Colorectal Cancer	COL-E	ICD-10: C18.0-C18.9, C19, C20, C21.2, C21.8, C78.5, Z85.038, Z85.048
Total Colectomy	COL-E	CPT: 44150-44153, 44155-44158, 44210-44212
Bilateral Mastectomy	BCS-E	ICD-10: OHTV0ZZ
Unilateral Mastectomy with Bilateral Modifier	BCS-E	CPT: 19180, 19200, 19220, 19240, 19303-19307
Bilateral Procedure/ Bilateral Modifier	BCS-E	Modifier: 50 – Used for bilateral procedure
History of Bilateral Mastectomy	BCS-E	ICD-10: Z90.13
Complete Hysterectomy with No Residual Cervix and Absence of Cervix Diagnosis	CCS-E	CPT: 57530, 57531, 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275, 58280, 58285, 58290-58294, 58548, 58550, 58552-58554, 58570-58573, 58575, 58951, 58953, 58954, 58956, 59135 ICD-10: Q51.5, Z90.710, Z90.712
ESRD Diagnosis	BPC-E, CBP, KED, SPC-E, SPD-E	ICD-10: N18.5-N18.6

Exclusions	Applicable Measures	Codes*
Kidney Transplant	CBP, BPC-E	CPT: 50360, 50365, 50380 HCPCS: S2065 ICD-10: OTY00Z0, OTY00Z1, OTY00ZZ, OTY10Z0, OTY10Z1, OTY10ZZ
Partial Nephrectomy	BPC-E, CBP	CPT: 50240 ICD-10: OTB00ZZ, OTB04ZZ, OTB07ZZ, OTB08ZZ, OTB10ZZ, OTB13ZZ, OTB14ZZ, OTB17ZZ, OTB18ZZ
Cirrhosis	SPD, SPC	ICD-10: K70.30, K70.31, K71.7, K74.3, K74.4, K74.5, K74.60, K74.69, P78.81
Total Nephrectomy	BPC-E, CBP	CPT: 50220, 50225, 50230, 50234, 50236, 50340, 50370, 50543, 50545, 50546, 50548 ICD-10: OTT00ZZ, OTT04ZZ, OTT10ZZ, OTT14ZZ, OTT20ZZ, OTT24ZZ
Non-Live Births	PPC	ICD-10: Z37.1, Z37.4, Z37.7

Exclusions	Applicable Measures	Codes*
Pregnancy Diagnosis	BPC-E, CBP, SPC-E, SPD-E, WCC* <i>*With a prescription of isotretinoin or an X-Ray within 6 days after pregnancy test</i>	ICD-10: O09.00-O09.03, O09.10-O09.13, O09.211-O09.213, O09.219, O09.291-O09.293, O09.299, O09.30-O09.33, O09.40-O09.43, O09.511-O09.513, O09.519, O09.521-O09.523, O09.529, O09.611-O09.613, O09.621-O09.623, O09.629, O09.70-O09.73, O09.811-O09.813, O09.819, O09.21-O09.823, O09.829, O09.891-O09.893, O09.899, O09.90-O09.93, O09.A0-O09.A3, O10.011-O10.013, O10.019, O10.111-O10.113, O10.119, O10.211-O10.213, O10.219, O10.311-O10.313, O10.319, O10.411-O10.413, O10.419, O10.911-O10.913, O10.919, O11.1, O11.2, O11.3, O11.9, O12.00-O12.03, O12.10-O12.13, O12.20-O12.23, O13.1-O13.3, O13.9, O14.00, O14.02-O14.03, O14.10, O14.12-O14.13, O14.20, O14.22-O14.23, O14.90, O14.92-O14.93, O15.00, O15.02-O15.03, O15.1, O15.9, O16.1-O16.3, O16.9, O20.0, O20.8, O20.9, O21.0-O21.2, O21.8-O21.9, O22.00-O22.03, O22.20-O22.23, O22.30-O22.33, O22.40-O22.43, O22.50-O22.53, O22.8X1-O22.8X3, O22.8X9, O22.90-O22.93, O23.00-O23.03, O23.10-O23.13, O23.20-O23.23, O23.30-O23.33, O23.40-O23.43, O23.511-O23513, O23.519, O23.521-O23.523, O23.529, O23.591-O23.593, O23.599, O23.90-O23.93, O24.011-O24.013, O24.019, O24.111-O24.113, O24.119, O24.311-O24.313, O24.319, O24.410, O24.414-O24.415, O24.419, O24.811-O24.813, O24.819, O24.911-O24.913, O24.919, O25.10-O25.13, O26.00-O26.03,

Exclusions	Applicable Measures	Codes*
Pregnancy Diagnosis (continued)		O26.11-O26.13, O26.20-O26.23, O26.30-O26.33, O26.40-O26.43, O26.50-O26.53, O26.611-O26613, O26.619, O26.711-O26.713, O26.719, O26.811-O26.813, O26.819, O26.821-O26.823, O26.829, O26.831-O26.833, O26.839, O26.841-O26.843, O26.849, O26.851-O26.853, O26.859, O26.86, O26.872-O26.873, O26.879, O26.891-O26.893, O26.899, O26.90-O26.93, O28.0-O285, O28.8-O28.9, O29.011-O29.013, O29.019, O29.021-O29.23, O29.029, O29.091-O29.093, O29.099, O229.111-O29.113, O29.119, O29.121-O29.123, O29.129, O29.191-O29.193, O29.199, O29.211-O29.213, O29.219, O29.291-O29.293, O29.299, O29.3X1-O29.3X3, O29.3X9, O29.40-O29.43, O29.5X1-O29.5X3, O29.5X9, O29.60-O29.63, O29.8X1-O29.8X3, O29.8X9, O29.90-O29.93, O30.001-O30.003, O30.009, O30.011-O30.013, O30.019, O30.021-O30.023, O30.029, O30.031-O30.033, O30.039, O30.041-O30.043, O30.049, O30.091-O30.093, O30.099, O30.101-O30.103, O30.109, O30.111-O30.113, O30.119, O30.121-O30.123, O30.129, O30.131-O30.133, O30.139, O30.191-O30.193, O30.199, O30.201-O30.203, O30.209, O30.211-O30.213, O30.219, O30.221-O30.223, O30.229, O30.231-O30.233, O30.239, O30.291-O30.293, O30.299, O30.801-O30.803, O30.809, O30.811-O30.813, O30.819, O30.821-O30.823, O30.829, O30.831-O30.833, O30.839,

Exclusions	Applicable Measures	Codes*
Pregnancy Diagnosis (continued)		O30.891-O30.893, O30.899, O30.90-O30.93, O31.00X0-O30.00X5, O31.00X9, O31.01X0-O31.01X5, O31.01X9, O31.02X0-O31.02X5, O31.02X9, O31.03X0-O31.03X5, O31.03X9, O31.10X0-O31.10X5, O31.10X9, O31.11X0-O31.11X5, O31.11X9, O31.12X0-O31.12X5, O31.12X9, O31.13X0-O31.13X5, O31.13X9, O31.20X0-O31.20X5, O31.20X9, O31.21X0-O31.21X5, O31.21X9, O31.22X0-O31.22X5, O31.22X9, O31.23X0-O31.23X5, O31.23X9, O31.30X0-O31.30X5, O31.30X9, O31.31X0-O31.31X5, O31.31X9, O31X0-O31.32X5, O31.32X9, O33X0-O31.33X5, O31.33X9, O31.8X10-O32.8X15, O31.8X19, O31.8X20-O31.8X25, O31.8X29, O31.8X30-O31.8X35, O31.8X39, O31.8X90-O31.8X95, O31.8X99, O32.0XX0-O32.0XX5, O32.0XX9, O32.1XX0-O32.1XX5, O32.1XX9, O32.2XX0-O32.2XX5, O32.2XX9, O32.3XX0-O32.3XX5, O32.3XX9, O32.4XX0-O32.4XX5, O32.4XX9, O32.6XX0-O32.6XX5, O32.6XX9, O32.8XX0-O32.8XX5, O32.8XX9, O32.9XX0-O32.9XX5, O32.9XX9, O33.0-O33.2, O33.3XX0-O33.3XX5, O33.3XX9, O33.4XX0-O33.4XX5, O33.4XX9, O33.5XX0-O33.5XX5, O33.5XX9, O33.6XX0-O33.6XX5, O33.6XX9, O33.7, O33.7XX0-O33.7XX5, O33.7XX9, O33.8, O33.9, 34.00-O34.030, O34.10-O34.13, O34.21, O34.211-O34.212, O34.218-O34.219, O34.22, O34.29-O34.33, O34.40-O34.43, O34.511-O34.513, O34.519, O34.521-O34.523, O34.529, O34.531-O34.-533, O34.539, O34.591-O34.593, O34.599, O34.60-O34.63, O34.70-O34.73,

Exclusions	Applicable Measures	Codes*
Pregnancy Diagnosis (continued)		O34.80-O34.83, O34.90-O34.93, O35.0XX0-O35.0XX5, O35.0XX9, O35.1XX0-O35.1XX5, O35.1XX9, O35.2XX0-O35.2XX5, O35.2XX9, O35.3XX0-O35.3XX5, O35.3XX9, O35.4XX0-O35.4XX5, O35.4XX9, O35.5XX0-O35.5XX5, O35.5XX9, O35.6XX0-O35.6XX5, O35.6XX9, O35.7XX0-O35.7XX5, O35.7XX9, O35.8XX0-O35.8XX5, O35.8XX9, O35.9XX0-O35.9XX5, O35.9XX9, O36.0110-O36.0115, O36.0119, O36.0120-O36.0125, O36.0129, O36.0130-O36.0135, O36.0139, O36.0190-O36.0195, O36.0199, O36.0910-O36.0915, O36.0919, O36.0920-O36.0925, O36.0929, O36.0930-O36.0935, O36.0939, O36.0990-O36.0995, O36.0999, O36.1110-O36.1115, O36.1119, O36.1120-O36.1125, O36.1129, O36.1130-O36.1135, O36.1139, O36.1190-O36.1195, O36.1199, O36.1910-O36.1915, O36.1919, O36.1920-O36.1925, O36.1929, O36.1930-O36.1935, O36.1939, O36.1990-O36.1995, O36.1999, O36.20X0-O36.20X5, O36.20X9, O36.21X0-O36.21X5, O36.21X9, O36.22X0-O36.22X5, O36.22X9, O36.23X0-O36.23X5, O36.23X9, O36.4XX0-O36.4XX5, O36.4XX9, O36.5110-O36.5115, O36.5119-O36.5125, O36.5129-O36.5135, O36.5139, O36.5190-O36.5195, O36.5199, O36.5910-O36.5915, O36.5919-O36.5925, O36.5929-O36.5935, O36.5939, O36.5990-O36.5995, O36.5999, O36.60X0-O36.60X5, O36.60X9, O36.61X0-O36.61X5, O36.61X9, O36.62X0-O36.621X5, O36.62X9, O36.63X0-O36.63X5, O36.63X9,

Exclusions	Applicable Measures	Codes*
Pregnancy Diagnosis (continued)		O36.70X0-O36.70X5, O36.70X9, O36.71X0-O36.71X5, O36.71X9, O36.72X0-O36.72X5, O36.72X9, O36.73X0-O36.73X5, O36.73X9, O36.80X0-O36.80X5, O36.80X9, O36.8120-O36.8125, O36.8129, O36.8130-O36.8135, O36.8139, O36.8190-O36.8195, O36.8199, O36.8210-O36.8215, O36.8219, O36.8220-O36.8225, O36.8229, O36.8230-O36.8235, O36.8239, O36.8290-O36.8295, O36.8299, O36.8310-O36.8315, O36.8319, O36.8320-O36.8325, O36.8329-O36.8335, O36.8339, O36.8390-O36.8395, O36.8399, O36.8910-O36.8915, O36.8919, O36.8920-O36.8925, O36.8929, O36.8930-O36.8935, O36.8939, O36.8990-O36.8995, O36.8999, O36.90X0-O36.90X5, O36.90X9, O36.91X0-O36.91X5, O36.91X9, O36.92X0-O36.92X5, O36.92X9, O36.93X0-O36.93X5, O36.93X9, O40.1XX0-O40.1XX5, O40.1XX9, O40.2XX0-O40.2XX5, O40.2XX9, O4.31XX0-O40.3XX5, O40.3XX9, O40.9XX0-O40.9XX5, O40.9XX9, O41.00X0-O41.00X5, O41.00X9, O41.01X0-O41.01X5, O41.01X9, O41.02X0-O41.02X5, O41.02X9, O41.03X0-O41.03X5, O41.03X9, O41.1010-O41.1015, O41.1019-O41.1025, O41.1029-O41.1035, O41.1039, O41.1090-O41.1095, O41.1099, O41.1210-O41.1215, O41.1219-O41.1225, O41.1229-O41.1235, O41.1239, O41.1290-O41.1295, O41.1299, O41.1410-O41.1415, O41.1419-O41.1425, O41.1429-O41.1435, O41.1439, O41.1490-O41.1495, O41.1499,

Exclusions	Applicable Measures	Codes*
Pregnancy Diagnosis (continued)		O41.8X10-O41.8X15, O41.8X19-O41.8X25, O41.8X29-O41.8X35, O41.8X39, O41.8X91-O41.8X95, O41.8X99, O41.90X0-O41.90X5, O41.90X9, O41.91X0-O41.91X5, O41.91X9, O41.92X0-O41.92X5, O41.92X9, O41.93X0-O41.93X5, O41.93X9, O42.00, O42.011-O42.013, O42.019, O42.02, O42.10, O42.111-O42.113, O42.119, O42.12, O42.011-O42.013, O42.90, O42.911-O42.913, O42.919, O42.92, O43.011-O43.013, O43.019, O43.021-O43.023, O43.029, O43.101-O43.103, O43.109, O43.111-O43.113, O43.119, O43.121-O43.123, O43.129, O43.191-O43.193, O43.199, O43.211-O43.213, O43.219, O43.221-O43.223, O43.229, O43.231-O43.233, O43.239, O43.811-O43.813, O43.819, O43.891-O43.893, O43.899, O43.90-O43.93, O44.00-O44.03, O44.10-O44.13, O44.20-O44.23, O44.30-O44.33, O44.40-O44.43, O44.50-O44.53, O45.001-O45.003, O45.009, O45.011-O45.013, O45.019, O45.021-O45.023, O45.029, O45.091-O45.093, O45.099, O45.8X1-O45.8X3, O45.8X9, O45.90-O45.93, O46.001-O46.003, O46.009, O46.011-O46.013, O46.019, O46.021-O46.023, O46.029, O46.091-O46.093, O46.099, O46.8X1-O46.8X3, O46.8X9, O46.90-O46.93, O47.00-O47.03, O47.1, O47.9, O48.0-O48.1, O60.00, O60.02-O60.03, O71.00-O71.02-O71.03, O71.1-O71.7, O71.81-O71.82, O71.89, O71.9, O88.011-O88.013, O88.019, O88.111-O88.113, O88.119, O88.211-O88.213, O88.219,

Exclusions	Applicable Measures	Codes*
Pregnancy Diagnosis (continued)		O88.311-O88.313, O88.319, O88.811-O88.813, O88.819, O91.011-O91.013, O91.019, O91.13, O91.211-O91.213, O91.219, O91.23, O92.011-O92.013, O92.019, O92.03, O92.111-O92.113, O92.119, O92.3-O92.6, O92.70, O92.79, O98.011-O98.013, O98.019, O98.111-O98.113, O98.119, O98.211-O98.213, O98.219, O98.311-O98.313, O98.319, O98.411-O98.413, O98.419, O98.511-O98.513, O98.519, O98.611-O98.613, O98.619, O98.711-O98.713, O98.719, O98.811-O98.813, O98.819, O98.911-O98.913, O98.919, O99.011-O99.013, O99.019, O99.111-O99.113, O99.119, O99.210-O99.213, O99.280-O99.283, O99.310-O99.313, O99.320-O99.323, O99.330-O99.333, O99.340-O99.343, O99.350-O99.353, O99.411-O99.413, O99.419, O99.511-O99.513, O99.519, O99.611-O99.613, O99.619, O99.711-O99.713, O99.719, O99.810, O99.820, O99.830, O99.840-O99.843, O99.891, O9A.111-O9A.113, O9A.119, O9A.211-O9A.213, O9A.219, O9A.311-O9A.313, O9A.319, O9A.411-O9A.413, O9A.419, O9A.511-O9A.513, O9A.519, Z03.71-Z03.75, Z03.79, Z32.01, Z34.00-Z34.03, Z34.80-Z34.83, Z34.90-Z34.93, Z36, Z36.0-Z36.5, Z36.81-Z36.89, Z36.8A, Z36.9
Dialysis Procedure	BPC-E, CBP, KED, SPC-E, SPD-E	CPT: 90935, 90937, 90945, 90947, 90997, 091999 HCPCS: G0257, S9339 ICD-10: 3E1M39Z, 5A1D00Z, 5A1D50Z, 5A1D70Z, 5A1D80Z, 5A1D90Z
Narcolepsy	ADD	ICD-10: G47.411, G47.419, G47.421, G47.429

Exclusions	Applicable Measures	Codes*
Contraindications to Childhood Vaccines		ICD-10-CM: B20, B97.35, C81.00, C81.01-C81.49, C81.70-C81.79, C81.90-C81.99, C82.00-C82.69, C82.80-C82.99, C83.00-C83.19, C83.30-C83.39, C83.50-C83.59, C83.70-C83.99, C84.00-C84.19, C84.40-C84.49, C84.60-C84.79, C84.7A, C84.90-C84.99, C84.A0-C84.A9, C84.Z0-C84.Z9, C85.10-C85.29, C85.80-C85.99, C86.0-C88.9, C90.00-C90.02, C90.10-C90.12, C90.20-C90.22, C90.30-C90.32, C91.00-C91.02, C91.10-C91.12, C91.30-C91.32, C91.40-C91.42, C91.50-C91.52, C91.60-C91.62, C91.90-C91.92, C91.A0-C91.A2, C91.Z0-C91.Z2, C92.00-C92.02, C92.10-C92.12, C92.20-C92.22, C92.30-C92.32, C92.40-C92.42, C92.50-C92.52, C92.60-C92.62, C92.90-C92.92, C92.A0-C92.A2, C92.Z0-C92.Z2, C93.00-C93.02, C93.10-C93.12, C93.30-C93.32, C93.90-C93.92, C93.Z0-C93.Z2, C94.00-C94.02, C94.20-C94.22, C94.30-C94.32, C94.80-C94.82, C95.00-C95.02, C95.10-C95.12, C95.90-C95.92, C96.0, C96.20-C96.22, C96.29, C96.4, C96.9, C96.A, C96.Z, D80.0-D80.9, D81.0, D81.1, D81.2, D81.4, D81.6, D81.7, D81.82, D81.89, D81.9, D82.0-D82.4, D82.8-D82.9, D83.0-D83.2, D83.8-D83.9, D84.0-D84.1, D84.8, D84.81, D84.821, D84.822, D84.89, D84.9, D89.3, D89.810, D89.811, D89.812, D89.813, D89.82, D89.831, D89.832, D89.833, D89.834, D89.835, D89.839, D89.89, D89.9, K56.1, Z21

*Codes listed are subject to change. Meridian recognizes that the circumstances around the services provided may not always directly support/match the codes. It is crucial that the medical record documentation describes the services rendered in order to support the medical necessity and use of these codes.

Point of Service Code Definitions



Code	Description
02	Telehealth Provided Other than in Patient's Home
03	School
05	Indian Health Service Free-standing Facility
07	Tribal 638 Free-standing Facility
09	Prison/Correctional Facility
10	Telehealth Provided in Patient's Home
11	Office
12	Home
13	Assisted Living Facility
14	Group Home
15	Mobile Unit
16	Temporary Lodging
17	Walk-in Retail Health Clinic
18	Place of Employment-Worksite
19	Off Campus-Outpatient Hospital
20	Urgent Care Facility
21	Inpatient Hospital
22	On Campus-Outpatient Hospital
31	Skilled Nursing Facility
32	Nursing Facility
33	Custodial Care Facility
49	Independent Clinic
50	Federally Qualified Health Center
51	Inpatient Psychiatric Facility
56	Psychiatric Residential Treatment Center
57	Non-residential Substance Abuse Treatment Facility
58	Non-residential Opioid Treatment Facility
71	Public Health Clinic
72	Rural Health Clinic