

# Clinical Policy: Ocrelizumab (Ocrevus)

Reference Number: IL.ERX.SPA.52 Effective Date: 06.01.21 Last Review Date: 05.21 Line of Business:Illinois Medicaid

Revision Log

## See Important Reminder at the end of this policy for important regulatory and legal information.

## Description

Ocrelizumab (Ocrevus<sup>®</sup>) is a CD20-directed cytolytic antibody.

### FDA Approved Indication(s)

Ocrevus is indicated for the treatment of:

- Relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults
- Primary progressive MS, in adults

#### **Policy/Criteria**

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

Health plan approved formularies should be reviewed for all coverage determinations. Requirements to use preferred alternative agents apply only when such requirements align with the health plan approved formulary.

It is the policy of health plans affiliated with Envolve Pharmacy Solutions<sup>™</sup> that Ocrevus is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

- A. Multiple Sclerosis (must meet all):
  - 1. Diagnosis of one of the following (a, b, c, or d):
    - a. Clinically isolated syndrome, and member is contraindicated to both, or has experienced clinically significant adverse effects to one, of the following at up to maximally indicated doses: an interferon-beta agent (*Betaseron<sup>®</sup> and Rebif<sup>®</sup> are preferred agents*), glatiramer (*Copaxone<sup>®</sup> 20 mg is preferred*);
    - Relapsing-remitting MS, and failure of the following at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced: dimethyl fumarate (*Tecfidera<sup>®</sup> brand is preferred*) and any of the following: an interferon-beta agent (*Betaseron and Rebif are preferred agents*) or glatiramer (*Copaxone 20 mg is preferred*);
      - \*Prior authorization is required for all disease modifying therapies for MS
    - c. Secondary progressive MS;
    - d. Primary progressive MS;
  - 2. Prescribed by or in consultation with a neurologist;
  - 3. Age  $\geq$  18 years;
  - 4. Ocrevus is not prescribed concurrently with other disease modifying therapies for MS (see *Appendix D*);
  - 5. Documentation of baseline number of relapses per year and expanded disability status scale (EDSS) score;
  - 6. At the time of request, member does not have active hepatitis B infection (positive results for hepatitis B surface antigen and anti-hepatitis B virus tests);
  - 7. Dose does not exceed the following:
    - a. Initial dose: 300 mg, followed by a second 300 mg dose 2 weeks later;
    - b. Maintenance dose: 600 mg every 6 months.

Approval duration: 6 months



## B. Other diagnoses/indications

1. Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

## II. Continued Therapy

- A. Multiple Sclerosis (must meet all):
  - 1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions or member has previously met initial approval criteria;
  - 2. Member meets one of the following (a or b):
    - a. If member has received < 1 year of total treatment: Member is responding positively to therapy;
    - b. If member has received ≥ 1 year of total treatment: Member meets one of the following (i, ii, iii, or iv):
      - i. Member has not had an increase in the number of relapses per year compared to baseline;
      - ii. Member has not had  $\geq$  2 new MRI-detected lesions;
      - iii. Member has not had an increase in EDSS score from baseline;
      - iv. Medical justification supports that member is responding positively to therapy;
  - 3. Ocrevus is not prescribed concurrently with other disease modifying therapies for MS (see *Appendix D*);

If request is for a dose increase, new dose does not exceed 600 mg every 6 months.
Approval duration: <u>first re-authorization</u>: 6 months; <u>second and subsequent re-authorizations</u>: 12 months

#### **B.** Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

 Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

#### III. Diagnoses/Indications for which coverage is NOT authorized:

**A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – ERX.PA.01 or evidence of coverage documents.

### IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key EDSS: expanded disability status scale FDA: Food and Drug Administration MS: multiple sclerosis

#### Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Rebif <sup>®</sup> (interferon beta-1a)	22 mcg or 44 mcg SC TIW	44 mcg TIW
Betaseron <sup>®</sup> (interferon beta-1b)	250 mcg SC QOD	250 mg QOD
glatiramer acetate (Copaxone®)	20 mg SC QD or 40 mg SC TIW	20 mg/day or 40 mg TIW
dimethyl fumarate (Tecfidera <sup>®</sup> )	120 mg PO BID for 7 days, followed by 240 mg PO BID	480 mg/day



Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): active hepatitis B virus infection; history of life-threatening infusion reaction to Ocrevus
- Boxed warning(s): none reported

Appendix D: General Information

Disease-modifying therapies for MS are: glatiramer acetate (Copaxone<sup>®</sup>, Glatopa<sup>®</sup>), interferon beta-1a (Avonex<sup>®</sup>, Rebif<sup>®</sup>), interferon beta-1b (Betaseron<sup>®</sup>, Extavia<sup>®</sup>), peginterferon beta-1a (Plegridy<sup>®</sup>), dimethyl fumarate (Tecfidera<sup>®</sup>), diroximel fumarate (Vumerity<sup>®</sup>), monomethyl fumarate (Bafiertam<sup>™</sup>), fingolimod (Gilenya<sup>®</sup>), teriflunomide (Aubagio<sup>®</sup>), alemtuzumab (Lemtrada<sup>®</sup>), mitoxantrone (Novantrone<sup>®</sup>), natalizumab (Tysabri<sup>®</sup>), ocrelizumab (Ocrevus<sup>®</sup>), cladribine (Mavenclad<sup>®</sup>), siponimod (Mayzent<sup>®</sup>), ozanimod (Zeposia<sup>®</sup>), and ofatumumab (Kesimpta<sup>®</sup>).

#### V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Relapsing and	Initial 300 mg IV infusion with a second 300 mg IV	600 mg/6 months
primary progressive	infusion two weeks later, followed by subsequent	
MS	doses of 600 mg via IV infusion every 6 months	

#### VI. Product Availability

Single-dose vial: 300 mg/10 mL

#### VII. References

- 1. Ocrevus Prescribing Information. South San Francisco, CA: Genentech, Inc; December 2020. Available at <u>www.ocrevus.com</u>. Accessed February 8, 2021.
- Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: diseasemodifying therapies for adults with multiple sclerosis: report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. Neurology. 2018; 90(17): 777-788. Full guideline available at: https://www.aan.com/Guidelines/home/GetGuidelineContent/904.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	04.19.21	05.21

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information.

This Clinical Policy is not intended to dictate to providers how to practice medicine, nor does it constitute a contract or guarantee regarding payment or results. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members.

This policy is the property of Envolve Pharmacy Solutions. Unauthorized copying, use, and distribution of this Policy or any information contained herein is strictly prohibited. By accessing this policy, you agree to

# CLINICAL POLICY Ocrelizumab



be bound by the foregoing terms and conditions, in addition to the Site Use Agreement for Health Plans associated with Envolve Pharmacy Solutions.

©2021 Envolve Pharmacy Solutions. All rights reserved. All materials are exclusively owned by Envolve Pharmacy Solutions and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Envolve Pharmacy Solutions. You may not alter or remove any trademark, copyright or other notice contained herein.