

Clinical Policy: Risankizumab-rzaa (Skyrizi)

Reference Number: IL.ERX.SPA.337

Effective Date: 06.01.21

Last Review Date: 05.21

Line of Business: Illinois Medicaid

[Revision Log](#)

See **Important Reminder** at the end of this policy for important regulatory and legal information.

Description

Risankizumab-rzaa (Skyrizi[™]) is an interleukin-23 (IL-23) blocker.

FDA Approved Indication(s)

Skyrizi is indicated for the treatment of moderate-to-severe plaque psoriasis (PsO) in adults who are candidates for systemic therapy or phototherapy.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

Health plan approved formularies should be reviewed for all coverage determinations. Requirements to use preferred alternative agents apply only when such requirements align with the health plan approved formulary.

It is the policy of health plans affiliated with Envolve Pharmacy Solutions[™] that Skyrizi is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Plaque Psoriasis (must meet all):

1. Diagnosis of moderate-to-severe PsO as evidenced by involvement of one of the following (a or b):
 - a. $\geq 3\%$ of total body surface area;
 - b. Hands, feet, scalp, face, or genital area;
2. Prescribed by or in consultation with a dermatologist or rheumatologist;
3. Age ≥ 18 years;
4. Failure of at least TWO of the following, each used for ≥ 3 consecutive months, unless contraindicated or clinically significant adverse effects are experienced: Enbrel[®], Humira[®], Cimzia[®];
**Prior authorization may be required for Enbrel, Humira, and Cimzia*
5. Member meets one of the following (a or b):
 - a. Failure of a ≥ 3 consecutive month trial of methotrexate (MTX) at up to maximally indicated doses;
 - b. Member has intolerance or contraindication to MTX (*see Appendix D*), and failure of a ≥ 3 consecutive month trial of cyclosporine at up to maximally indicated doses, unless clinically significant adverse effects are experienced or both are contraindicated;
6. Dose does not exceed 150 mg at weeks 0 and 4, then every 12 weeks thereafter.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy

A. Plaque Psoriasis (must meet all):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions or member has previously met initial approval criteria;

2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed 150 mg every 12 weeks.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via a health plan affiliated with Envolve Pharmacy Solutions and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to ERX.PA.01 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – ERX.PA.01 or evidence of coverage documents;

B. Combination use of biological disease-modifying antirheumatic drugs (bDMARDs), including any tumor necrosis factor (TNF) antagonists [Cimzia®, Enbrel®, Simponi®, Avsola™, Inflectra™, Remicade®, Renflexis™], interleukin agents [Arcalyst® (IL-1 blocker), Ilaris® (IL-1 blocker), Kineret® (IL-1RA), Actemra® (IL-6RA), Kevzara® (IL-6RA), Stelara® (IL-12/23 inhibitor), Cosentyx® (IL-17A inhibitor), Taltz® (IL-17A inhibitor), Siliq™ (IL-17RA), Ilumya™ (IL-23 inhibitor), Skyrizi™ (IL-23 inhibitor), Tremfya® (IL-23 inhibitor)], janus kinase inhibitors (JAKi) [Xeljanz®/Xeljanz® XR, Rinvoq™], anti-CD20 monoclonal antibodies [Rituxan®, Riabni™, Ruxience™, Truxima®, and Rituxan Hycela®], selective co-stimulation modulators [Orencia®], or integrin receptor antagonists [Entyvio®] because of the possibility of increased immunosuppression, neutropenia and increased risk of infection.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

IL-23: interleukin-23

MTX: methotrexate

PsO: plaque psoriasis

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria.

The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
cyclosporine (Sandimmune®, Neoral®)	PsO 2.5 – 4 mg/kg/day PO divided BID	4 mg/kg/day
methotrexate (Rheumatrex®)	PsO 10 – 25 mg/week PO or 2.5 mg PO Q12 hr for 3 doses/week	30 mg/week
Enbrel® (etanercept)	PsO <i>Adults:</i> <u>Initial dose:</u> 50 mg SC twice weekly for 3 months <u>Maintenance dose:</u> 50 mg SC once weekly <i>Pediatrics:</i> Weight < 63 kg: 0.8 mg/kg SC once weekly Weight ≥ 63 kg: 50 mg SC once weekly	50 mg/week
Humira® (adalimumab)	PsO <u>Initial dose:</u> 80 mg SC	40 mg every other week

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	<u>Maintenance dose:</u> 40 mg SC every other week starting one week after initial dose	
Cimzia® (certolizumab)	PsO 400 mg SC every other week. For some patients (with body weight ≤ 90 kg), a dose of 400 mg SC at 0, 2 and 4 weeks, followed by 200 mg SC every other week may be considered.	400 mg every other week

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

Appendix D: General Information

- Definition of failure of MTX or DMARDs
 - Child-bearing age is not considered a contraindication for use of MTX. Each drug has risks in pregnancy. An educated patient and family planning would allow use of MTX in patients who have no intention of immediate pregnancy.
 - Social use of alcohol is not considered a contraindication for use of MTX. MTX may only be contraindicated if patients choose to drink over 14 units of alcohol per week. However, excessive alcohol drinking can lead to worsening of the condition, so patients who are serious about clinical response to therapy should refrain from excessive alcohol consumption.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
PsO	150 mg (two 75 mg injections) SC at Week 0, Week 4 and every 12 weeks thereafter	150 mg every 12 weeks

VI. Product Availability

Single-dose prefilled syringe: 75 mg/0.83 mL

VII. References

1. Skyrizi Prescribing Information. North Chicago, IL: Abbvie Inc. March 2020. Available at: https://www.rxabbvie.com/pdf/skyrizi_pi.pdf. Accessed January 13, 2021.
2. Menter A, Gottlieb A, Feldman SR, et al. American Academy of Dermatology. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 1. Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2008;58:826-850.
3. Menter A, Gottlieb A, Feldman, SR, et al. American Academy of Dermatology. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. *J Am Acad Dermatol*. 2008; 58(5): 826-50.
4. Menter A, Korman NF, Elmetts CA, et al. American Academy of Dermatology. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4. Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol*. 2008; doi: 10.1016/j.jaad.2009.03.027
5. Menter A, Korman, NJ, Elmetts CA, et al. American Academy of Dermatology. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 3. Guidelines of care for the management and treatment of psoriasis with topical therapies. *J Am Acad Dermatol*. 2009;60:643-659.

6. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019;80:1029-72. doi:10.1016/j.aad.201811.057.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	04.22.21	05.21

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information.

This Clinical Policy is not intended to dictate to providers how to practice medicine, nor does it constitute a contract or guarantee regarding payment or results. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members.

This policy is the property of Envolve Pharmacy Solutions. Unauthorized copying, use, and distribution of this Policy or any information contained herein is strictly prohibited. By accessing this policy, you agree to be bound by the foregoing terms and conditions, in addition to the Site Use Agreement for Health Plans associated with Envolve Pharmacy Solutions.

©2021 Envolve Pharmacy Solutions. All rights reserved. All materials are exclusively owned by Envolve Pharmacy Solutions and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Envolve Pharmacy Solutions. You may not alter or remove any trademark, copyright or other notice contained herein.