

**Clinical Policy: Adalimumab (Humira), Adalimumab-atto (Amjevita), Adalimumab-adbm (Cyltezo), Adalimumab-bwwd (Hadlima), Adalimumab-adaz (Hyrimoz), Adalimumab-aacf (Idacio), Adalimumab-ryvk (Simlandi), Adalimumab-aaty (Yuflyma), Adalimumab-aqvh (Yusimry)**

Reference Number: MDN.CP.PHAR.242

Effective Date: 04.01.22

Last Review Date: 03.07.25

Line of Business: Meridian IL Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**Description**

Adalimumab (Humira<sup>®</sup>), adalimumab-atto (Amjevita<sup>™</sup>), adalimumab-adbm (Cyltezo<sup>™</sup>), adalimumab-bwwd (Hadlima<sup>™</sup>), adalimumab-adaz (Hyrimoz<sup>™</sup>), and adalimumab-aacf (Idacio<sup>®</sup>), adalimumab-ryvk (Simlandi<sup>®</sup>), adalimumab-aaty (Yuflyma<sup>®</sup>), and adalimumab-aqvh (Yusimry<sup>™</sup>) are tumor necrosis factor (TNF) blockers.

**FDA Approved Indication(s)**

Indications	Description	Humira	Abrilada, Amjevita, Cyltezo/adalimumab-adbm, Hadlima, Hulio/adalimumab-fkjp, Hyrimoz/adalimumab-adaz, Idacio, adalimumab-aacf./Simlandi/adalimumab-ryvk, Yuflyma/adalimumab-aaty, Yusimry
Rheumatoid arthritis (RA)	Reducing signs and symptoms, inducing major clinical response, inhibiting the progression of structural damage, and improving physical function in adult patients with moderately to severely active RA	X	X
Juvenile idiopathic arthritis (JIA)	Reducing signs and symptoms of moderately to severely active polyarticular JIA in patients 2 years of age and older	X	X
Psoriatic arthritis (PsA)	Reducing signs and symptoms, inhibiting the progression of structural damage, and improving physical function in adult patients with active PsA	X	X

Indications	Description	Humira	Abrilada, Amjevita, Cyltezo/adalimumab-adbm, Hadlima, Hulio/adalimumab-fkjp, Hyrimoz/adalimumab-adaz, Idacio, adalimumab-aacf,/Simlandi/adalimumab-ryvk, Yuflyma/adalimumab-aaty, Yusimry
Ankylosing spondylitis (AS)	Reducing signs and symptoms in adult patients with active AS	X	X
Crohn's disease (CD)	Treatment of moderately to severely active CD in adults and pediatric patients 6 years of age and older	X	X
Adult ulcerative colitis (UC)	Treatment of moderately to severely active ulcerative colitis in adult patients  <u>Limitation of use:</u> Effectiveness has not been established in patients who have lost response to or were intolerant to TNF blockers	X	X
Pediatric UC	Treatment of moderately to severely active UC in pediatric patients 5 years of age and older  <u>Limitation of use:</u> Effectiveness has not been established in patients who have lost response to or were intolerant to TNF blockers	X	–
Plaque psoriasis (PsO)	The treatment of adult patients with moderate to severe chronic plaque psoriasis who are candidates for systemic therapy or phototherapy, and when other systemic therapies are medically less appropriate	X	X
Pediatric hidradenitis suppurativa (HS)	The treatment of moderate to severe hidradenitis suppurativa in patients 12 years of age and older	X	–
Adult HS	The treatment of moderate to severe hidradenitis suppurativa in adult patients	X	X
Pediatric uveitis (UV)	The treatment of non-infectious intermediate, posterior and panuveitis in adults and pediatric patients 2 years of age and older	X	–
Adult UV	The treatment of non-infectious intermediate, posterior, and panuveitis in adult patients	X	X

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation® that Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, and Yusimry are **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Ankylosing Spondylitis (must meet all):

1. Diagnosis of AS;
2. Prescribed by or in consultation with a rheumatologist;
3. Age  $\geq$  18 years;
4. Request is for adalimumab-adbm or Simlandi®;
5. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
6. Failure of at least TWO NSAIDs at up to maximally indicated doses, each used for  $\geq$  4 weeks unless contraindicated or clinically significant adverse effects are experienced;
7. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
8. Dose does not exceed 40 mg every other week.

**Approval duration: 6 months**

##### B. Crohn's Disease (must meet all):

1. Diagnosis of CD;
2. Prescribed by or in consultation with a gastroenterologist;
3. Request is for adalimumab-adbm or Simlandi®;
4. Age  $\geq$  6 years;
5. Member meets one of the following (a or b):
  - a. Failure of a  $\geq$  3 consecutive month trial of at least ONE immunomodulator (e.g., azathioprine, 6-mercaptopurine [6-MP], MTX) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
  - b. Medical justification supports inability to use immunomodulators (*see Appendix E*);
6. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
7. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the

- following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
8. Dose does not exceed one of the following (a or b):
    - a. Adults: 160 mg on Day 1 and 80 mg on Day 15, followed by maintenance dose of 40 mg every other week starting Day 29;
    - b. Pediatrics (i or ii):
      - i. Weight 17 kg (37 lbs.) to < 40 kg (88 lbs.): 80 mg on Day 1 and 40 mg on Day 15, followed by maintenance dose of 20 mg every other week starting Day 29;
      - ii. Weight  $\geq$  40 kg (88 lbs): 160 mg on Day 1 and 80 mg on Day 15, followed by maintenance dose of 40 mg every other week starting Day 29.

**Approval duration: 6 months**

**C. Hidradenitis Suppurativa (must meet all):**

1. Diagnosis of HS;
2. Prescribed by or in consultation with a dermatologist, rheumatologist, or gastroenterologist;
3. Member meets one of the following (a or b):
  - a. For Humira: Age  $\geq$  12 years;
  - b. For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry: Age  $\geq$  18 years;
4. If member is  $\geq$  18 years and request is for Abrilada, Amjevita, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, , member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
5. Documentation of Hurley stage II or stage III (*see Appendix D*);
6. Failure of at least TWO of the following, each tried for  $\geq$  3 consecutive months from different therapeutic classes, at up to maximally indicated doses, unless clinically significant adverse effects are experienced or all are contraindicated:
  - a. Systemic antibiotic therapy (e.g., clindamycin, minocycline, doxycycline, rifampin);
  - b. Oral retinoids (e.g., isotretinoin);
  - c. Hormonal treatment (e.g., estrogen-containing combined oral contraceptives, spironolactone);
7. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
8. Dose does not exceed 160 mg on Day 1 and 80 mg on Day 15, followed by maintenance dose of 40 mg every week starting Day 29.

**Approval duration: 6 months**

**D. Plaque Psoriasis (must meet all):**

1. Diagnosis of moderate-to-severe PsO as evidenced by involvement of one of the following (a or b):
  - a.  $\geq 3\%$  of total body surface area;
  - b. Hands, feet, scalp, face, or genital area;
2. Prescribed by or in consultation with a dermatologist or rheumatologist;
3. Request is for adalimumab-adbm or Simlandi<sup>®</sup>;
4. Age  $\geq 18$  years;
5. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
5. Member meets one of the following (a,b, or c):
  - a. Failure of a  $\geq 3$  consecutive month trial of MTX at up to maximally indicated doses;
  - b. Member has intolerance or contraindication to MTX (*see Appendix D*), and failure of a  $\geq 3$  consecutive month trial of cyclosporine at up to maximally indicated doses, unless clinically significant adverse effects are experienced or both are contraindicated;
  - c. Member has intolerance or contraindication to MTX, cyclosporine, and failure of phototherapy, unless contraindicated or clinically significant adverse effects are experienced;
6. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
7. Dose does not exceed 80 mg initial dose, followed by maintenance dose of 40 mg every other week starting one week after initial dose.

**Approval duration: 6 months**

**E. Polyarticular Juvenile Idiopathic Arthritis (must meet all):**

1. Diagnosis of PJIA as evidenced by  $\geq 5$  joints with active arthritis;
2. Prescribed by or in consultation with a rheumatologist;
3. Request is for adalimumab-adbm or Simlandi<sup>®</sup>;
4. Age  $\geq 2$  years;
5. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
6. Documented baseline 10-joint clinical juvenile arthritis disease activity score (cJADAS-10) (*see Appendix J*);
8. Member meets one of the following (a, b, c, or d):
  - a. Failure of a  $\geq 3$  consecutive month trial of MTX at up to maximally indicated doses;
  - b. Member has intolerance or contraindication to MTX (*see Appendix D*), and failure of a  $\geq 3$  consecutive month trial of leflunomide or sulfasalazine at up to maximally indicated doses, unless clinically significant adverse effects are experienced or both are contraindicated;

- c. For sacroiliitis/axial spine involvement (i.e., spine, hip), failure of a ? 4 week trial of an NSAID at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
  - d. Documented presence of high disease activity as evidenced by a cJADAS-10 > 8.5 (*see Appendix J*);
9. Dose does not exceed one of the following (a, b, or c):
- a. Weight 10 kg (22 lbs) to <15 kg (33 lbs): 10 mg every other week;
  - b. Weight 15 kg (33 lbs) to < 30 kg (66 lbs): 20 mg every other week;
  - c. Weight ≥ 30 kg (66 lbs): 40 mg every other week.

**Approval duration: 6 months**

**F. Psoriatic Arthritis** (must meet all):

1. Diagnosis of PsA;
2. Prescribed by or in consultation with a dermatologist or rheumatologist;
3. Request is for adalimumab-adbm or adalimumab-ryvk (Simlandi®);
4. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
5. Age ≥ 18 years;
6. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
7. Dose does not exceed 40 mg every other week.

**Approval duration: 6 months**

**G. Rheumatoid Arthritis** (must meet all):

1. Diagnosis of RA per American College of Rheumatology (ACR) criteria (*see Appendix G*);
2. Prescribed by or in consultation with a rheumatologist;
3. Request is for adalimumab-adbm or adalimumab-ryvk (Simlandi®);
4. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
5. Age ≥ 18 years;
6. Member meets one of the following (a or b):
  - a. Failure of a ≥ 3 consecutive month trial of MTX at up to maximally indicated doses;
  - b. Member has intolerance or contraindication to MTX (*see Appendix D*), and failure of a ≥ 3 consecutive month trial of at least ONE conventional disease-modifying antirheumatic drug [DMARD] (e.g., sulfasalazine, leflunomide, hydroxychloroquine) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
7. Documentation of one of the following baseline assessment scores (a or b):
  - a. Clinical disease activity index (CDAI) score (*see Appendix H*);
  - b. Routine assessment of patient index data 3 (RAPID3) score (*see Appendix I*);

8. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
9. Dose does not exceed 40 mg every other week.

**Approval duration: 6 months**

**H. Ulcerative Colitis (must meet all):**

1. Diagnosis of UC;
2. Prescribed by or in consultation with a gastroenterologist;
3. Member meets one of the following (a or b):
  - a. For Humira: Age  $\geq$  5 years;
  - b. For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry: Age  $\geq$  18 years;
4. If member is  $\geq$  18 years and request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, , member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
5. Documentation of a Mayo Score  $\geq$  6 (*see Appendix F*);
6. Failure of an 8-week trial of systemic corticosteroids, unless contraindicated or clinically significant adverse effects are experienced;
7. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
8. Dose does not exceed one of the following (a, b, or c):
  - a. For adults: 160 mg on Day 1 and 80 mg on Day 15, followed by maintenance dose of 40 mg every other week starting Day 29.
  - b. For pediatric patients weighing more than 20 kg, but less than 40 kg: 80 mg on Day 1, 40 mg on Day 8 and Day 15, followed by maintenance doses of 40 mg every other week or 20 mg every week
  - c. For pediatric patients weighing more than 40 kg: 160 mg on Day 1 and 80 mg on Day 8 and 15, followed by maintenance doses of 80 mg every other week or 40 mg every week.

**Approval duration: 6 months**

**I. Uveitis (must meet all):**

1. Diagnosis of non-infectious intermediate, posterior or panuveitis;
2. Prescribed by or in consultation with an ophthalmologist or rheumatologist;
3. Member meets one of the following (a or b):
  - c. For Humira: Age  $\geq$  2 years;
  - d. For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry: Age  $\geq$  18 years;

4. If member is  $\geq 18$  years and request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwbd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
5. Failure of a  $\geq 2$  week trial of a systemic corticosteroid (e.g., prednisone) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
6. Failure of a trial of a non-biologic immunosuppressive therapy (e.g., azathioprine, methotrexate, mycophenolate mofetil, cyclosporine, tacrolimus, cyclophosphamide, chlorambucil) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
7. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
8. Dose does not exceed 80 mg initial dose, followed by maintenance dose of 40 mg every other week starting one week after initial dose.

**Approval duration: 6 months**

**J. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.PMN.53 for Medicaid

**II. Continued Therapy**

**A. Rheumatoid Arthritis** (must meet all):

1. Member meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. If request is for Abrilada, Amjevita, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwbd, member must use ALL of the following,

- unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-adbm;
3. Member is responding positively to therapy as evidenced by one of the following (a or b):
    - a. A decrease in CDAI (*see Appendix H*) or RAPID3 (*see Appendix I*) score from baseline;
    - b. Medical justification stating inability to conduct CDAI re-assessment, and submission of RAPID3 score associated with disease severity that is similar to initial CDAI assessment or improved;
  3. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
  4. If request is for a dose increase, new dose does not exceed one of the following (a or b):\*
    - a. 40 mg every other week;
    - b. 40 mg every week and both of the following (i and ii):
      - i. Documentation supports inadequate response to a  $\geq 3$  month trial of 40 mg every other week or member is not a candidate for concurrent methotrexate and Humira due to contraindications or intolerance;

**Approval duration: 12 months\***

*\*(If new dosing regimen, approve for 6 months)*

**B. All Other Indications in Section I (must meet all):**

1. Member meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member meets one of the following (a, b, or c):
  - a. For CD (both i and ii):
    - i. Age  $\geq 6$  years;
    - ii. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi, adalimumab-adbm\* and adalimumab-ryvk
  - b. For pJIA (both i and ii):
    - i. Age  $\geq 2$  years;
    - ii. If request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following, unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi and adalimumab-ryvk;
  - c. For PsA, AS, UC, PsO, HS, UV: If member is  $\geq 18$  years and request is for Abrilada, Amjevita, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, adalimumab-aacf, or adalimumab-bwwd, member must use ALL of the following,

- unless clinically significant adverse effects are experienced or all are contraindicated: Simlandi, adalimumab-adbm\* and adalimumab-ryvk
3. Member meets one of the following (a, b or c):
    - a. HS, at least a 25% reduction in inflammatory nodules and abscesses;
    - b. For pJIA, member is responding positively to therapy as evidenced by a decrease in cJADAS-10 from baseline (*see Appendix J*)
    - c. For all other indications: member is responding positively to therapy;
  4. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
  5. If request is for a dose increase, new dose does not exceed one of the following (a, b, or c):
    - a. PJIA, PsA, AS, CD, PsO, UV: 40 mg every other week;
    - b. HS: 40 mg every week;
    - c. For UC, one of the following (i or ii)
      - i. 40 mg every other week or 20 mg every week;
      - ii. 80 mg every other week or 40 mg every week, and member initiated Humira prior to 18 years of age.

**Approval duration: 12 months\***

*\*(If new dosing regimen, approve for 6 months)*

**C. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 for Medicaid or evidence of coverage documents;
- B. Combination use with biological disease-modifying antirheumatic drugs (bDMARDs) or potent immunosuppressants, including but not limited to any tumor necrosis factor (TNF) antagonists [e.g., Cimzia<sup>®</sup>, Enbrel<sup>®</sup>, Humira<sup>®</sup> and its biosimilars, Remicade<sup>®</sup> and its biosimilars (Avsola<sup>™</sup>, Inflectra<sup>™</sup>, Renflexis<sup>™</sup>, Zymfentra<sup>®</sup>), Simponi<sup>®</sup>], interleukin agents [e.g., Actemra<sup>®</sup> (IL-6RA), Arcalyst<sup>®</sup> (IL-1 blocker),

Bimzelx<sup>®</sup> (IL-17A and F antagonist), Cosentyx<sup>®</sup> (IL-17A inhibitor), Ilaris<sup>®</sup> (IL-1 blocker), Ilumya<sup>™</sup> (IL-23 inhibitor), Kevzara<sup>®</sup> (IL-6RA), Kineret<sup>®</sup> (IL-1RA), Omvoh<sup>™</sup> (IL-23 antagonist), Siliq<sup>™</sup> (IL-17RA), Skyrizi<sup>™</sup> (IL-23 inhibitor), Stelara<sup>®</sup> (IL-12/23 inhibitor), Taltz<sup>®</sup> (IL-17A inhibitor), Tofidence<sup>™</sup> (IL-6), Tremfya<sup>®</sup> (IL-23 inhibitor), Tyenne<sup>®</sup> (IL-6), Wezlana<sup>™</sup> (IL-12/23 inhibitor), Janus kinase inhibitors (JAKi) [e.g., Cibinqo<sup>™</sup>, Olumiant<sup>™</sup>, Rinvoq<sup>™</sup>, Xeljanz<sup>®</sup>/Xeljanz<sup>®</sup> XR,], anti-CD20 monoclonal antibodies [Rituxan<sup>®</sup> and its biosimilars (Riabni<sup>™</sup>, Ruxience<sup>™</sup>, Truxima<sup>®</sup>), Rituxan Hycela<sup>®</sup>], selective co-stimulation modulators [Orencia<sup>®</sup>], integrin receptor antagonists [Entyvio<sup>®</sup>], tyrosine kinase 2 inhibitors [Sotyktu<sup>™</sup>], and sphingosine 1-phosphate receptor modulator [Velsipity<sup>™</sup>] because of the additive immunosuppression, increased risk of neutropenia, as well as increased risk of serious infections.

#### IV. Appendices/General Information

##### Appendix A: Abbreviation/Acronym Key

6-MP: 6-mercaptopurine	HS: hidradenitis suppurative
AS: ankylosing spondylitis	MTX: methotrexate
CD: Crohn's disease	NSAIDs: nonsteroidal anti-inflammatory drugs
CDAI: clinical disease activity index	PJIA: polyarticular juvenile idiopathic arthritis
cJADAS: clinical juvenile arthritis disease activity score	PsA: psoriatic arthritis
DMARD: disease-modifying antirheumatic drug	PsO: psoriasis
FDA: Food and Drug Administration	RA: rheumatoid arthritis
GI: gastrointestinal	
RAPID3: routine assessment of patient index data 3	UC: ulcerative colitis
TNF: tumor necrosis factor	UV: uveitis

##### Appendix B: Therapeutic Alternatives

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
acitretin (Soriatane <sup>®</sup> )	<b>PsO, HS</b> 25 or 50 mg PO QD	50 mg/day
azathioprine (Azasan <sup>®</sup> , Imuran <sup>®</sup> )	<b>RA</b> 1 mg/kg/day PO QD or divided BID  <b>CD*, UV*</b> 1.5 – 2 mg/kg/day PO	2.5 mg/kg/day
chlorambucil (Leukeran <sup>®</sup> )	<b>UV*</b> 0.2 mg/kg PO QD, then taper to 0.1 mg/kg PO QD or less	0.2 mg/kg/day

clindamycin (Cleocin <sup>®</sup> ) + rifampin (Rifadin <sup>®</sup> )	<b>HS*</b> clindamycin 300 mg PO BID and rifampin 300 mg PO BID	clindamycin: 1,800 mg/day rifampin: 600 mg/day
corticosteroids	<b>CD*</b> prednisone 40 mg PO QD for 2 weeks or IV 50 – 100 mg Q6H for 1 week  budesonide (Entocort EC <sup>®</sup> ) 6 – 9 mg PO QD  <b>UV*</b> prednisone 5 – 60 mg/day PO in 1 – 4 divided doses	Various
Cuprimine <sup>®</sup> (d-penicillamine)	<b>RA*</b> Initial dose: 125 or 250 mg PO QD Maintenance dose: 500 – 750 mg/day PO QD	1,500 mg/day
cyclophosphamide (Cytoxan <sup>®</sup> )	<b>UV*</b> 1 – 2 mg/kg/day PO	N/A
cyclosporine (Sandimmune <sup>®</sup> , Neoral <sup>®</sup> )	<b>PsO</b> 2.5 mg/kg/day PO divided BID  <b>RA</b> 2.5 – 4 mg/kg/day PO divided BID	PsO, RA: 4 mg/kg/day  UV: 5 mg/kg/day

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	<b>UV*</b> 2.5 – 5 mg/kg/day PO in divided doses	
doxycycline (Acticlate <sup>®</sup> )	<b>HS*</b> 50 – 100 mg PO BID	300 mg/day
Hormonal agents (e.g., estrogen- containing combined oral contraceptives, spironolactone)	<b>HS</b> varies	varies
hydroxychloroquine (Plaquenil <sup>®</sup> )	<b>RA*</b> Initial dose: 400 – 600 mg/day PO QD Maintenance dose: 200 – 400 mg/day PO QD	600 mg/day

Isotretinoin (Absorica <sup>®</sup> , Amnesteem <sup>®</sup> , Claravis <sup>®</sup> , Myorisan <sup>®</sup> , Zenatane <sup>®</sup> )	<b>HS</b> varies	varies 1.6 to 2 mg/kg/day
leflunomide (Arava <sup>®</sup> )	<b>PJIA*</b> Weight < 20 kg: 10 mg every other day PO Weight 20 - 40 kg: 10 mg/day PO Weight > 40 kg: 20 mg/day PO  <b>RA</b> 100 mg PO QD for 3 days, then 20 mg PO QD	20 mg/day
6-mercaptopurine (Purixan <sup>®</sup> )	<b>CD*</b> 50 mg PO QD or 1 – 2 mg/kg/day PO	2 mg/kg/day
methotrexate (Rheumatrex <sup>®</sup> )	<b>CD*</b> 15 – 25 mg/week IM or SC  <b>PsO</b> 10 – 25 mg/week PO or 2.5 mg PO Q12 hr for 3 doses/week  <b>PJIA*</b> 10 – 20 mg/m <sup>2</sup> /week PO, SC, or IM  <b>RA</b> 7.5 mg/week PO, SC, or IM or 2.5 mg PO Q12 hr for 3 doses/week  <b>UV*</b>	30 mg/week

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	7.5 – 20 mg/week PO	
minocycline (Minocin <sup>®</sup> )	<b>HS*</b> 50 – 100 mg PO BID	200 mg/day
mycophenolate mofetil (Cellcept <sup>®</sup> )	<b>UV*</b> 500 – 1,000 mg PO BID	3 g/day
NSAIDs (e.g., indomethacin, ibuprofen, naproxen, celecoxib)	<b>AS</b> Varies	Varies
Pentasa <sup>®</sup> (mesalamine)	<b>CD</b> 1,000 mg PO QID	4 g/day
Ridaura <sup>®</sup> (auranofin)	<b>RA</b> 6 mg PO QD or 3 mg PO BID	9 mg/day (3 mg TID)
sulfasalazine (Azulfidine <sup>®</sup> )	<b>PJIA*</b> 30-50 mg/kg/day PO divided BID  <b>RA</b> 2 g/day PO in divided doses	PJIA: 2 g/day RA: 3 g/day UC: 4 g/day
tacrolimus (Prograf <sup>®</sup> )	<b>CD*</b> 0.27 mg/kg/day PO in divided doses or 0.15 – 0.29 mg/kg/day PO  <b>UV*</b> 0.1-0.15 mg/kg/day PO	N/A
Actemra <sup>®</sup> (tocilizumab)	<b>RA</b> IV: 4 mg/kg every 4 weeks followed by an increase to 8 mg/kg every 4 weeks based on clinical response  <b>SC:</b> Weight < 100 kg: 162 mg SC every other week, followed by an increase to every week based on clinical response Weight ? 100 kg: 162 mg SC every week	IV: 800 mg every 4 weeks SC: 162 mg every week
Enbrel <sup>®</sup> (etanercept)	<b>AS</b> 50 mg SC once weekly  <b>PJIA</b> Weight < 63 kg: 0.8 mg/kg SC once weekly Weight ? 63 kg: 50 mg SC once weekly	50 mg/week

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	<b>PsA, RA</b> 25 mg SC twice weekly or 50 mg SC once weekly	
Cimzia® (certolizumab)	<b>AS</b> Initial dose: 400 mg SC at 0, 2, and 4	400 mg every 4 weeks
	weeks	
	Maintenance dose: 200 mg SC every other week (or 400 mg SC every 4 weeks)	
Kevzara® (sarilumab)	<b>RA</b> 200 mg SC once every two weeks	200 mg/2 weeks
Otezla® (apremilast)	<b>PsA</b> Initial dose:	60 mg/day
	Day 1: 10 mg PO QAM	
	Day 2: 10 mg PO QAM and 10 mg PO QPM	
Day 3: 10 mg PO QAM and 20 mg PO QPM		
Day 4: 20 mg PO QAM and 20 mg PO QPM		
Day 5: 20 mg PO QAM and 30 mg PO QPM		
Day 6 and thereafter: 30 mg PO BID		
Maintenance dose:		
Day 6 and thereafter: 30 mg PO BID		
Taltz® (ixekizumab)	<b>AS, PsA</b> Initial dose: 160 mg (two 80 mg	80 mg every 4 weeks
	injections) SC at week 0	
	Maintenance dose:	
	80 mg SC every 4 weeks	
	<b>PsO</b> Initial dose:	
	160 mg (two 80 mg injections) SC at week 0, then 80 mg SC at weeks 2, 4, 6, 8, 10, and 12	
Maintenance dose:		
80 mg SC every 4 weeks		
Xeljanz® (tofacitinib)	<b>PsA, RA</b> 5 mg PO BID	<b>PsA, RA</b> 10 mg/day

Xeljanz XR <sup>®</sup> (tofacitinib extended-release)	<b>PsA, RA</b> 11 mg PO QD	<b>PsA,RA</b> 11 mg/day
---	-------------------------------	----------------------------

Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

\*Off-label

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): none reported
- Boxed warning(s):
  - Serious infections
  - Malignancy

#### Appendix D: General Information

- Definition of failure of MTX or DMARDs
  - Child-bearing age is not considered a contraindication for use of MTX. Each drug has risks in pregnancy. An educated patient and family planning would allow use of MTX in patients who have no intention of immediate pregnancy.
  - Social use of alcohol is not considered a contraindication for use of MTX. MTX may only be contraindicated if patients choose to drink over 14 units of alcohol per week. However, excessive alcohol drinking can lead to worsening of the condition, so patients who are serious about clinical response to therapy should refrain from excessive alcohol consumption.
- Examples of positive response to therapy may include, but are not limited to:
  - Reduction in joint pain/swelling/tenderness
  - Improvement in ESR/CRP levels
  - Improvements in activities of daily living
- Hidradenitis suppurativa:
  - HS is sometimes referred to as: "acne inversa, acne conglobata, apocrine acne, apocrinitis, Fox-den disease, hidradenitis axillaris, HS, pyoderma sinifica fistulans, Velpeau's disease, and Verneuil's disease."
  - In HS, Hurley stages are used to determine severity of disease. Hurley stage II indicates moderate disease, and is characterized by recurrent abscesses, with sinus tracts and scarring, presenting as single or multiple widely separated lesions. Hurley stage III indicates severe disease, and is characterized by diffuse or near-diffuse involvement presenting as multiple interconnected tracts and abscesses across an entire area.
- Ulcerative colitis: there is insufficient evidence to support the off-label weekly dosing of Humira for the treatment of moderate-to-severe UC. It is the position of Centene Corporation<sup>®</sup> that the off-label weekly dosing of Humira for the treatment of moderate-to-severe UC is investigational and not medically necessary at this time.
  - The evidence from the *post hoc* study of the Humira pivotal trial suggests further studies are needed to confirm the benefit of weekly Humira dosing for the treatment of UC in patients with inadequate or loss of therapeutic response to treatment with Humira every other week. No large, randomized or prospective studies have been published to support the efficacy of the higher frequency of dosing, while national and international treatment guidelines also do not strongly support dose escalation of

Humira for UC. The current market consensus is that weekly dosing of Humira is not medically necessary due to lack of evidence to support its benefit.

*Appendix E: Immunomodulator Medical Justification*

- The following may be considered for medical justification supporting inability to use an immunomodulator for Crohn’s disease:
  - Inability to induce short-term symptomatic remission with a 3-month trial of systemic glucocorticoids
  - High-risk factors for intestinal complications may include:
    - Initial extensive ileal, ileocolonic, or proximal GI involvement
    - Initial extensive perianal/severe rectal disease
    - Fistulizing disease (e.g., perianal, enterocutaneous, and rectovaginal fistulas)
    - Deep ulcerations
    - Penetrating, stricturing or stenosis disease and/or phenotype
    - Intestinal obstruction or abscess
  - High risk factors for postoperative recurrence may include:
    - Less than 10 years duration between time of diagnosis and surgery
    - Disease location in the ileum and colon
    - Perianal fistula
    - Prior history of surgical resection
    - Use of corticosteroids prior to surgery

*Appendix F: Mayo Score*

- Mayo Score: evaluates ulcerative colitis stage, based on four parameters: stool frequency, rectal bleeding, endoscopic evaluation and Physician’s global assessment. Each parameter of the score ranges from zero (normal or inactive disease) to 3 (severe activity) with an overall score of 12.

Score	Decoding
0 – 2	Remission
3 – 5	Mild activity
6 – 10	Moderate activity
>10	Severe activity

- The following may be considered for medical justification supporting inability to use an immunomodulator for ulcerative colitis:
  - Documentation of Mayo Score 6 – 12 indicative of moderate to severe ulcerative colitis.

*Appendix G: The 2010 ACR Classification Criteria for RA*

Add score of categories A through D; a score of  $\geq 6$  out of 10 is needed for classification of a patient as having definite RA.

A	Joint involvement	Score
	1 large joint	0
	2-10 large joints	1
	1-3 small joints (with or without involvement of large joints)	2
	4-10 small joints (with or without involvement of large joints)	3
	> 10 joints (at least one small joint)	5

<b>B Serology (at least one test result is needed for classification)</b>		
	Negative rheumatoid factor (RF) <i>and</i> negative anti-citrullinated protein antibody (ACPA)	0
	Low positive RF <i>or</i> low positive ACPA <i>* Low: &lt; 3 x upper limit of normal</i>	2
	High positive RF <i>or</i> high positive ACPA <i>* High: ≥ 3 x upper limit of normal</i>	3
<b>C Acute phase reactants (at least one test result is needed for classification)</b>		
	Normal C-reactive protein (CRP) and normal erythrocyte sedimentation rate (ESR)	0
	Abnormal CRP or abnormal ESR	1
<b>D Duration of symptoms</b>		
	< 6 weeks	0
	≥ 6 weeks	1

*Appendix H: Clinical Disease Activity Index (CDAI) Score*

The Clinical Disease Activity Index (CDAI) is a composite index for assessing disease activity in RA. CDAI is based on the simple summation of the count of swollen/tender joint count of 28 joints along with patient and physician global assessment on VAS (0–10 cm) Scale for estimating disease activity. The CDAI score ranges from 0 to 76.

<b>CDAI Score</b>	<b>Disease state interpretation</b>
< 2.8	Remission
> 2.8 to ≤ 10	Low disease activity
> 10 to ≤ 22	Moderate disease activity
> 22	High disease activity

*Appendix I: Routine Assessment of Patient Index Data 3 (RAPID3) Score*

The Routine Assessment of Patient Index Data 3 (RAPID3) is a pooled index of the three patient-reported ACR core data set measures: function, pain, and patient global estimate of status. Each of the individual measures is scored 0 – 10, and the maximum achievable score is 30.

<b>RAPID3 Score</b>	<b>Disease state interpretation</b>
< 3	Remission
3.1 to 6	Low disease activity
6.1 to 12	Moderate disease activity
> 12	High disease activity

*Appendix J: Clinical Juvenile Arthritis Disease Activity Score based on 10 joints (cJADAS-10)*

The cJADAS10 is a continuous disease activity score specific to JIA and consisting of the following three parameters totaling a maximum of 30 points:

- Physician’s global assessment of disease activity measured on a 0-10 visual analog scale (VAS), where 0 = no activity and 10 = maximum activity;
- Parent global assessment of well-being measured on a 0-10 VAS, where 0 = very well and 10 = very poor;

- Count of joints with active disease to a maximum count of 10 active joints\*  
\*ACR definition of active joint: presence of swelling (not due to currently inactive synovitis or to bony enlargement) or, if swelling is not present, limitation of motion accompanied by pain, tenderness, or both

cJADAS-10	Disease state interpretation
≤ 1	Inactive disease
1.1 to 2.5	Low disease activity
2.51 to 8.5	Moderate disease activity
> 8.5	High disease activity

## V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Adalimumab and biosimilars (Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry)	RA	40 mg SC every other week  Some patients with RA not receiving concomitant methotrexate may benefit from increasing the frequency to 40 mg every week or 80 mg every other week.	40 mg/week
	PJIA	<b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hyrimoz, Idacio:</b> Weight 10 kg (22 lbs) to < 15 kg (33 lbs): 10 mg SC every other week  <b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Idacio, Simlandi, Yuflyma:</b> Weight 15 kg (33 lbs) to < 30 kg (66 lbs): 20 mg SC every other week  <b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry:</b> Weight ≥ 30 kg (66 lbs): 40 mg SC every other week	40 mg every other week
	PsA	40 mg SC every other week	40 mg every other week
	AS		
	CD	<u>Initial dose:</u> <i>Adults:</i> 160 mg SC on Day 1, then 80 mg SC on Day 15  <i>Pediatrics:</i> <b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Idacio, Simlandi, Yuflyma:</b> Weight 17 kg (37 lbs) to < 40 kg (88 lbs): 80 mg SC on Day 1, then 40 mg SC on Day 15 <b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry:</b> Weight ≥ 40 kg (88 lbs): 160 mg SC on Day 1, then 80 mg SC on Day 15	40 mg every other week

Drug Name	Indication	Dosing Regimen	Maximum Dose
		<p><u>Maintenance dose:</u> <i>Adults:</i> 40 mg SC every other week starting on Day 29</p> <p><i>Pediatrics:</i> <b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Idacio, Simlandi, Yuflyma:</b> Weight 17 kg (37 lbs) to &lt; 40 kg (88 lbs): 20 mg SC every other week starting on Day 29 <b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry:</b> Weight ≥ 40 kg (88 lbs): 40 mg SC every other week starting on Day 29</p>	
	UC	<p><u>Initial dose:</u> <i>Adults:</i> 160 mg SC on Day 1, then 80 mg SC on Day 15</p> <p><u>Maintenance dose:</u> <i>Adults:</i> 40 mg SC every other week starting on Day 29</p>	40 mg every week
	PsO	<p><u>Initial dose:</u> 80 mg SC</p> <p><u>Maintenance dose:</u> 40 mg SC every other week starting one week after initial dose</p>	40 mg every other week
	HS	<p><b>Humira:</b> <i>For patients 12 years of age and older weighing at least 30 kg:</i> <u>Initial dose:</u> Weight 30 kg (66 lbs) to &lt; 60 kg (132 lbs): 80 mg SC on Day 1, then 40 mg on Day 8 Weight ≥ 60 kg (132 lbs): 160 mg SC on Day 1, then 80 mg SC on Day 15</p> <p><u>Maintenance dose:</u> Weight 30 kg (66 lbs) to &lt; 60 kg (132 lbs): 40 mg every other week Weight ≥ 60 kg (132 lbs): 40 mg SC every week or 80 mg SC every other week starting on Day 29</p> <p><b>Humira, Abrilada, Amjevita, Cyltezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry:</b> <u>Initial dose:</u> <i>Adults:</i> 160 mg SC on day 1, then 80 mg SC on Day 15</p> <p><u>Maintenance dose:</u> <i>Adults:</i> 40 mg SC every week or 80 mg SC every other week starting on Day 29</p>	40 mg/week

Drug Name	Indication	Dosing Regimen	Maximum Dose												
	UV	<p><b>Humira:</b> <i>Pediatrics:</i> Weight 10 kg (22 lbs) to &lt; 15 kg (33 lbs): 10 mg SC every other week Weight 15 kg (33 lbs) to &lt; 30 kg (66 lbs): 20 mg SC every other week Weight ≥ 30 kg (66 lbs): 40 mg SC every other week</p> <p><b>Humira, Abrilada, Amjevita, Cytezo, Hadlima, Hulio, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry:</b> <i>Adults:</i> Initial dose of 80 mg SC, followed by 40 mg SC every other week starting one week after the initial dose</p>	40 mg every other week												
Adalimumab (Humira)	Pediatric UC	<p><u>Initial dose:</u> <i>Pediatrics:</i></p> <table border="1"> <thead> <tr> <th>Weight</th> <th>Days 1 through 15</th> </tr> </thead> <tbody> <tr> <td>20 kg to less than 40 kg</td> <td>Day 1: 80 mg Day 8: 40 mg Day 15: 40 mg</td> </tr> <tr> <td>40 kg and greater</td> <td>Day 1: 160 mg (single dose or split over two consecutive days) Day 8: 80 mg Day 15: 80 mg</td> </tr> </tbody> </table> <p><i>Pediatrics:</i></p> <table border="1"> <thead> <tr> <th>Weight</th> <th>Starting on Day 29*</th> </tr> </thead> <tbody> <tr> <td>20 kg to less than 40 kg</td> <td>40 mg every other week or 20 mg every week</td> </tr> <tr> <td>40 kg and greater</td> <td>80 mg every other week or 40 mg every week</td> </tr> </tbody> </table> <p><i>*Continue the recommended pediatric dosage in patients who turn 18 years of age and who are well-controlled on Humira regimen.</i></p>	Weight	Days 1 through 15	20 kg to less than 40 kg	Day 1: 80 mg Day 8: 40 mg Day 15: 40 mg	40 kg and greater	Day 1: 160 mg (single dose or split over two consecutive days) Day 8: 80 mg Day 15: 80 mg	Weight	Starting on Day 29*	20 kg to less than 40 kg	40 mg every other week or 20 mg every week	40 kg and greater	80 mg every other week or 40 mg every week	80 mg every other week or 40 mg every week
Weight	Days 1 through 15														
20 kg to less than 40 kg	Day 1: 80 mg Day 8: 40 mg Day 15: 40 mg														
40 kg and greater	Day 1: 160 mg (single dose or split over two consecutive days) Day 8: 80 mg Day 15: 80 mg														
Weight	Starting on Day 29*														
20 kg to less than 40 kg	40 mg every other week or 20 mg every week														
40 kg and greater	80 mg every other week or 40 mg every week														

## VI. Product Availability

Drug Name	Availability
Adalimumab (Humira)	<ul style="list-style-type: none"> <li>• Single-dose prefilled pen: 80 mg/0.8 mL, 40 mg/0.8 mL, 40 mg/0.4 mL</li> <li>• Single-dose prefilled syringe: 80 mg/0.8 mL, 40 mg/0.8 mL, 40 mg/0.4 mL, 20 mg/0.4 mL, 20 mg/0.2 mL, 10 mg/0.2 mL, 10 mg/0.1 mL</li> <li>• Single-use vial for institutional use only: 40 mg/0.8 mL</li> </ul>
Adalimumab-afzb (Abrilada)	<ul style="list-style-type: none"> <li>• Single-dose prefilled pen (Abrilada Pen): 40 mg/0.8 mL</li> </ul>

Drug Name	Availability
	<ul style="list-style-type: none"> <li>• Single dose prefilled syringe: 40 mg/0.8 mL, 20 mg/0.4 mL, 10 mg/0.2 mL</li> <li>• Single-dose glass vial for institutional use only: 40 mg/0.8 mL</li> </ul>
Adalimumab-atto (Amjevita)	<ul style="list-style-type: none"> <li>• Single-dose prefilled SureClick autoinjector: 80 mg/0.8 mL, 40 mg/0.8 mL, 40 mg/0.4 mL</li> <li>• Single-dose prefilled syringe: 80 mg/0.8 mL, 40 mg/0.8 mL, 40 mg/0.4 mL, 20 mg/0.4 mL, 20 mg/0.2 mL, 10 mg/0.2 mL</li> </ul>
Adalimumab-adbm (Cyltezo)	<ul style="list-style-type: none"> <li>• Single-dose prefilled syringe: 40 mg/0.4 mL, 40 mg/0.8 mL, 20 mg/0.4 mL, 10 mg/0.2 mL</li> <li>• Single-dose prefilled pen (Cyltezo Pen): 40 mg/0.4 mL, 40 mg/0.8 mL</li> </ul>
Adalimumab-bwwd (Hadlima)	<ul style="list-style-type: none"> <li>• Single-dose prefilled autoinjector (Hadlima PushTouch): 40 mg/0.8 mL, 40 mg/0.4 mL (citrate-free)</li> <li>• Single-dose prefilled syringe: 40 mg/0.8 mL, 40 mg/0.4 mL (citrate-free)</li> <li>• Single-dose glass vial for institutional use only: 40 mg/0.8 mL</li> </ul>
Adalimumab-fkjp (Hulio)	<ul style="list-style-type: none"> <li>• Single-dose prefilled pen (Hulio Pen): 40 mg/0.8 mL</li> <li>• Single-dose prefilled syringe: 40 mg/0.8 mL, 20 mg/0.4 mL</li> </ul>
Adalimumab-adaz (Hyrimoz)	<ul style="list-style-type: none"> <li>• Single-dose prefilled glass syringe (with BD UltraSafe Passive™ Needle Guard): 20 mg/0.4 mL, 40 mg/0.8 mL, 40 mg/0.4 mL, 80 mg/0.8 mL</li> <li>• Single-dose prefilled pen (Sensoready® Pen): 40 mg/0.8 mL, 40 mg/0.4 mL, 80 mg/0.8 mL</li> <li>• Single-dose prefilled glass syringe: 10 mg/0.2 mL, 10 mg/0.1 mL, 20 mg/0.2 mL</li> </ul>
Adalimumab-aacf (Idacio)	<ul style="list-style-type: none"> <li>• Single-dose prefilled pen (Idacio Pen): 40 mg/0.8 mL</li> <li>• Single-dose prefilled glass syringe: 40 mg/0.8 mL</li> <li>• Single-dose institutional use vial kit: 40 mg/0.8 mL</li> </ul>
Adalimumab-ryvk (Simlandi)	<ul style="list-style-type: none"> <li>• Single-dose autoinjector: 40 mg/0.4 mL</li> <li>• Single-dose prefilled glass syringe: 20 mg/0.2 mL, 40 mg/0.4 mL, 80 mg/0.8 mL</li> </ul>
Adalimumab-aaty (Yuflyma)	<ul style="list-style-type: none"> <li>• Single-dose prefilled auto-injector (Yuflyma AI): 40 mg/0.4 mL, 80 mg/0.8 mL</li> <li>• Single-dose prefilled syringe with safety guard: 40 mg/0.4 mL, 80 mg/0.8 mL</li> <li>• Single-dose prefilled syringe: 20 mg/0.2 mL, 40 mg/0.4 mL, 80 mg/0.8 mL</li> </ul>
Adalimumab-aqvh (Yusimry)	<ul style="list-style-type: none"> <li>• Single-dose prefilled pen (Yusimry Pen): 40 mg/0.8 mL</li> <li>• Single-dose prefilled glass syringe: 40 mg/0.8 mL</li> </ul>

## VII. References

1. Humira Prescribing Information. North Chicago, IL: AbbVie, Inc.; November 2023. Available at:

- [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/125057s423lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/125057s423lbl.pdf). Accessed February 8, 2024.
2. Abrilada Prescribing Information. New York, NY: Pfizer Inc.; October 2023. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761118s007lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761118s007lbl.pdf). Accessed February 8, 2024.
  3. Amjevita Prescribing Information. Thousand Oaks, CA: Amgen Inc.; August 2023. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761024s015lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761024s015lbl.pdf). Accessed February 8, 2024.
  4. Cyltezo Prescribing Information. Ridgefield, CT: Boehringer Ingelheim Pharmaceuticals, Inc.; April 2024. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/761058s026lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/761058s026lbl.pdf). Accessed May 13, 2024.
  5. Hadlima Prescribing Information. Jersey City, NJ: Organon & Co.; July 2023. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761059s008lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761059s008lbl.pdf). Accessed February 8, 2024.
  6. Hulio Prescribing Information. Morgantown, WV: Myland Pharmaceuticals Inc.; August 2023. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761154s005lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761154s005lbl.pdf). Accessed February 8, 2024.
  7. Hyrimoz Prescribing Information. Princeton, NJ: Sandoz Inc.; September 2023. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761071s016lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761071s016lbl.pdf). Accessed February 8, 2024.
  8. Idacio Prescribing Information. Lake Zurich, IL. Fresenius Kabi.; January 2024. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/761255Orig2s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/761255Orig2s000lbl.pdf). Accessed January 24, 2024.
  9. Simlandi Prescribing Information. Parsippany, NJ. Teva Pharmaceuticals.; June 2024. Available at: <https://www.simlandihcp.com/globalassets/simlandi/prescribing-information.pdf>. Accessed August 13, 2024.
  10. Yuflyma Prescribing Information. Incheon, Republic of Korea. Celltrion, Inc.; December 2023. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761219s006s007s008lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761219s006s007s008lbl.pdf). Accessed February 8, 2024.
  11. Yusimry Prescribing Information. Redwood City, CA. Coherus BioSciences, Inc.; September 2023. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761216s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761216s004lbl.pdf). Accessed February 8, 2024.

### ***Rheumatoid Arthritis***

8. Aletaha D, Neogi T, Silman AJ, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum.* 2010; 62(9):2569-2581. DOI 10.1002/art.27584

9. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care Res.* 2011; 63(4):465-482.

10. Singh JA, Furst DE, Bharat A, et al. 2012 update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care Res.* 2012; 64(5):625-639.

11. Ringold, S, Weiss PF, et al. 2013 Update of the 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis. *Arthritis Care Res.* 2013; 65(10):2499-2512.

12. Smolen JS, Landewé R, Breedveld FC, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2013 update. *Ann Rheum Dis.* 2014; 73:492-509.

13. Dhaon P, Das SK, Srivastava R, et al. Performances of clinical disease activity index (CDAI) and simplified disease activity index (SDAI) appear to be better than the gold standard disease assessment score (DAS-28-CRP) to assess rheumatoid arthritis patients. *Int J Rheum Dis.* 2018; 21:1933-1939.

#### ***Psoriasis and Psoriatic Arthritis***

14. Menter A, Gottlieb A, Feldman SR, et al. Guidelines for the management of psoriasis and psoriatic arthritis. Section 1: Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2008; 58(5):826-850.

15. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011; 65(1):137-174.

16. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis.* 2015;0:1-12. doi:10.1136/annrheumdis-2015-208337

17. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80:102972. doi:10.1016/j.aad.201811.057.

18. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the treatment of psoriatic arthritis. *American College of Rheumatology.* 2019; 71(1):5-32. doi: 10.1002/art.40726

#### ***Spondylitis***

19. Braun J, van den berg R, et al. 2010 Update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Am Rheu Dis.* 2011; 70:896-904.

20. Ward MM, Deodhar A, Aki EA, et al. American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network 2015 recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Rheumatol.* 2015. DOI 10.1002/ART.39298.

#### ***Crohn's Disease and Ulcerative Colitis***

21. Lichtenstein GR, Loftus Jr. EV, Isaacs KI, Regueiro MD, Gerson LB, and Sands BE. ACG clinical guideline: management of Crohn’s disease in adults. *Am J Gastroenterol.* 2018; 113:481-517.

22. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. ACG Clinical Guideline: Ulcerative Colitis

in Adults. *Am J Gastroenterol.* 2019;114:384-413

**Hidradenitis Suppurativa**

23. Zouboulis CC, Desai N, Emtestam L, et al. European S1 guideline for the treatment of hidradenitis suppurativa/acne inversa. *J Eur Acad Dermatol Venereol.* April 2015; 29(4):61944. Epub 2015 Jan 30.

24. Gulliver W, Zouboulis CC, Prens E, et al. Evidence-based approach to the treatment of hidradenitis suppurativa/acne inversa, based on the European guidelines for hidradenitis suppurativa. *Rev Endocr Metab Disord.* February 1, 2016. doi: 10.1007/s11154-016-9328-5.

25. Zouboulis, CC. Adalimumab for the treatment of hidradenitis suppurativa/acne inversa. *Expert Review of Clinical Immunology.* August 29, 2016. doi: 10.1080/1744666X.2016.1221762.

26. Alikhan A, Sayed C, Alavi A, et al. North American Clinical Management Guidelines for Hidradenitis Suppurativa: a publication from the United States and Canadian Hidradenitis Suppurativa Foundations. Part II: topical, intralesional, and systemic medical management. *J Am Acad Dermatol.* 2019; pii: S0190-9622(19)30368-8. doi: 10.1016/j.jaad.2019.02.068.

**Uveitis**

27. Levy-Clarke G, Jabs DA, Read RW, et al. Expert panel recommendations for the use of antitumor necrosis factor biologic agents in patients with ocular inflammatory disorders. *Ophthalmology.* 2014; 121:785.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J0139	Injection, adalimumab, 1 mg
Q5140	Injection, adalimumab-fkjp, biosimilar, 1 mg
Q5141	Injection, adalimumab-aaty, biosimilar, 1 mg
Q5142	Injection, adalimumab-ryvk biosimilar, 1 mg
Q5143	Injection, adalimumab-adbm, biosimilar, 1 mg
Q5144	Injection, adalimumab-aacf (idacio), biosimilar, 1 mg
Q5145	Injection, adalimumab-afzb (abrilada), biosimilar, 1 mg
C9399	Unclassified drugs or biologicals
J3590	Unclassified biologics

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created, adapted from CP.PHAR.242	04.01.22	04.22
Policy retired- t/f criteria removed	12.22.22	
Policy readapted to align with HFS PDL due to February SDC adaptation of Amjevita; added biosimilar Idacio; added Humira as preferred product in Section I	02.20.23	
RT4: for Cyltezo, added new dosage form (single-dose prefilled pen 40 mg/0.8 mL) and single-dose prefilled syringe 10 mg/0.2 mL to policy; RT4: added Yuflyma biosimilar to policy.	6.26.23	
2Q 2024 annual review: RT4: for Yuflyma, added newly approved UV indication to criteria; added HCPCS codes [C9399] and [J3590]; added Bimzelx, Zymfentra, Omvoh, Sotyktu, and Velsipity to section III.B;	7.26.24	
1Q2025 Annual Review: Added adalimumab-aacf and adalimumab-bwwd to criteria; for HS, added adalimumab-aacf, adalimumab-adbm, adalimumab-bwwd, and adalimumab-ryvk to age ≥ 18 years criteria; for UC and UV, added adalimumab-aacf, adalimumab-bwwd, and adalimumab-ryvk to age ≥ 18 years criteria; updated section I for HFS PDL updates; Added HCPCS codes [J0139, Q5140, Q5141, Q5142, Q5143, Q5144, Q5145] and removed [J0135, Q5131, Q5132]; for Simlandi, added new prefilled syringe formulation and strengths [20 mg/0.2 mL, 40 mg/0.4 mL, 80 mg/0.8 mL]; for section V, added Simlandi pediatric dose for pJIA [15 kg to less than 30 kg: 20 mg every other week] and pediatric dose for CD [17 kg to less than 40 kg: 80 mg SC on Day 1, 40 mg SC on Day 15, then 20 mg SC every other week starting on Day 29]; references reviewed and updated.	3.12.25	
Criteria modified to allow for 90 day transition period for HFS PDL until 6/30/25	<u>3.20.25</u>	
Humira removed as preferred agent per State PDL effective 6.30.25	<u>3.27.25</u>	
Updated preferred biosimilar agents	<u>6.18.25</u>	

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health

plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene

Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.