

Clinical Policy: Isotretinoin (Absorica, Absorica LD, Amnesteem, Claravis, Zenatane)

Reference Number: CP.PMN.143

Effective Date: 12.01.14

Last Review Date: 11.25

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Isotretinoin (Absorica[®], Absorica LD[®], Amnesteem[®], Claravis[™], Zenatane[®]) is a systemic retinoid.

FDA Approved Indication(s)

Absorica, Absorica LD, Amnesteem, Claravis, and Zenatane are indicated for severe recalcitrant nodular acne. Absorica and Absorica LD are specifically indicated in non-pregnant patients 12 years of age and older with multiple inflammatory nodules with a diameter of 5 mm or greater.

Limitation(s) of use: If a second course of Absorica/Absorica LD therapy is needed, it is not recommended before a two-month waiting period because the patient's acne may continue to improve following a 15 to 20-week course of therapy.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results, or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that isotretinoin, Absorica, Absorica LD, Amnesteem, Claravis, and Zenatane are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Acne (must meet all):**

1. Diagnosis of nodular acne;
2. Age $\geq$ 12 years;
3. Failure of $\geq$ 2 of the following topical anti-acne agents[^] (must be from 2 different classes listed below), unless clinically significant adverse effects are experienced or all are contraindicated: *
 - a. Topical antibiotics: clindamycin, erythromycin;
 - b. Topical anti-infectives: benzoyl peroxide;
 - c. Topical retinoids: tretinoin;

[^]Prior authorization may be required for tretinoin and combination products
^{*}For Illinois HIM requests, the step therapy requirements above do not apply as of 1/1/2026 per IL HB 5395
4. At least one of the topical agents above was used concurrently with one of the following oral antibiotics for $\geq$ 60 days: doxycycline, erythromycin, minocycline,

tetracycline, unless clinically significant adverse effects are experienced or all are contraindicated;*

**For Illinois HIM requests, the step therapy requirements above do not apply as of 1/1/2026 per IL HB 5395*

5. If request is for Absorica or Absorica LD, member must use Amnesteem, Claravis, Zenatane, and generic isotretinoin unless clinically significant adverse effects are experienced or all are contraindicated;*

**For Illinois HIM requests, the step therapy requirements above do not apply as of 1/1/2026 per IL HB 5395*

6. Dose does not exceed one of the following (a or b):
 - a. Absorica, Amnesteem, Claravis, Zenatane: 2 mg/kg per day;
 - b. Absorica LD: 1.6 mg/kg per day.

Approval duration:

Medicaid/HIM – 6 months

Commercial – 12 months or duration of request, whichever is less

B. Hidradenitis Suppurativa (off-label) (must meet all):

1. Diagnosis of hidradenitis suppurativa (HS);
 2. Prescribed by or in consultation with a dermatologist, rheumatologist, or gastroenterologist;
 3. Age $\geq$ 12 years;
 4. Failure of one systemic antibiotic therapy (e.g., clindamycin, minocycline, doxycycline, rifampin), tried for $\geq$ 3 consecutive months at up to maximally indicated doses, unless clinically significant adverse effects are experienced or all are contraindicated;
 5. If request is for Absorica or Absorica LD, member must use Amnesteem, Claravis, Zenatane, and generic isotretinoin, unless clinically significant adverse effects are experienced or all are contraindicated;*
- *For Illinois HIM requests, the step therapy requirements above do not apply as of 1/1/2026 per IL HB 5395*
6. Request meets one of the following (a or b):
 - a. Dose does not exceed 1 mg/kg per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration:

Medicaid/HIM – 6 months

Commercial – 12 months or duration of request, whichever is less

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or

- b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
- 2. For acne: Both of the following (a and b):
 - a. Member is responding positively to therapy;
 - b. If member has received 20 consecutive weeks of treatment, an 8-week treatment-free interval must be allowed prior to reinitiating isotretinoin treatment;
- 3. For HS: At least a 25% reduction in inflammatory nodules and abscesses;
- 4. If request is for Absorica or Absorica LD, member must use Amnesteem, Claravis, Zenatane, and generic isotretinoin, unless clinically significant adverse effects are experienced or all are contraindicated;*

**For Illinois HIM requests, the step therapy requirements above do not apply as of 1/1/2026 per IL HB 5395*

- 5. If request is for a dose increase, request meets one of the following (a or b):
 - a. For acne: New dose does not exceed one of the following (i or ii):
 - i. Absorica, Amnesteem, Claravis, Zenatane: 2 mg/kg per day;
 - ii. Absorica LD: 1.6 mg/kg per day;
 - b. For HS: Request meets one of the following (i or ii):
 - i. New dose does not exceed 1 mg/kg per day;
 - ii. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration:

Medicaid/HIM – 6 months

Commercial – 12 months or duration of request, whichever is less

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:

- CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
- b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

HS: hidradenitis suppurativa

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Topical Agents		
clindamycin 1% (Cleocin T [®] , Clindagel [®] , Clindacin [®])	Acne Apply a thin film BID	Not applicable
erythromycin 2% (Erygel [®] , Ery [®])	Acne Apply to the affected area BID	Not applicable
benzoyl peroxide (Benzac [®]) BPO [®] , Panoxyl [®])	Acne Apply or wash QD to TID	Not applicable
tretinoin (Retin-A [®])	Acne Apply a thin film QD	Not applicable
Example combination products: - clindamycin/benzoyl peroxide - erythromycin/benzoyl peroxide - clindamycin/tretinoin	Acne Varies	Not applicable

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Oral Antibiotics		
doxycycline	Acne, HS* 50 to 100 mg PO QD	300 mg per day
clindamycin (Cleocin [®]) + rifampin (Rifadin [®])	HS* clindamycin 300 mg PO BID and rifampin 300 mg PO BID	clindamycin: 600 mg/day rifampin: 600 mg/day
erythromycin (EES [®] , Erythromycin Base [®] , Ery- Tab [®])	Acne 250 to 500 mg PO twice daily, followed by twice daily dosing	4 gm per day
minocycline	Acne IR: 100 mg PO BID ER: 1 mg/kg PO QD HS* 50 – 100 mg PO BID	200 mg per day
tetracycline	Acne 125 to 250 mg PO every 6 hours for 2 weeks, then 125 to 500 mg PO daily or every other day HS* 500 mg PO BID	1,000 mg per day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

*Off-label

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): pregnancy (category X), hypersensitivity to the medication or any of its components
- Boxed warning(s): if pregnancy occurs during isotretinoin use, there is an extremely high risk for severe birth defects (iPLEDGE REMS program enrollment is required for prescribers, patients, pharmacies, and distributors)

Appendix D: General Information

- The American Academy of Dermatology recognizes that acne patients with psychosocial burden or scarring should be considered as having severe acne and to be candidates for isotretinoin.
- Micromedex classifies the use of isotretinoin for the non-FDA labeled indication of rosacea as a Class IIa strength of recommendation.
- The American Acne and Rosacea Society Consensus Recommendations recognize that isotretinoin has been shown to be effective in treating some refractory cases of papulopustular rosacea, but therapeutic benefit may require continued use. Due to the

limited data on the management of refractory rosacea, isotretinoin should only be considered in select cases.

- Because of the risk of teratogenicity and to minimize fetal exposure, isotretinoin is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called iPLEDGE. Isotretinoin must only be prescribed by prescribers who are registered and activated with the iPLEDGE program. Isotretinoin must only be dispensed by a pharmacy registered and activated with iPLEDGE, and must only be dispensed to patients who are registered and meet all the requirements of iPLEDGE. Registered and activated pharmacies must receive isotretinoin only from wholesalers registered with iPLEDGE. For more information call 866-495-0654 or visit <http://www.ipledgeprogram.com>.
- HS is sometimes referred to as: acne inversa, acne conglobata, apocrine acne, apocrinitis, Fox-den disease, hidradenitis axillaris, pyoderma sinifica fistulans, Velpeau's disease, and Verneuil's disease.

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Isotretinoin (Absorica, Amnesteem, Claravis, Zenatane)	Acne	0.5 to 1 mg/kg/day PO given in two divided doses	2 mg/kg/day
	HS*	0.5 to 1 mg/kg/day PO given in two divided doses	Various
Isotretinoin (Absorica LD)	Acne	0.4 to 0.8 mg/kg/day PO given in two divided doses	1.6 mg/kg/day
	HS*	0.5 to 1 mg/kg/day PO given in two divided doses	Various

*Off-label use

VI. Product Availability

Drug Name	Availability
Isotretinoin (Absorica)	Capsules: 10 mg, 20 mg, 25 mg, 30 mg, 35 mg, and 40 mg
Isotretinoin (Absorica LD)	Capsules: 8 mg, 16 mg, 20 mg, 24 mg, 28 mg, and 32 mg
Isotretinoin (Amnesteem)	Capsules: 10 mg, 20 mg, 40 mg
Isotretinoin (Claravis, Zenatane)	Capsules: 10 mg, 20 mg, 30 mg, and 40 mg

VII. References

- Absorica and Absorica LD Prescribing Information. Cranbury, NJ: Sun Pharmaceutical Industries, Inc.; July 2023. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/021951s024,211913s0111bl.pdf. Accessed August 8, 2025.
- Amnesteem Prescribing Information. Morgantown, WV: Mylan Pharmaceuticals Inc; June 2024. Available at: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=b2cb63c9-f825-4991-9a2c-6260f1bbcc2c>. Accessed August 8, 2025.

3. Claravis Prescribing Information. North Wales, PA: Teva Pharmaceuticals USA, Inc.; August 2022. Available at: <https://dailymed.nlm.nih.gov/dailymed/lookup.cfm?setid=a31fd109-d0fd-4ab9-ba98-a3d64333c18d>. Accessed August 8, 2025.
4. Zenatane Prescribing Information. Princeton, NJ: Dr. Reddy's Laboratories Inc.; February 2025. Available at: <https://dailymed.nlm.nih.gov/dailymed/fda/fdaDrugXsl.cfm?setid=27b3cf26-f22e-5b70-1c24-009933b7c6ee>. Accessed August 8, 2025.
5. Clinical Pharmacology [database online]. Tampa, FL: Elsevier, Inc.; 2025. Available at: <https://www.clinicalkey.com/pharmacology/>. Accessed August 8, 2025.
6. Zaenglein AL, Pathy AL, Schlosser BJ, Alikhan A, Baldwin HE, Berson DS, et al. Guidelines of care for the management of acne vulgaris. *J Am Acad Dermatol*. 2016 Feb 15;74(5):945-973.e33. doi: 10.1016/j.jaad.2015.12.037.
7. Reynolds RV, Yeung H, Cheng CE, et al. Guidelines of care for the management of acne vulgaris. *J Am Acad Dermatol*. 2024 May;90(5):1006.e1-1006.e30. doi: 10.1016/j.jaad.2023.12.017.
8. Alikhan A, Sayed C, Alavi A, et al. North American Clinical Management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations. Part II: topical, intralesional, and systemic medical management. *J Am Acad Dermatol*. 2019; pii: S0190-9622(19)30368-8. Doi: 10.1016/j.jaad.2019.02.068.
9. Hendricks A, J, Hsiao J, L, Lowes M, et al: A comparison of international management guidelines for hidradenitis suppurativa. *Dermatology* 2021;237:81-96. doi: 10.1159/000503605.
10. Dagenet CB, Lee KH, Fragoso NM et al. Approach to the patient with hidradenitis suppurativa: Evaluating severity to guide therapy. *J Am Acad Dermatol*. 2024; 91:S22-6. Doi:10.1016/j.jaad.20024.09.007.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
4Q 2021 annual review: no significant changes; updated reference for HIM off-label use to HIM.PA.154 (replaces HIM.PHAR.21); references reviewed and updated.	08.09.21	11.21
Revised approval duration for Commercial line of business from length of benefit to 12 months or duration of request, whichever is less.	04.27.22	08.22
4Q 2022 annual review: no significant changes; converted prior trial language to “member must use” language; references reviewed and updated. Template changes applied to other diagnoses/indications and continued therapy section.	08.23.22	11.22
4Q 2023 annual review: no significant changes; updated FDA approved indications section to align with Absorica/ Absorica LD prescriber information; removed commercially unavailable brand therapeutic alternatives; references reviewed and updated.	08.02.23	11.23
In initial approval criteria, removed “for age $\geq$ 30 years” from asterisk note stating prior authorization may be required for tretinoin.	02.13.24	

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Added off-label criteria for hidradenitis suppurativa per local market request.	05.07.24	08.24
4Q 2024 annual review: in initial approval criteria, clarified that combination products may require prior authorization, removed redirection to specific concentrations for benzoyl peroxide and tretinoin, removed redirection to trimethoprim-sulfamethoxazole per acne guideline; updated Appendix B per Clinical Pharmacology; references reviewed and updated.	07.31.24	11.24
4Q 2025 annual review: removed obsolete brand Myorisan; for all indications, added step therapy bypass for IL HIM per IL HB 5395; references reviewed and updated.	08.08.25	11.25

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise

professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.